

Table S1. Comparison of mortality rate: no suicide attempt group vs suicide attempt group.

	Full cohort			<i>P</i>	PS matched cohort			<i>P</i>
	No suicide attempt (n = 2531)	Suicide attempt (n = 183)			No suicide attempt (n = 139)	Suicide attempt (n = 139)		
Outcome				< 0.001				0.002
Dead	362 (14.3)	69 (37.7)			29 (20.9)	52 (37.4)		
Alive	2169 (85.7)	114 (62.3)			110 (79.1)	87 (62.6)		

Data are expressed as n (%). PS, propensity score.

Table S2. Disposition location among survivors: no suicide attempt group vs suicide attempt group.

	Full cohort			OR (95% CI)	PS matched cohort			OR (95% CI)
	No suicide attempt (n = 2169) ^a	Suicide attempt (n = 114)			No suicide attempt (n = 110) ^a	Suicide attempt (n = 87)		
Long-term care facility	165 (7.6)	6 (5.3)	0.68 (0.29–1.56)	7 (6.4)	5 (5.7)	0.90 (0.28–2.93)		
Psychiatric hospital	3 (0.1)	9 (7.9)	61.9 (16.5–232.0)	1 (0.9)	8 (9.2)	11.0 (1.35–90.0)		

^aThe reference set was the no suicide attempt group. Data are expressed as n (%). PS, propensity score.




Figure S1. Distribution of propensity score in the full (A) and propensity score-matched (B) cohorts.