

Table S5. Change in urinalysis results (full analysis set)

Items	Group	n	At screening			n	6W				(method)
			mean	±	SD		mean	±	SD	p value ¹⁾	
Urine pH	Placebo	60	6.18	±	0.51	60	6.18	±	0.54	0.922	(Student's t-test)
	LJ88	60	6.24	±	0.65	59	6.17	±	0.51	0.533	
Urine specific gravity	Placebo	60	1.018	±	0.009	60	1.017	±	0.007	0.176	(Student's t-test)
	LJ88	60	1.019	±	0.009	59	1.019	±	0.009	0.793	
Urine protein (qualitative)*	Placebo	60	0.10	±	0.54	60	0.05	±	0.29	0.680	
	LJ88	60	0.05	±	0.22	59	0.05	±	0.22	1.000	
Urine glucose (qualitative)*	Placebo	60	0.00	±	0.00	60	0.00	±	0.00	1.000	
	LJ88	60	0.00	±	0.00	59	0.00	±	0.00	1.000	
Urine urobilinogen (qualitative)*	Placebo	60	0.00	±	0.00	60	0.00	±	0.00	1.000	(Wilcoxon signed-rank test)
	LJ88	60	0.00	±	0.00	59	0.00	±	0.00	1.000	
Urine bilirubin (qualitative)*	Placebo	60	0.00	±	0.00	60	0.00	±	0.00	1.000	(Wilcoxon signed-rank test)
	LJ88	60	0.00	±	0.00	59	0.00	±	0.00	1.000	
Urine keton body*	Placebo	60	0.00	±	0.00	60	0.00	±	0.00	1.000	
	LJ88	60	0.00	±	0.00	59	0.00	±	0.00	1.000	
Urinary occult blood reation*	Placebo	60	0.20	±	0.73	60	0.17	±	0.69	0.596	
	LJ88	60	0.00	±	0.00	59	0.29	±	0.87	0.016	

¹⁾Qualitative data were scored and used for analyses.