

Table S1

Primer and probe sequences used in RTqPCR.

Target	Primers (5' → 3')	Probe
<i>Cdh2</i>	For tgggaatccgacgaatgg Rev gcagatcggaccggatactg	6-FAM-tgaaagacccatccacgctgagcc-BHQ-1
<i>ATP5β</i>	For tactgtcgcgtgccattgct Rev cacgggcaacatcgtaatgc	6-FAM-atcccaacattgttggcagt-BHQ-1
<i>Adam10</i>	For aaacaccagcgtgccaaaag Rev ccctcttcattcgtagggtgaaa	
<i>Plin2</i>	For tgagatggcagagaacggtgtgaa Rev ttgcggctctagcttctggatgat	
<i>Rpl38</i>	For gttgctgcttgctgtgagt Rev cgaatttggcatcctttcgtc	
<i>Tspan5</i>	For acaagggctcctgaagtcagtt Rev tgatggaagagetgttgacaga	
<i>Tspan10</i>	For ctgctcaagtatctgatcttcc Rev aagccacgtaacagggagg	
<i>Tspan14</i>	For ggctctgcgggagaatatctg Rev gcactggtagctttctgaagg	
<i>Tspan15</i>	For acttcctgaacgacaacattcg Rev cgccacagcactgaactgattt	
<i>Tspan17</i>	For ctgctgcgggaaatacttctt Rev gatgttcgagagaacgccctt	
<i>Tspan33</i>	For ctacgatcggctaataagca Rev tgagcaggaacatgaggacac	