

Supporting Information

Amino group-driven adsorption of sodium p-perfluorous nonenoxybenzene sulfonate in water by the modified graphene oxide

Mengyuan Lu ^{1, #}, Yang Liu ^{1, #, *}, Xinning Zheng ², Wenjuan Liu ³, Yang Liu ⁴, Jia Bao ^{1, *}, Ao Feng ¹, Yueyao Bao ¹, Jiangyong Diao ⁵ and Hongyang Liu ⁵

¹ School of Environmental and Chemical Engineering, Shenyang University of Technology, Shenyang 110870, China;

² Shenyang Zhenxing Sewage Treatment Co., LTD, Shenyang 110143, China;

³ Dalian Xigang District Center for Disease Control and Prevention, Dalian 116021, China;

⁴ Shenyang Hoper Group CO., LTD, Shenyang 110112, China;

⁵ Shenyang National Laboratory for Materials Science, Institute of Metal Research, Chinese Academy of Sciences, Shenyang 110016, China

* Correspondence: liuyang@sut.edu.cn (Y.L.); baojia@sut.edu.cn (J.B.)

Both authors contributed equally to this paper.

Figure S1. Geometry optimization of GH-OBS solution (a), GO-OBS solution (b), and CS-GO-OBS solution (c)




Figure S2. MSD trajectories of three different solutions

