

Proceeding Paper

# Zn-Catalyzed Direct Synthesis of 3-Iodo-1,3-dienes from $\alpha$ -Allenols <sup>†</sup>

Mireia Toledano-Pinedo <sup>1,\*</sup>, Beatriz Peñín <sup>1</sup>, Teresa Martínez del Campo <sup>1</sup> and Pedro Almendros <sup>2</sup>

<sup>1</sup> Grupo de Lactamas y Heterociclos Bioactivos, Unidad Asociada al CSIC, Departamento de Química Orgánica, Facultad de Ciencias Químicas, Universidad Complutense de Madrid, 28040 Madrid, Spain; beapenin@ucm.es (B.P.); tmcampo@quim.ucm.es (T.M.d.C.)

<sup>2</sup> Instituto de Química Orgánica General, IQOG-CSIC, Juan de la Cierva 3, 28006 Madrid, Spain; palmendros@iqog.csic.es

\* Correspondence: mireiato@ucm.es

<sup>†</sup> Presented at the 24th International Electronic Conference on Synthetic Organic Chemistry, 15 November–15 December 2020; Available online: <https://ecsoc-24.sciforum.net/>.

**Abstract:** We wish to report herein a new protocol that allows one to obtain 3-iodo-1,3-dienes through a sustainable process starting from  $\alpha$ -allenols. In this new synthetic route, zinc (II) derivatives are used as metallic promoter, which are an inexpensive and environmentally friendly species.

**Keywords:** allenes; 3-iodo-1,3-dienes and zinc

**Citation:** Toledano-Pinedo, M.; Peñín, B.; del Campo, T.; Almendros, P. Zn-Catalyzed Direct Synthesis of 3-Iodo-1,3-dienes from  $\alpha$ -Allenols. *Chem. Proc.* **2021**, *3*, 113. <https://doi.org/10.3390/ecsoc-24-08091>

Academic Editors: Julio A. Seijas and M. Pilar Vázquez-Tato

Published: 13 November 2020

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

In recent years, the chemistry of allenes has attracted the attention of many chemists giving rise to numerous studies. The presence of a cumulated diene in the structure of the allene compound allows for the selective synthesis of different functionalized organic molecules [1–3].

Allenes are the starting point of a large number of synthetic routes, due to the possibility of having four substituents in its structure. Furthermore, both the electronic density and the reactivity of each carbon of the allenes can be modulated as a function of the substitution. Besides, the inherent axial chirality allows the stereoselective synthesis of optically active allenes and the transfer of chirality from the allenes to the final product.


On the other hand, iododienes are interesting species for organic chemistry because of their wide reactivity. These compounds are precursors in Diels–Alder reactions [4], palladium-catalyzed reactions [5] such as Stille [6], Negishi [7], and Sonogashira [8] and carbonylation cross-couplings [9,10], solvolysis [11,12], and metalation reactions [8].

In the literature there are several methods of synthesis of iododienes from allenes. Okamoto et al. described the oxidative addition of ( $\eta^2$ -propene)Ti(O-*i*-Pr)<sub>2</sub> to  $\alpha$ -allenyl carbonates to form titanium complexes that by reacting with molecular iodine resulted in 2-iodo-1,3-dienes (Scheme 1a) [13]. Nishimaya et al. observed the formation of 2-iodo-1,3-butadiene by treatment of 1-trimethylsilyl-2,3-butadiene with I<sub>2</sub> and tetra-*n*-butylammonium fluoride (TBAF) (Scheme 1b) [9].


Based on these precedents, the main aim of this project is the development of a new synthetic method that allows one to obtain 3-iodo-1,3-dienes directly from  $\alpha$ -allenols through a sustainable process.

First, the allenylation reaction was carried out on the aldehydes **1** (Scheme 2). Experiments were performed according to previous conditions developed in our research group [14,15]. Barbier-type conditions [16] were used where aldehydes **1** reacted with differently substituted propargyl bromides, using indium and a THF/NH<sub>4</sub>Cl mixture (1:5) as a metal promoter and solvent, respectively. In this way,  $\alpha$ -allenols **2** were prepared

with total regioselectivity.




**Scheme 1.** Previous reports on the synthesis of iododienes from allenes [9,13].



**Scheme 2.** Synthesis of  $\alpha$ -allenols [14,15].

Once the starting materials were synthesized, allenol **2a** was selected as model substrate. Initial experiments were performed though the reaction of **2a** with ZnI<sub>2</sub> and *p*-toluenesulfonic acid monohydrate in dichloromethane (Scheme 3), which provided the 3-iodo-1,3-diene **3a** as the only reaction product.



**Scheme 3.** Synthesis of 3-iodo-1,3-diene **3a**.


This reactivity involving addition of the iodine atom and double bond rearrangement was surprising, because it was expected to obtain the corresponding dihydrofuran through the well-developed metal-catalyzed oxycyclization of  $\alpha$ -allenols [17].

However, the yield was very low and the reaction conditions were optimized (see Table 1). The highest yield of the product was achieved by employing 1.2 equiv. of ZnI<sub>2</sub> at room temperature in dichloromethane without the presence of *p*-toluenesulfonic acid (Table 1, entry 4).

**Table 1.** Optimization of reaction conditions for the formation of 3-iodo-1,3-dienes **3**.


Entry	ZnI <sub>2</sub> (Equiv.)	TsOH·H <sub>2</sub> O (Equiv.)	Yield (%)
1	0.025	0.025	2
2	0.1	0.1	33
3	1.2	0.1	37
4	1.2	-	48

The scope of the reaction was explored in different  $\alpha$ -allenols giving rise to the corresponding 3-iodo-1,3-dienes **3a–h** in reasonable yields (22–69%) and good (*E*)-diastereoselectivity (Scheme 4). It should be noted that in products **3a** and **3b** the reaction temperature had to be increased to 80 °C to consume the starting material.



**Scheme 4.** Preparation of 3-iodo-1,3-dienes **3a–h**.

A possible pathway for the generation of 3-iodo-1,3-dienes **3** from  $\alpha$ -allenols **2** is outlined in Scheme 5. It is assumed that the initial step is the coordination of the zinc salt to the terminal double bond of the allene to give the corresponding complex **2-Zn**. Subsequently, the loss of HI occurs followed by an addition of an iodine atom. Final elimination of the hydroxyl group of the former  $\alpha$ -allenol assisted by Zn generated the iodo-diene.



**Scheme 5.** Possible mechanism for the formation of 3-iodo-1,3-dienes **3**.

## 2. Experimental Section

General methods:  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were recorded on a Bruker Avance-300 spectrometer. NMR spectra were recorded in  $\text{C}_6\text{D}_6$ , except otherwise stated. Chemical shifts are given in ppm relative to TMS ( $^1\text{H}$ , 0.00 ppm), or  $\text{C}_6\text{D}_6$  ( $^1\text{H}$ , 7.16 ppm;  $^{13}\text{C}$ , 128.0 ppm). Low and high resolution mass spectra were taken on an AGILENT 6520 Accurate-Mass QTOF LC/MS spectrometer using the electrospray mode (ES) unless otherwise stated. IR spectra were recorded on a Bruker Tensor 27 spectrometer. All commercially available compounds were used without further purification.


### 2.1. General Procedure for the Preparation of $\alpha$ -Allenols **2a–h**

These precursors were readily obtained as described in the literature [14,15].


### 2.2. General Procedure for the Preparation of 3-Iodo-1,3-dienes **3a–h**

$\text{ZnI}_2$  (1.2 mmol) was added to a well stirred solution of  $\alpha$ -allenol (1 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL). After the disappearance of the starting material (TLC) the mixture was filtered through a short pad of celite, washed with  $\text{CH}_2\text{Cl}_2$ , and then concentrated under reduced pressure to give the product.


Chromatography of the residue eluting with hexanes/ethyl acetate mixtures gave analytically pure compounds **3**. It should be noted that in products **3a** and **3b** the reaction temperature had to be increased to  $80^\circ\text{C}$  to consume the starting material.




**3-iodo-1,3-diene 3a**. From 40 mg (0.23 mmol) of allenol **2a**, and after chromatography of the residue using hexanes/ethyl acetate (60:1) as the eluent gave compound **3a** (31 mg, 48%) as a yellow oil;  $^1\text{H}$  NMR (300 MHz,  $\text{C}_6\text{D}_6$ ,  $25^\circ\text{C}$ ):  $\delta$  = 6.95 (s, 1H), 6.88 (d, 2H,  $J$  = 6.0 Hz), 6.78 (d, 2H,  $J$  = 6.1 Hz), 5.87 (d, 1H,  $J$  = 3.2 Hz), 5.70 (d, 1H,  $J$  = 3.0 Hz), 1.94 (s, 3H), 1.70 (s, 3H);  $^{13}\text{C}$  NMR (75 MHz  $\text{C}_6\text{D}_6$ ,  $25^\circ\text{C}$ ):  $\delta$  = 137.0, 136.5, 136.3 (CH), 134.8, 129.6 ( $2\text{CH}_{\text{Ar}}$ ), 129.2 ( $2\text{CH}_{\text{Ar}}$ ), 126.3 ( $\text{CH}_2$ ), 114.6, 21.1 ( $\text{CH}_3$ ), 15.6 ( $\text{CH}_3$ ); IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ):  $\nu$  = 3051, 2898, 553.




**3-iodo-1,3-diene 3b**. From 100 mg (0.53 mmol) of allenol **2b**, and after chromatography of the residue using hexanes/ethyl acetate (60:1) as the eluent gave compound **3b** (50 mg, 32%) as a yellow oil;  $^1\text{H}$  NMR (300 MHz,  $\text{C}_6\text{D}_6$ ,  $25^\circ\text{C}$ ):  $\delta$  = 7.08 (s, 1H), 7.04 (d, 2H,  $J$  = 8.4 Hz), 6.72 (d, 2H,  $J$  = 8.8 Hz), 6.04 (d, 1H,  $J$  = 1.5 Hz), 5.87 (d, 1H,  $J$  = 1.6 Hz), 3.28 (s, 3H), 1.87 (d, 3H,  $J$  = 1.1 Hz);  $^{13}\text{C}$  NMR (75 MHz,  $\text{C}_6\text{D}_6$ ,  $25^\circ\text{C}$ ):  $\delta$  = 159.3, 136.0 (CH), 135.6, 131.0 ( $2\text{CH}_{\text{Ar}}$ ), 130.0, 128.5 ( $2\text{CH}_{\text{Ar}}$ ), 126.0 ( $\text{CH}_2$ ), 114.9, 54.7 ( $\text{CH}_3$ ), 15.7 ( $\text{CH}_3$ ); IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ):  $\nu$  = 3079, 2854, 554.




**3-iodo-1,3-diene 3c**. From 100 mg (0.46 mmol) of allenol **2c**, and after chromatography of the residue using hexanes/ethyl acetate (13:1) as the eluent gave compound **3c** (62 mg, 41%) as a yellow oil;  $^1\text{H}$  NMR (300 MHz,  $\text{C}_6\text{D}_6$ ,  $25^\circ\text{C}$ ):  $\delta$  = 8.06 (m, 2H), 7.16 (s, 1H), 6.93 (m, 2H), 6.00 (d, 1H,  $J$  = 1.7 Hz), 5.85 (d, 1H,  $J$  = 1.7 Hz), 3.51 (s, 3H), 1.69 (d, 3H,  $J$  = 1.1 Hz);  $^{13}\text{C}$  NMR (75 MHz,  $\text{C}_6\text{D}_6$ ,  $25^\circ\text{C}$ ):  $\delta$  = 166.4, 142.0, 138.7, 135.1 (CH), 129.8 ( $2\text{CH}_{\text{Ar}}$ ), 129.4 ( $2\text{CH}_{\text{Ar}}$ ), 129.1, 127.7 ( $\text{CH}_2$ ), 113.6, 51.6 ( $\text{CH}_3$ ), 15.6 ( $\text{CH}_3$ ); IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ):  $\nu$  = 3001, 2854, 760, 548.




**3-iodo-1,3-diene 3d.** From 75 mg (0.39 mmol) of allenol **2d**, and after chromatography of the residue using hexanes/ethyl acetate (60:1) as the eluent gave compound **3d** (38 mg, 32%) as a yellow oil;  $^1\text{H}$  NMR (300 MHz,  $\text{C}_6\text{D}_6$ , 25  $^\circ\text{C}$ ):  $\delta$  7.03 (d, 2H,  $J$  = 8.5 Hz), 6.85 (s, 1H), 6.71 (d, 2H,  $J$  = 8.2 Hz), 5.99 (d, 1H,  $J$  = 1.6 Hz), 5.84 (d, 1H,  $J$  = 1.7 Hz), 1.67 (d, 3H,  $J$  = 1.2 Hz);  $^{13}\text{C}$  NMR (75 MHz,  $\text{C}_6\text{D}_6$ , 25  $^\circ\text{C}$ ):  $\delta$  = 137.6, 135.9, 134.9 (CH), 131.2, 130.8 (2 $\text{CH}_{\text{Ar}}$ ), 128.6 (2 $\text{CH}_{\text{Ar}}$ ), 126.9 ( $\text{CH}_2$ ), 113.8, 15.5 ( $\text{CH}_3$ ); IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ):  $\nu$  = 3047, 2937, 1095, 564.




**3-iodo-1,3-diene 3e.** From 50 mg (0.21 mmol) of allenol **2e**, and after chromatography of the residue using hexanes/ethyl acetate (16:1) as the eluent gave compound **3e** (32 mg, 44%) as a yellow oil;  $^1\text{H}$  NMR (300 MHz,  $\text{C}_6\text{D}_6$ , 25  $^\circ\text{C}$ ):  $\delta$  = 7.53 (d, 2H,  $J$  = 8.5 Hz), 7.50 (s, 1H), 6.98 (d, 2H,  $J$  = 8.2 Hz), 6.33 (d, 1H,  $J$  = 1.6 Hz), 6.18 (d, 1H,  $J$  = 1.7 Hz), 1.00 (d, 3H,  $J$  = 1.2 Hz);  $^{13}\text{C}$  NMR (75 MHz,  $\text{C}_6\text{D}_6$ , 25  $^\circ\text{C}$ ):  $\delta$  = 137.7, 136.3, 134.9 (CH), 131.5 (2 $\text{CH}_{\text{Ar}}$ ), 131.0 (2 $\text{CH}_{\text{Ar}}$ ), 127.0 ( $\text{CH}_2$ ), 121.5, 113.8, 15.4 ( $\text{CH}_3$ ); IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ):  $\nu$  = 3047, 2900, 1100, 528.



**3-iodo-1,3-diene 3f.** From 100 mg (0.51 mmol) of allenol **2f**, and after chromatography of the residue using hexanes as the eluent gave compound **3f** (35 mg, 22%) as a yellow oil;  $^1\text{H}$  NMR (300 MHz,  $\text{C}_6\text{D}_6$ , 25  $^\circ\text{C}$ ):  $\delta$  = 7.04 (m, 1H), 6.99 (ddd, 1H,  $J$  = 7.5, 2.6, 1.0 Hz), 6.77 (m, 3H), 5.95 (d, 1H,  $J$  = 1.6 Hz), 5.82 (d, 1H,  $J$  = 1.7 Hz), 1.64 (d, 3H,  $J$  = 1.1 Hz);  $^{13}\text{C}$  NMR (75 MHz,  $\text{C}_6\text{D}_6$ , 25  $^\circ\text{C}$ ):  $\delta$  = 139.4, 138.3, 134.6 (CH), 134.4, 129.6 (CH), 129.4 (CH), 127.5 (CH), 127.3 (CH), 127.1 ( $\text{CH}_2$ ), 113.5, 15.4 ( $\text{CH}_3$ ); IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ):  $\nu$  = 3052, 2958, 1073, 601.



**3-iodo-1,3-diene 3g.** From 100 mg (0.57 mmol) of allenol **2g**, and after chromatography of the residue using hexanes as the eluent gave compound **3g** (54 mg, 33%) as a yellow oil;  $^1\text{H}$  NMR (300 MHz,  $\text{C}_6\text{D}_6$ , 25  $^\circ\text{C}$ ):  $\delta$  = 7.09 (s, 1H), 7.00 (m, 4H), 6.03 (d, 1H,  $J$  = 1.5 Hz), 5.87 (d, 1H,  $J$  = 1.7 Hz), 2.07 (s, 3H), 1.71 (d, 3H,  $J$  = 1.1 Hz);  $^{13}\text{C}$  NMR (75 MHz,  $\text{C}_6\text{D}_6$ , 25  $^\circ\text{C}$ ):  $\delta$  = 136.9, 136.8, 136.7 (CH), 130.2 ( $\text{CH}_{\text{Ar}}$ ), 129.3 ( $\text{CH}_{\text{Ar}}$ ), 128.0, 127.7 ( $\text{CH}_{\text{Ar}}$ ), 126.5 ( $\text{CH}_2$ ), 125.7 ( $\text{CH}_{\text{Ar}}$ ), 114.4, 19.9 ( $\text{CH}_3$ ), 15.2 ( $\text{CH}_3$ ); IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ):  $\nu$  = 3039, 2906, 587.



**3-iodo-1,3-diene 3h.** From 50 mg (0.24 mmol) of allenol **2h**, and after chromatography of the residue using hexanes/ethyl acetate (12:1) as the eluent gave compound **3h** (53 mg, 69%) as a yellow oil;  $^1\text{H}$  NMR (300 MHz,  $\text{C}_6\text{D}_6$ , 25  $^\circ\text{C}$ ):  $\delta$  = 7.42 (s, 1H), 7.24 (m, 1H), 7.02 (m, 3H), 6.15 (m, 1H), 5.94 (d, 1H,  $J$  = 1.7 Hz), 1.99 (s, 3H), 1.81 (d, 3H,  $J$  = 1.1 Hz);  $^{13}\text{C}$  NMR (75 MHz,  $\text{C}_6\text{D}_6$ , 25  $^\circ\text{C}$ ):  $\delta$  = 139.5, 137.6, 136.2, 135.2 (CH), 129.7 ( $\text{CH}_{\text{Ar}}$ ), 128.1 ( $\text{CH}_{\text{Ar}}$ ), 126.7 ( $\text{CH}_2$ ), 125.4 ( $\text{CH}_{\text{Ar}}$ ), 124.4 ( $\text{CH}_{\text{Ar}}$ ), 113.7, 15.4 ( $\text{CH}_3$ ), 15.1 ( $\text{CH}_3$ ); IR ( $\text{CHCl}_3$ ,  $\text{cm}^{-1}$ ):  $\nu$  = 3100, 2252, 524.

### 3. Conclusions

In conclusion, we present the controlled preparation of 3-iodo-1,3-dienes, directly from allenes through the coordination of the zinc salt to the double terminal bond of the allene by a subtle variation of the substituents.

**Supplementary Materials:** The poster presentation is available online at <https://www.mdpi.com/article/10.3390/ecsoc-24-08091/s1>.

**Author Contributions:** M.T.-P. and B.P. planned and conducted experiments. M.T.-P. and B.P. analyzed the data for the compounds and compiled most of the Supplementary Information. T.M.d.C. analyzed data to support the mechanistic proposal. P.A. designed and directed the project. M.T.-P. wrote the manuscript. T.M.d.C. and P.A. contributed to discussion. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was supported in part by AEI (MICIU) and FEDER (Project PGC2018-095025-B-I00).

**Conflicts of Interest:** The authors declare no conflict of interest.

### References

- Schuster, H.F.; Coppola, G.M. *Allenes in Organic Synthesis*; John Wiley & Sons: New York, NY, USA, 1984.
- Patai, S. *The Chemistry of Ketenes, Allenes and Related Compounds Part 1*; John Wiley & Sons: New York, NY, USA, 1980.
- Alcaide, B.; Almendros, P. Progress in allene chemistry. *Chem. Soc. Rev.* **2014**, *43*, 2886–2887.
- Choi, H.; Shirley, H.J.; Aitken, H.R.; Schulte, T.; Shöhnel, T.; Hume, P.A.; Brimble, M.A.; Furkert, D.P. Intermolecular Diels–Alder Cycloaddition/Cross-Coupling Sequences of 2-Bromo-1, 3-butadienes. *Org. Lett.* **2020**, *22*, 1022–1027.
- Clayden, J.; Greeves, N.; Warren, S. *Organometallic Chemistry” Organic Chemistry*, 2nd ed.; Oxford University Press Inc.: New York, NY, USA; Oxford, UK, 2012.
- Coleman, R.S.; Walczak, M.C. Tandem Stille/Suzuki–Miyaura coupling of a hetero-bis-metalated diene. rapid, one-pot assembly of polyene systems. *Org. Lett.* **2005**, *7*, 2289–2291.
- Bradford, T.A.; Payne, A.D.; Willis, A.C.; Paddon-Row, M.N.; Sherburn, M.S. Cross-coupling for cross-conjugation: Practical synthesis and Diels–Alder reactions of [3] dendralenes. *Org. Lett.* **2007**, *9*, 4861–4864.
- Yu, L.; Meng, B.; Huang, X.J. Halohydroxylation of 1-cyclopropylallenes: An efficient and stereoselective method for the preparation of multisubstituted olefins. *Org. Chem.* **2008**, *73*, 6895–6898.
- Nishiyama, T.; Esumi, T.; Iwabuchi, Y.; Irie, H.; Hatakeyama, S. Preparation of 2-Iodo-1, 3-butadienes from 1-Trimethylsilyl-2, 3-butadienes and their Functionalizations. *Tetrahedron Lett.* **1998**, *39*, 43–46.
- Froese, J.; Reed, J.; Hudlicky, T. Palladium-catalyzed carbonylation of halo arene-cis-dihydrodiols to the corresponding carboxylates. Access to compounds unavailable by toluene dioxygenase-mediated dihydroxylation of the corresponding benzoate esters. *Org. Biomol. Chem.* **2014**, *12*, 7810–7819.
- Wade, L.C. *Química Orgánica*, 5th ed.; Pearson Educación, S.A.: Madrid, Spain, 2004.
- Grob, C.A.; Spaar, R. Mesomeric Vinyl Cations. Part II. Solvolysis of 2-bromo-1, 3-butadienes. *Helvetica Chim. Acta* **1970**, *53*, 2119–2129.
- Okamoto, S.; Sato, H.; Sato, F. Highly efficient synthesis of alka-1, 3-dien-2-yltitanium compounds from alka-2, 3-dienyl carbonates. A new, practical synthesis of 1, 3-dienes and 2-iodo-1, 3-dienes. *Tetrahedron Lett.* **1996**, *37*, 8865–8868.
- Alcaide, B.; Almendros, P.; Aragoncillo, C.; Redondo, M.C. Stereoselective Synthesis of 1,2,3-Trisubstituted 1,3-Dienes through Novel [3,3]-Sigmatropic Rearrangements in  $\alpha$ -Allenic Methanesulfonates: Application to the Preparation of Fused Tricyclic Systems by Tandem Rearrangement/Diels–Alder Reaction. *Eur. J. Org. Chem.* **2005**, 98–106, doi:10.1002/ejoc.200400527.
- Alcaide, B.; Almendros, P.; Rodríguez-Acebes, R. Efficient entry to diversely functionalized spirocyclic oxindoles from isatins through carbonyl-addition/cyclization reaction sequences. *J. Org. Chem.* **2006**, *71*, 2346–2351.
- Isaac, M.B.; Chan, T.-H. Indium-mediated coupling of aldehydes with prop-2-ynyl bromides in aqueous media. *J. Chem. Soc. Chem Commun.* **1995**, 1003, doi:10.1039/C39950001003.
- Chou, T.-H.; Yu, B.-H.; Chein, R.-J.  $ZnI_2/Zn(OTf)_2-TsOH$ : A versatile combined-acid system for catalytic intramolecular hydrofunctionalization and polyene cyclization. *Chem. Commun.* **2019**, *55*, 13522–13525.