

Proceeding Paper

Human Risks Due to *Acanthamoeba* spp. in Grass from Public Parks across Leicester City, England [†]

Umar Anjum ¹, Angela Magnet ², Fernando Izquierdo ² and Antonio Peña-Fernández ^{1,3,*}

¹ Leicester School of Allied Health Sciences, De Montfort University, Leicester LE1 9BH, UK

² Facultad de Farmacia, Universidad San Pablo-CEU, CEU Universities, Urbanización Montepríncipe, Boadilla del Monte, 28660 Madrid, Spain

³ Department of Surgery, Medical and Social Sciences, Faculty of Medicine and Health Sciences, University of Alcalá, Ctra. Madrid-Barcelona, Km. 33.600, 28871 Alcalá de Henares, Spain

* Correspondence: antonio.penafer@uah.es or antonio.pena-fernandez@dmu.ac.uk

[†] Presented at the 2nd International Electronic Conference on Microbiology, 1–15 December 2023; Available online: <https://ecm2023.sciforum.net/>.

Abstract: *Acanthamoeba* spp., an emerging opportunistic human parasite, has been previously detected in open-water systems across Leicester city and surrounding rural areas (Leicestershire, England), suggesting a potential risk for the population. A total of 190 grass samples were collected over a year (June 2017 to August 2018) to study the temporal variation, in and around Leicester city and adjacent rural areas. After appropriate pre-treatment to resuspend potential cysts/amoebas present in the grass avoiding potential soil contamination, DNA was extracted from each pre-concentrated sample using the Fast DNA[®] Spin kit. *Acanthamoeba* spp. were detected using a triplex real-time TaqMan PCR assay in 22 grass samples (11.57%). The seasonal study revealed the following prevalence percentages from summer 2017 to summer 2018: 15.7, 10.5, 13.1, 7.89, and 10.5%. To our knowledge, this is the first report reporting the presence and seasonal distribution of *Acanthamoeba* spp. in grass across Leicestershire.

Keywords: free-living amoebas; *Acanthamoeba* spp.; grass; public parks; Leicester; England



Citation: Anjum, U.; Magnet, A.; Izquierdo, F.; Peña-Fernández, A. Human Risks Due to *Acanthamoeba* spp. in Grass from Public Parks across Leicester City, England. *Biol. Life Sci. Forum* **2024**, *31*, 23. <https://doi.org/10.3390/ECM2023-16459>

Academic Editor: James White

Published: 30 November 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Acanthamoeba spp., an emerging opportunistic free-living heterotrophic protist that can affect humans [1], are the most predominant amoeba in diverse ecological habitats from an ample range of climatic regions [2]. *Acanthamoeba*'s life cycle consists of an active and feeding trophozoite and a dormant double-walled polygonal cyst, which are highly resistant to environmental harsh conditions and can remain viable for many years (for more than twenty years) [2,3]. As a result, *Acanthamoeba* has been found in a myriad of different environments, including in soils, sewage, domestic water supplies, ponds, seawater, and water reservoirs across the world. Moreover, Fatemi et al. [4] had recently identified the presence of the highly pathogenic genotypes of T4, T5, and T9 in garden cress, chives, mint, parsley, and basil collected from municipal public markets.

Owing to the increasing relevance of *Acanthamoeba* spp. as an emerging human concern in recent years, our group is carrying out a comprehensive environmental monitoring study to identify the presence and circulation of these amoebas in different environmental habitats across Leicester city and surrounding rural areas (Leicestershire, England). We initiated this study because of the reported high incidence of *Acanthamoeba* keratitis (AK) in the United Kingdom, which is about 15 times that recorded in the United States and 7 times that in the Netherlands [5]. These authors have related the high incidence with the presence of water storage tanks, which might promote colonisation of domestic water by these amoebas, although a preliminary study carried out by our group recently has not detected the presence of any human-pathogenic free-living amoebas in the tap water

in Leicester [6]. However, we have isolated *Acanthamoeba* spp. in different open water systems across Leicestershire, including the River Soar, which crosses Leicester city and attracts a large number of users [7]. Thus, the main aim was to determine the presence, distribution/circulation of this emerging opportunistic pathogen in grass from public parks and recreational areas in and around Leicester city and adjacent rural areas, and to identify potential risks to their citizens and users.

2. Material and Methods

No ethical approval was required for the described study. A total of 190 grass samples were collected over a year (June 2017 to August 2018), with, specifically, 38 samples collected during 5 seasons. Grass samples were collected by cutting the top portion (5–10 cm) to avoid contamination with the topsoils and stored in 50 mL Falcon tubes. Each sample was washed with 15–20 mL of PBS (1% sterile) during a 24 h agitation to obtain a homogenised supernatant, and any potential cyst or amoeba present was resuspended. Supernatants were collected in sterile 50 mL Falcon tubes and centrifuged at 3000 rpm for 30 min. Pellets were collected and suspended in 5 mL of sterile PBS (1%) for the detection of these amoebas.

DNA was extracted from appropriately pre-concentrated grass samples using the FastDNA[®] Spin Kit (MP Biomedicals, Solon, OH, USA), following the manufacturer protocol with the modifications described by Gomes et al. [1]. These modifications included the introduction of an additional 1/4 inch ceramic sphere in each Fastprep tube and the lysing cycles were performed in triplicate in the homogeniser FastPrep-24[™] 5G (MP Biomedicals, Solon, OH, USA). DNA materials were then purified with NucleoSpin[®] Gel and PCR Clean-up (Macherey-Nagel, Düren, Germany), following the manufacturers' instructions.

The triplex real-time TaqMan PCR assay described by Qvarnstrom et al. [8] was used to molecularly detect the presence of *Acanthamoeba* spp. in the collected grass samples. Positive controls were used as previously described by our group in Peña-Fernández et al. [9].

3. Results and Discussion

Acanthamoeba spp. were detected in 22 grass samples (11.57%). The seasonal study revealed the following prevalence percentages: 15.7 (summer 2017), 10.5 (autumn 2017), 13.1 (winter 2017–2018), 7.89 (spring 2018), and 10.5% (summer 2018). The highest frequency of *Acanthamoeba* spp. (15.7%; 8/38) was determined in summer 2017, meanwhile the lowest prevalence was observed in spring 2018 (7.9%; 3/38). The seasonal variations in the prevalence detected did not show statistical differences. To our knowledge, this is the first report reporting the presence and seasonal distribution of *Acanthamoeba* spp. in grass across different public parks and recreational green areas in Leicester city and surrounding rural areas in Leicestershire. The moderate to high seasonal prevalence found would be logical, owing to the high environmental resistance of their cysts [1–5]. Our results would suggest a potential unknown source for *Acanthamoeba* spp. in Leicester city/Leicestershire, which could represent a serious hazard to human health that should be appropriately tackled, to prevent and minimise the exposure to the users of these public and recreational areas in the East Midlands region explored.

Although data on the presence and/or prevalence of *Acanthamoeba* spp. and other free-living amoebas in grass are limited in the literature, different recent studies have reported their presence in a wide range of vegetables, including lettuce, carrots, cauliflower, radishes, onions, spinach, and tomatoes [4], which might explain their presence found in the grass.

The presence and seasonal circulation of *Acanthamoeba* spp. found in the grass could be explained by their presence in topsoils. However, and although our group has detected their presence in some topsoils sampled in a pilot study carried out on different public parks in Leicester, further monitoring studies would be needed to fully understand the potential role of the (top)soils on the presence and distribution of *Acanthamoeba* spp. found in the grass samples. Moreover, although their interactions with domestic and wild animals

are not well-understood, a potential zoonotic reservoir might explain the moderate to high distribution found in a similar fashion as for other parasites. Thus, although limited, *Acanthamoeba* species have been found in domestic and wild animals [10].

4. Conclusions

This is the first report reporting the presence and seasonal distribution of *Acanthamoeba* spp. in grass collected from public parks/recreational areas in Leicester city and in different rural areas across Leicestershire, suggesting a moderate to high seasonal prevalence, which highlights a concern for public health. Leicestershire's citizens could be exposed when playing sports, or by contact of the cornea or tissue wounds with the grass. Further genotyping studies are required for a better characterisation of the distribution and circulation of *Acanthamoeba* spp. in Leicestershire, and to identify appropriate public health interventions and decontamination techniques to minimise the risks identified, especially to protect individuals with their immune systems compromised.

Author Contributions: Conceptualization, A.P.-F.; methodology, U.A., A.M., F.I., A.P.-F.; validation, U.A., A.P.-F.; formal analysis, U.A., A.P.-F.; investigation, U.A., A.M., F.I., A.P.-F.; resources, A.M., F.I., A.P.-F.; data curation, U.A., A.P.-F.; writing—original draft preparation, A.P.-F.; writing—review and editing, U.A., A.M., F.I., A.P.-F.; visualization, U.A., A.M., F.I., A.P.-F.; supervision, A.P.-F.; project administration, A.P.-F.; internal funding acquisition, A.P.-F. All authors have read and agreed to the published version of the manuscript.

Funding: This project has been also partially supported with other internal funds from the Leicester School of Allied Health Sciences, De Montfort University. This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author. The data are not publicly available due to further processing for a future submission as a manuscript.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Gomes, T.S.; Vaccaro, L.; Magnet, A.; Izquierdo, F.; Ollero, D.; Martínez-Fernández, C.; Mayo, L.; Moran, M.; Pozuelo, M.J.; Fenoy, S.; et al. Presence and interaction of free-living amoebae and amoeba-resisting bacteria in water from drinking water treatment plants. *Sci. Total. Environ.* **2020**, *719*, 137080. [[CrossRef](#)] [[PubMed](#)]
2. Rayamajhee, B.; Williams, N.L.; Siboni, N.; Rodgers, K.; Willcox, M.; Henriquez, F.L.; Seymour, J.R.; Potts, J.; Johnson, C.; Scanes, P.; et al. Identification and quantification of *Acanthamoeba* spp. within seawater at four coastal lagoons on the east coast of Australia. *Sci. Total. Environ.* **2023**, *901*, 165862. [[CrossRef](#)] [[PubMed](#)]
3. Sriram, R.; Shoff, M.; Booton, G.; Fuerst, P.; Visvesvara, G.S. Survival of *Acanthamoeba* Cysts after Desiccation for More than 20 Years. *J. Clin. Microbiol.* **2008**, *46*, 4045–4048. [[CrossRef](#)] [[PubMed](#)]
4. Fatemi, M.; Niyiyati, M.; Rouhani, S.; Karamati, S.A.; Mirjalali, H.; Karanis, P. Contamination of fresh vegetables in municipal stores with pathogenic *Acanthamoeba* genotypes; a public health concern. *Int. J. Environ. Heal. Res.* **2022**, *33*, 1010–1021. [[CrossRef](#)] [[PubMed](#)]
5. Kilvington, S.; Gray, T.; Dart, J.; Morlet, N.; Beeching, J.R.; Frazer, D.G.; Matheson, M. *Acanthamoeba* keratitis: The role of domestic tap water contamination in the United Kingdom. *Investig. Ophthalmology Vis. Sci.* **2004**, *45*, 165–169. [[CrossRef](#)]
6. Anjum, U.; Acosta, L.; Peña-Fernández, A. Determination of human-pathogenic free living amoebas in drinking water supplies in Leicester, UK. In Proceedings of the ISEE 2022: 34th Annual Conference of the International Society of Environmental Epidemiology, Athens, Greece, 18–21 September 2022; Volume 2022.
7. Anjum, U.; Peña Fernández, A. Annual presence and distribution of human-pathogenic *Acanthamoeba* spp. in River Soar, Leicester, UK. In Proceedings of the ISEE 2021: 33rd Annual Conference of the International Society of Environmental Epidemiology, virtual event, 23–26 August 2021; Volume 2021.
8. Qvarnstrom, Y.; Visvesvara, G.S.; Sriram, R.; da Silva, A.J. Multiplex real-time PCR assay for simultaneous detection of *Acanthamoeba* spp., *Balamuthia mandrillaris*, and *Naegleria fowleri*. *J. Clin. Microbiol.* **2006**, *44*, 3589–3595. [[CrossRef](#)]

9. Peña-Fernández, A.; Guetiya Wadoum, R.E.; Anjum, U. Human pathogenic free-living amoebas in faeces from cows and pigs from Bombali and Tonkolili Districts, Sierra Leone. *Biol. Life Sci. Forum* 2023, *submitted for publication*.
10. Schuster, F.L.; Visvesvara, G.S. Free-living amoebae as opportunistic and non-opportunistic pathogens of humans and animals. *Int. J. Parasitol.* **2004**, *34*, 1001–1027. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.