

Supplementary Material

Electrochemical Determination of 17- β -Estradiol Using a Glassy Carbon Electrode Modified with α - Fe_2O_3 Nanoparticles Supported on Carbon Nanotubes

Juliana Costa Rolim Galvão, Mayara da Silva Araujo, Maiyara Carolyne Prete, Vanildo Leão Neto, Luiz Henrique Dall'Antonia, Roberto Matos, Cesar Ricardo Texeira Tarley and Roberta Antigo Medeiros *

Department of Chemistry, State University of Londrina, Londrina 86057-970, PR, Brazil; juliana.costa.rolim@uel.br (J.C.R.G.); asmayara@uel.br (M.d.S.A.); mayprete@gmail.com (M.C.P.); vanildosln@gmail.com (V.L.N.); luizh@uel.br (L.H.D.); rmatos@uel.br (R.M.); tarley@uel.br (C.R.T.T.)

* Correspondence: ramedeiros@uel.br; Tel.: +55-43-33714811




Figure S1: Cyclic voltammograms obtained for 0.1 mmol L^{-1} of E2 using $\alpha\text{-Fe}_2\text{O}_3\text{-CNT/GCE}$ at different pHs (A) E_p versus pH plot (B). Supporting electrolyte: 0.1 mol L^{-1} KCl solution (10% v/v ethanol) in the presence of $0.1 \mu\text{mol L}^{-1}$ E2. Scan rate: 50 mV s^{-1} .