

Supplementary Materials

Antimicrobial Properties of Flavonoid Derivatives with Bromine, Chlorine, and Nitro Group Obtained by Chemical Synthesis and Biotransformation Studies

Martyna Perz^{1,*}, Daria Szymanowska^{2,3}, Tomasz Janeczko¹, and Edyta Kostrzewska-Susłow^{1,*}

¹ Department of Food Chemistry and Biocatalysis, Faculty of Biotechnology and Food Science, Wrocław University of Environmental and Life Sciences, 50-375 Wrocław, Poland;

² Department of Biotechnology and Food Microbiology, Faculty of Food Science and Nutrition, Poznań University of Life Sciences, 60-627 Poznań, Poland;

³ Department of Pharmacognosy and Biomaterials, Faculty of Pharmacy, University of Medical Sciences, Rokietnicka 3, 60-806 Poznań, Poland;

* Correspondence: martyna.perz@upwr.edu.pl (M.P.); edyta.kostrzewska-suslow@upwr.edu.pl (E.K.-S.)

Compound name: 5'-chloro-2'-hydroxy-3'-nitrochalcone

Molecular Formula: C₁₅H₁₀ClNO₄

Formula Weight: 303.697

Ionization mode: positive

Precursor [M + H]⁺: 304.029

Monoisotopic Mass: 303.029

Collision energy (CE): -15.0





Figure S1. LC-MS spectrum of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S2. ¹H NMR spectrum of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S3. ¹H NMR spectrum expansion of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S4. ^{13}C NMR of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S5. ^{13}C NMR spectrum expansion of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S6. ^{13}C NMR spectrum expansion of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S7. COSY contour map – ^1H x ^1H of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S8. COSY contour map – ^1H x ^1H expansion of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S9. HSQC contour map – ^1H x ^{13}C of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S10. HSQC contour map – ¹H x ¹³C expansion of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S11. HMBC contour map – ¹H x ¹³C of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S12. HMBC contour map – ^1H x ^{13}C expansion of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).




Figure S13. HMBC contour map – ^1H x ^{13}C expansion of 5'-chloro-2'-hydroxy-3'-nitrochalcone (4).

Compound name: 6-chloro-8-nitroflavanone

Molecular Formula: C₁₅H₁₀ClNO₄

Formula Weight: 303.697

Ionization mode: negative

Precursor [M + H]⁺: 302.029

Monoisotopic Mass: 303.029

Collision energy (CE): 25.0





Figure S14. LC-MS spectrum of 6-chloro-8-nitroflavanone.




Figure S15. ¹H NMR spectrum of 6-chloro-8-nitroflavanone.




Figure S16. ¹H NMR spectrum expansion of 6-chloro-8-nitroflavanone.




Figure S17. ¹³C NMR spectrum of 6-chloro-8-nitroflavanone.




Figure S18. ^{13}C NMR spectrum expansion of 6-chloro-8-nitroflavanone.




Figure S19. ^{13}C NMR spectrum expansion of 6-chloro-8-nitroflavanone




Figure S20. COSY contour map – ^1H x ^1H expansion of 6-chloro-8-nitroflavanone.




Figure S21. COSY contour map – ^1H x ^1H expansion of 6-chloro-8-nitroflavanone.




Figure S22. HSQC contour map – ^1H x ^{13}C of 6-chloro-8-nitroflavanone.




Figure S23. HSQC contour map – ^1H x ^{13}C expansion of 6-chloro-8-nitroflavanone.




Figure S24. HSQC contour map – ^1H x ^{13}C expansion of 6-chloro-8-nitroflavanone.




Figure S25. HMBC contour map – ^1H x ^{13}C of 6-chloro-8-nitroflavanone.




Figure S26. HMBC contour map – ^1H x ^{13}C expansion of 6-chloro-8-nitroflavanone.




Figure S27. HMBC contour map – ^1H x ^{13}C expansion of 6-chloro-8-nitroflavanone.

Compound name: 6-chloro-8-nitroflavone

Molecular Formula: C₁₅H₈ClNO₄

Formula Weight: 301.681

Ionization mode: positive

Precursor [M + H]⁺: 302.014

Monoisotopic Mass: 301.014

Collision energy (CE): -25.0





Figure S28. LC-MS spectrum of 6-chloro-8-nitroflavone.




Figure S29. ^1H NMR spectrum of 6-chloro-8-nitroflavone.




Figure S30. ¹H NMR spectrum expansion of 6-chloro-8-nitroflavone.




Figure S31. ¹³C NMR spectrum of 6-chloro-8-nitroflavone.




Figure S32. ^{13}C NMR spectrum expansion of 6-chloro-8-nitroflavone.




Figure S33. ^{13}C NMR spectrum expansion of 6-chloro-8-nitroflavone.




Figure S34. COSY contour map – ^1H x ^1H of 6-chloro-8-nitroflavone.




Figure S35. COSY contour map – ^1H x ^1H expansion of 6-chloro-8-nitroflavone.




Figure S36. HSQC contour map – ^1H x ^{13}C of 6-chloro-8-nitroflavone.




Figure S37. HSQC contour map – ^1H x ^{13}C expansion of 6-chloro-8-nitroflavone.




Figure S38. HSQC contour map – ^1H x ^{13}C expansion of 6-chloro-8-nitroflavone.




Figure S39. HMBC contour map – ^1H x ^{13}C of 6-chloro-8-nitroflavone.




Figure S40. HMBC contour map – ^1H x ^{13}C expansion of 6-chloro-8-nitroflavone.




Figure S41. HMBC contour map – ^1H x ^{13}C expansion of 6-chloro-8-nitroflavone.

Compound name: 8-amino-6-chloroflavone

4'-O- β -D-(4''-O-methyl)-glucopyranoside

Molecular Formula: C₂₂H₂₂ClNO₈

Formula Weight: 463.865

Ionization mode: positive

Precursor [M + H]⁺: 464.103

Monoisotopic Mass: 463.103

Collision energy (CE): -15.0




Figure S42. LC-MS spectrum of 8-amino-6-chloroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S43. ¹H NMR spectrum of 8-amino-6-chloroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S44. ¹H NMR spectrum expansion of 8-amino-6-chloroflavone 4'-O-β-D-(4''-O-methyl)-glucopyranoside.




Figure S45. ¹H NMR spectrum expansion of 8-amino-6-chloroflavone 4'-O-β-D-(4''-O-methyl)-glucopyranoside.




Figure S46. ^{13}C NMR spectrum of 8-amino-6-chloroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S47. ^{13}C NMR spectrum expansion of 8-amino-6-chloroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S48. ¹³C NMR spectrum expansion of 8-amino-6-chloroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S49. COSY contour map - ¹H x ¹H of 8-amino-6-chloroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S50. COSY contour map – ^1H x ^1H expansion of 8-amino-6-chloroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S51. COSY contour map – ^1H x ^1H expansion of 8-amino-6-chloroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S52. HSQC contour map – ^1H x ^{13}C of 8-amino-6-chloroflavone 4'-O- β -D-(4"-O-methyl)-glucopyranoside.




Figure S53. HSQC contour map – ^1H x ^{13}C expansion of 8-amino-6-chloroflavone 4'-O- β -D-(4"-O-methyl)-glucopyranoside.




Figure S54. HSQC contour map – ¹H x ¹³C expansion of 8-amino-6-chloroflavone 4'-O-β-D-(4''-O-methyl)-glucopyranoside.




Figure S55. HMBC contour map – ¹H x ¹³C of 8-amino-6-chloroflavone 4'-O-β-D-(4''-O-methyl)-glucopyranoside.

Compound name: 5'-bromo-2'-hydroxy-3'-nitrochalcone

Molecular Formula: C₁₅H₁₀BrNO₄

Formula Weight: 348.148

Ionization mode: positive

Precursor [M + H]⁺: 347.979

Monoisotopic Mass: 346.979

Collision energy (CE): -15.0





Figure S56. LC-MS spectrum of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S57. ¹H NMR spectrum of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S58. ¹H NMR spectrum expansion of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S59. ¹³C NMR spectrum of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S60. ^{13}C NMR spectrum expansion of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S61. ^{13}C NMR spectrum expansion of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S62. COSY contour map – ^1H x ^1H of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S20. COSY contour map – ^1H x ^1H expansion of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S64. HSQC contour map ${}^1\text{H}$ x ${}^{13}\text{C}$ of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S65. HSQC contour map ${}^1\text{H}$ x ${}^{13}\text{C}$ expansion of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S66. HMBC contour map – ^1H x ^{13}C of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S67. HMBC contour map – ^1H x ^{13}C expansion of 5'-bromo-2'-hydroxy-3'-nitrochalcone.




Figure S68. HMBC contour map – ^1H x ^{13}C expansion of 5'-bromo-2'-hydroxy-3'-nitrochalcone.

Compound name: 6-bromo-8-nitroflavanone

Molecular Formula: $\text{C}_{15}\text{H}_{10}\text{BrNO}_4$

Formula Weight: 348.148

Ionization mode: negative

Precursor $[\text{M} + \text{H}]^+$: 345.979

Monoisotopic Mass: 346.979

Collision energy (CE): +25.0





Figure S69. LC-MS spectrum of 6-bromo-8-nitroflavanone.




Figure S70. ¹H NMR spectrum of 6-bromo-8-nitroflavanone.




Figure S71. ¹H NMR spectrum expansion of 6-bromo-8-nitroflavanone.




Figure S72. ^{13}C NMR spectrum of 6-bromo-8-nitroflavanone.




Figure S73. ^{13}C NMR spectrum expansion of 6-bromo-8-nitroflavanone.




Figure S74. ^{13}C NMR spectrum expansion of 6-bromo-8-nitroflavanone.




Figure S75. ^{13}C NMR spectrum expansion of 6-bromo-8-nitroflavanone.




Figure S76. COSY contour map – ^1H x ^1H of 6-bromo-8-nitroflavanone.




Figure S77. COSY contour map – ^1H x ^1H expansion of 6-bromo-8-nitroflavanone.




Figure S78. COSY contour map – ^1H x ^1H expansion of 6-bromo-8-nitroflavanone.




Figure S79. HSQC contour map – ^1H x ^{13}C of 6-bromo-8-nitroflavanone.




Figure S80. HSQC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavanone.




Figure S81. HSQC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavanone.




Figure S82. HMBC contour map – ^1H x ^{13}C of 6-bromo-8-nitroflavanone.




Figure S83. HMBC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavanone.




Figure S84. HMBC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavanone.

Compound name: 6-bromo-8-nitroflavone

Molecular Formula: $\text{C}_{15}\text{H}_8\text{BrNO}_4$

Formula Weight: 346.132

Ionization mode: positive

Precursor $[\text{M} + \text{H}]^+$: 345.964

Monoisotopic Mass: 344.964

Collision energy (CE): -25.0





Figure S85. LC-MS spectrum of 6-bromo-8-nitroflavone.




Figure S86. ¹H NMR spectrum of 6-bromo-8-nitroflavone.




Figure S87. ¹H NMR spectrum expansion of 6-bromo-8-nitroflavone.




Figure S88. ^{13}C NMR spectrum of 6-bromo-8-nitroflavone.




Figure S89. ^{13}C NMR spectrum expansion of 6-bromo-8-nitroflavone.




Figure S90. ^{13}C NMR spectrum expansion of 6-bromo-8-nitroflavone.




Figure S91. COSY contour map - ^1H x ^1H of 6-bromo-8-nitroflavone.




Figure S92. COSY contour map – ^1H x ^1H expansion of 6-bromo-8-nitroflavone.




Figure S93. HSQC contour map – ^1H x ^{13}C of 6-bromo-8-nitroflavone.




Figure S94. HSQC contour map – ${}^1\text{H} \times {}^{13}\text{C}$ expansion of 6-bromo-8-nitroflavone.




Figure S95. HMBC contour map – ${}^1\text{H} \times {}^{13}\text{C}$ of 6-bromo-8-nitroflavone.




Figure S96. HMBC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavone.




Figure S97. HMBC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavone.




Figure S98. HMBC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavone.

Compound name: 6-Bromo-8-nitroflavone
4'-O- β -D-(4"-O-methyl)-glucopyranoside
Molecular Formula: C₂₂H₂₀BrNO₁₀
Formula Weight: 538.299
Ionization mode: positive
Precursor [M + H]⁺: 539.030
Monoisotopic Mass: 537.027
Collision energy (CE): -25.0





Figure S99. LC-MS spectrum of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S100. ¹H NMR spectrum of 6-bromo-8-nitroflavone 4'-O-β-D-(4''-O-methyl)-glucopyranoside.




Figure S101. ¹H NMR spectrum expansion of 6-bromo-8-nitroflavone 4'-O-β-D-(4''-O-methyl)-glucopyranoside.




Figure S102. ¹H NMR spectrum expansion of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S103. ¹³C NMR spectrum of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S104. ^{13}C NMR spectrum expansion of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S105. ^{13}C NMR spectrum expansion of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S106. COSY contour map – ^1H x ^1H of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S107. COSY contour map – ^1H x ^1H expansion of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S108. COSY contour map – ^1H x ^1H expansion of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S109. HSQC contour map – ^1H x ^{13}C of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S110. HSQC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S111. HSQC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S112. HMBC contour map – ^1H x ^{13}C of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S113. HMBC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.




Figure S114. HMBC contour map – ^1H x ^{13}C expansion of 6-bromo-8-nitroflavone 4'-O- β -D-(4''-O-methyl)-glucopyranoside.