

## **Supplementary material**

**Title: Associations between NK Cells in Different Immune Organs and Cellular SIV DNA and RNA in Regional HLA<sup>DR</sup><sup>-</sup> CD4<sup>+</sup> T Cells in Chronically SIV<sub>mac239</sub>-Infected, Treatment-Naïve Rhesus Macaques**

## Supplementary Figures and Tables

### 1 Supplementary Table

**Table S1.** The characteristics of SIV-infected Chinese rhesus macaques in this study.

Animal No.	Sex	Weight (kg)	Age (months)	Viral stock	Routine of inoculation	Viral quantification	Time of inoculation	Time of euthanasia
G0101R	F	4.15	49	SIV <sub>mac</sub> 239	Ir	100 TCID50	Jan 3, 2013	Jan 21, 2015
G0102R	F	4.25	51	SIV <sub>mac</sub> 239	Ir	100 TCID50	Jan 3, 2013	Jan 21, 2015
G0104R	F	4.55	54	SIV <sub>mac</sub> 239	Ir	100 TCID50	Jan 3, 2013	Jan 21, 2015
G0105R	F	4.8	47	SIV <sub>mac</sub> 239	Ir	100 TCID50	Jan 3, 2013	Jan 21, 2015
G0106R	F	3.9	55	SIV <sub>mac</sub> 239	Ir	100 TCID50	Jan 3, 2013	Jan 21, 2015

Ir, intrarectally; SIV, simian immunodeficiency virus; TCID, tissue culture infectious dose.

## 2 Supplementary Figures

### Supplementary Figure S1




Figure S1. Dynamics of peripheral CD4<sup>+</sup> T and CD8<sup>+</sup> T cell counts in the bloodstream of 5 SIV-infected Chinese Rhesus macaques.

## Supplementary Figure S2




Figure S2. Dynamics of plasma SIV viral load in 5 SIV-infected Chinese Rhesus Macaques.