

Table S1



Univariable analysis of factors associated with overall survival.



			Hazard ratio (95% CI, p-value)
Age (years)	<i>Mean (SD)</i>	26.7 (16.3)	1.03 (1.00-1.06, p=0.050)
WHO CNS grade	2	58 (65.2)	-
	1	17 (19.1)	0.38 (0.11-1.37, p=0.139)
Location	<i>Frontotemporal</i>	31 (34.8)	-
	<i>Visual/thalamic</i>	43 (48.3)	0.28 (0.08-1.02, p=0.054)
	<i>Other</i>	15 (16.9)	1.23 (0.34-4.48, p=0.758)
Indication	<i>Primary, definitive</i>	18 (20.2)	-
	<i>Primary, postoperative</i>	28 (31.5)	1.91 (0.26-14.21, p=0.528)
	<i>Recurrence/progression, definitive</i>	33 (37.1)	2.84 (0.60-13.45, p=0.189)
	<i>Recurrence/progression, postoperative</i>	10 (11.2)	2.91 (0.41-20.88, p=0.289)
Initial seizures	<i>No</i>	55 (61.8)	-
	<i>Yes</i>	34 (38.2)	1.55 (0.54-4.40, p=0.415)



Figure S1



Quality of life: additional scores.

EORTC QLQ-C30












Diarrhoe (C30)




Financial difficulties (C30)





Short breath (C30)




Sleeping (C30)




EORTC QLQ-BN20




Motor dysfunction (BN20)



Visual disorder (BN20)



Bladder control (BN20)



Drowsiness (BN20)

