

Article

More Insights into Symmetries in Multisymplectic Field Theories

Arnoldo Guerra IV ¹  and Narciso Román-Roy ^{2,*} ¹ Departament de Física Quàntica i Astrofísica, Universitat de Barcelona, 08028 Barcelona, Spain² Department of Mathematics, Campus Nord, Universitat Politècnica de Catalunya, 08034 Barcelona, Spain

* Correspondence: narciso.roman@upc.edu

Abstract: This work provides a general overview for the treatment of symmetries in classical field theories and (pre)multisymplectic geometry. The geometric characteristics of the relation between how symmetries are interpreted in theoretical physics and in the geometric formulation of these theories are clarified. Finally, a general discussion is given on the structure of symmetries in the presence of constraints appearing in singular field theories. Symmetries of some typical theories in theoretical physics are analyzed through the construction of the relevant multimomentum maps which are the conserved quantities (by Noether's theorem) on the (pre)multisymplectic phase spaces.

Keywords: Lagrangian and Hamiltonian field theories; jet bundles; multisymplectic forms; symmetries; noether symmetries; gauge symmetries; conserved quantities; multimomentum maps

MSC: Primary: 53D42, 70S05, 83C05; Secondary: 35Q75, 35Q76, 53Z05, 70H50, 83C99

1. Introduction

One of the main research topics in mathematical physics is the study of symmetries in dynamical (mechanical) systems and field theories. From a more mathematical point of view, such studies of symmetries involve the analysis of ordinary and partial differential equations; this is because symmetries are associated with conserved quantities or conservation laws, which give fundamental information about the physical system. The foundational work on symmetries in physical systems is that of *Emmy Noether*, ref. [1]. The general underlying concept of symmetry in a physical system was first presented as the invariance of the equations of motion under a transformation on the phase space of the system. In the more modern geometric formulation of classical mechanics and classical field theories, symmetries are usually characterized by demanding the invariance of some underlying geometric structure from which the preservation of the equations of motions is included as a consequence.

The *multisymplectic formulation* of classical field theories takes place on the *multivelocety* and *multimomentum phase spaces* where the Lagrangian and De Donder–Weyl Hamiltonian formulations are developed. These phase spaces are fiber bundles $q: \mathcal{M} \rightarrow M$ over an orientable m -dimensional manifold M (typically spacetime) where $m > 1$. In particular, \mathcal{M} is either a jet bundle in the Lagrangian formalism or a bundle of forms (or a quotient of them) in the Hamiltonian formalism. This formulation of classical field theories can be viewed as a geometric extension of the classical mechanics of non-autonomous systems for which $m = 1$. Furthermore, the phase spaces for field theories are endowed with a characteristic geometric structure: a multisymplectic or premultisymplectic form $\Omega \in \Omega^{m+1}(\mathcal{M})$ (depending on the regularity of the theory). In practice, the (pre)multisymplectic structures are constructed by starting from the Lagrangian function of the field theory under investigation [2–7]. Then, the field equations (the Euler–Lagrange equations or the the Hamilton–De Donder–Weyl equations) are stated geometrically using the corresponding (pre)multisymplectic forms.



Citation: Guerra, A., IV; Román-Roy, N. More Insights into Symmetries in Multisymplectic Field Theories. *Symmetry* **2023**, *15*, 390. <https://doi.org/10.3390/sym15020390>

Academic Editor: Jorge Segovia

Received: 30 December 2022

Revised: 26 January 2023

Accepted: 28 January 2023

Published: 1 February 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

All symmetries, along with their associated conserved quantities and conservation laws, have been studied extensively in the geometric framework, both in the Lagrangian setting [2–5,7–12] and in the corresponding De Donder–Weyl Hamiltonian setting [6,9,13], where the symmetries which preserve the (pre)multisymplectic forms are called *Noether* or *Cartan symmetries*. The geometric presentation of Noether symmetries in field theories culminates with the statement of the geometric version of Noether’s theorem [4,9,13–17]. We have paid special attention to the *multimomentum maps* which are the fundamental conserved quantities associated with Noether symmetries. These are the generalization of the *momentum maps* of symplectic mechanics to the multisymplectic setting for field theory. (Multi)momentum maps are the relevant quantities for performing a *symmetry reduction procedure*. However, the multisymplectic reduction of field theories is currently, in general, an unsolved problem under research [18–21]. For the analogous geometric construction of symmetries in (first-order) non-autonomous mechanical systems in which $M = \mathbb{R}$, the corresponding results for the Lagrangian and Hamiltonian formalisms can be found, for instance, in [22–24].

Singular field theories are invariant under a family of Noether symmetries called *geometric gauge symmetries* which are linked to the degeneracy of the Lagrangian and are generated on the premultisymplectic phase spaces by vector fields which lie in the kernel of the premultisymplectic forms. However, the term “gauge symmetry” is used in various different contexts throughout the literature [25]; such contexts will be discussed in Section 2.4. The so-called geometric gauge symmetries form *gauge orbits* on the premultisymplectic phase space which define equivalence classes. Physical states which lie in the same gauge orbit differ from one another by a geometric gauge transformation and are said to be *gauge equivalent*.

This paper has three main aims which are as follows. The first aim of this work is to review and broaden the discussion about the geometric structures and properties regarding the study of symmetries of first-order classical field theories and is mainly based on the previous papers [14,15]. The second main goal of this work is to clarify and enhance the relationship between the geometric treatment of classical field theories found in the mathematics literature and the equivalent approaches which are standard in theoretical physics. We believe this second goal is especially important as there seem to be several gaps in the dialogue between the communities of theoretical physics and differential geometry in this regard. In the theoretical physics literature, it is standard to work on spaces of sections of some of the fiber bundles mentioned here, while in the differential geometry literature, the analysis of field theories is conducted on the fiber bundles themselves. For example, in theoretical physics it is standard to develop the Lagrangian formulation of field theories on the space of sections of the multivelocity phase space where the Lagrangian function is taken to be a functional of the fields and its spacetime derivatives, while in the mathematics literature, typically, the Lagrangian function is taken to be a function on the multivelocity phase space itself as mentioned earlier. The equivalence between these approaches will be made precise throughout the paper with Sections 3.2 and 3.3 being of utmost importance to this aim. Finally, the third aim of this paper is to detail some of the subtleties that arise in the analysis of symmetries when dealing with field theories that are singular in the De Donder–Weyl sense. The main goals of this paper mentioned above will be further pursued through the geometric analysis of specific field theories that are relevant in theoretical physics.

This paper is organized as follows. Section 2 reviews the main definitions and results regarding the various types of symmetries along with the corresponding conserved quantities and multimomentum maps in the geometric framework of generic (pre)multisymplectic bundles. Some new clarifying ideas regarding Noether and gauge symmetries are also provided. Section 3 is devoted to explaining how to lift diffeomorphisms and vector fields from the base space to the corresponding jet bundle; this is important in order to characterize the most common types of symmetries in physics. In Section 4, which contains the main contribution of the work, the definitions and results of Section 2 are adapted to

the Lagrangian and Hamiltonian descriptions of classical field theories. Additionally, the so-called *Lagrangian symmetries* arising in Lagrangian field theories are discussed. The connection with the physical interpretation of all these ideas is detailed. Finally, in Section 5, the symmetries of some typical theories in theoretical physics are displayed. Basic notions on multivector fields are given in Appendix A.

The classical field theories discussed here exhibit various kinds of symmetries that are typical in theoretical physics. The Nambu–Goto action for bosonic strings, which is invariant under the full group of diffeomorphisms (local point transformations) of the string worldsheet, is studied in the multisymplectic setting in Section 5.1. The string action can be thought of as a matter action of a two-dimensional field theory coupled to gravity; it is well-known that, in this sense, the Nambu–Goto action is invariant only under spacetime isometries and not under the full group of spacetime diffeomorphisms as is General Relativity. This is typical for matter theories, including Yang–Mills which is presented subsequently. The symmetry of spacetime isometries, as well as the $SU(N)$ gauge symmetry, of Yang–Mills is discussed Section 5.2. The Chern–Simons theory with gauge group $SU(N)$ is the only theory discussed in this work (in Section 5.3) that is invariant under the full group of spacetime diffeomorphisms. Finally, in Sections 5.4 and 5.5, we discuss the electric and magnetic Carrollian scalar field theories which are invariant only under Carroll transformations of spacetime; for more information on Carrollian field theories see, for example, ref. [26] and references therein.

Some comments about notation: $\mathfrak{X}(\mathcal{M})$ and $\Omega^k(\mathcal{M})$ denote the $C^\infty(\mathcal{M})$ -modules of vector fields and differential k -forms on a manifold \mathcal{M} , respectively; $L(X)\Omega$ denotes the Lie derivative of a differential form Ω with respect to a vector field X , and $i(X)\Omega$ is the inner contraction between X and Ω .

All manifolds are taken to be real, second countable, and C^∞ . All maps are also C^∞ . Sum over repeated indices (Einstein notation) is understood.

2. Symmetries on (Pre)multisymplectic Fiber Bundles

This section is devoted to the introduction of the main concepts and properties regarding symmetries and conserved quantities (i.e., conservation laws) on (pre)multisymplectic manifolds. This presentation is made in the generic geometric framework of (pre)multisymplectic fiber bundles which are the relevant structures on which classical field theories are constructed. The structures discussed in this section will later be particularized to the specific contexts of the Lagrangian and Hamiltonian formalisms for classical field theories. The fundamental ideas in this section are taken from [14]. For the proofs of the results and other details, see also [13–15].

2.1. (Pre)Multisymplectic Bundles

Given a differentiable manifold \mathcal{M} , a differential form $\Omega \in \Omega^m(\mathcal{M})$ is 1-nondegenerate if, for every $p \in \mathcal{M}$ and $Y \in \mathfrak{X}(\mathcal{M})$, it follows that $i(Y)\Omega|_p = 0 \iff Y|_p = 0$. Then, $\Omega \in \Omega^m(\mathcal{M})$ is said to be a **multisymplectic form** if it is closed and 1-nondegenerate and (\mathcal{M}, Ω) is called a **multisymplectic manifold**. Alternatively, Ω is called a **premultisymplectic form** if it is closed and 1-degenerate and (\mathcal{M}, Ω) is called a **premultisymplectic manifold**. If Ω is an exact form, then it is called an **exact (pre)multisymplectic form** and the couple (\mathcal{M}, Ω) is called a **(pre)multisymplectic system**.

The geometric framework for Lagrangian and Hamiltonian field theories consists of a fiber bundle $q: \mathcal{M} \rightarrow M$ over an orientable manifold M ($\dim M = m > 1$, $\dim \mathcal{M} = N + m$). The volume form on M is denoted $\omega \in \Omega^m(M)$ (with ω also denoting $q^*\omega$), and \mathcal{M} is endowed with an exact multisymplectic or premultisymplectic form (depending on the regularity of the theory) $\Omega = -d\Theta \in \Omega^{m+1}(\mathcal{M})$ for some $\Theta \in \Omega^m(\mathcal{M})$. Furthermore, the (pre)multisymplectic form satisfies the *variational condition* $i(Z_1)i(Z_2)i(Z_3)\Omega = 0$, for all q -vertical vector fields $Z_1, Z_2, Z_3 \in \mathfrak{X}^{V(q)}(\mathcal{M})$, which allows for the field equations to be obtained from a variational principle.

The solution to the variational problem on \mathcal{M} is sections $\psi: M \rightarrow \mathcal{M}$ of the projection ϱ which solve the field equations given by

$$\psi^* i(X)\Omega = 0, \quad \text{for every } X \in \mathfrak{X}(\mathcal{M}).$$

Equivalently, the solutions to the variational problem are integral sections of a m -multivector field (see Appendix A) contained in a class of integrable, ϱ -transverse (i.e., $\mathbf{X} \in \ker^m \Omega$), m -multivector fields $\{\mathbf{X}\} \subset \mathfrak{X}^m(\mathcal{M})$ which satisfy the field equations which are now written as

$$i(\mathbf{X})\Omega = 0, \tag{1}$$

where the ϱ -transversality condition can be set, without loss of generality, as

$$i(\mathbf{X})\omega = 1. \tag{2}$$

The couple (\mathcal{M}, Ω) is usually called a (pre)multisymplectic system.

In general, solvable premultisymplectic systems admit stable solutions to the field equations on some submanifold $\mathcal{S} \subseteq \mathcal{M}$ which is obtained by applying a *constraint algorithm* (see Section 4.4 and [27] for more details); the submanifold \mathcal{S} is called the *final constraint submanifold*, and the physical states of the field theory under investigation are the sections of the projection ϱ whose images are on \mathcal{S} .

2.2. Conserved Quantities and Symmetries

In geometric mechanics, conserved quantities are functions on the phase space which are invariant under the advance along the dynamical trajectories given by the integral curves of some dynamical vector fields. The invariance condition is stated by demanding the vanishing of the Lie derivative of the functions representing the conserved quantity with respect to the dynamical vector field solution to the equations of motion. For field theories, there is an analogous geometrical way to introduce this concept using multivector fields to represent the solutions to the field equations.

Let (\mathcal{M}, Ω) be a (pre)multisymplectic system.

Definition 1. A *conserved quantity* of the (pre)multisymplectic system (\mathcal{M}, Ω) is a form $\alpha \in \Omega^{m-1}(\mathcal{M})$ which satisfies $L(\mathbf{X})\alpha = (-1)^{m+1} i(\mathbf{X})d\alpha = 0$ for every locally decomposable and ϱ -transverse multivector field $\mathbf{X} \in \ker^m \Omega$ (i.e., which satisfies Equations (1) and (2)).

Conserved quantities are characterized by the following property: if $\alpha \in \Omega^{m-1}(\mathcal{M})$ is a conserved quantity and $\mathbf{X} \in \ker^m \Omega$ is a ϱ -transverse integrable multivector field, then α is closed on the integral submanifolds of \mathbf{X} ; that is, if $j_S: S \hookrightarrow \mathcal{M}$ is an integral submanifold of \mathbf{X} , then $dj_S^* \alpha = 0$.

Remark 1. Conserved quantities in field theories appear as conservation laws or conserved currents as, for every $\alpha \in \Omega^{m-1}(\mathcal{M})$ and $\mathbf{X} \in \mathfrak{X}^m(\mathcal{M})$, if $\psi: M \rightarrow \mathcal{M}$ is an integral section of \mathbf{X} such that $\psi^* \alpha \in \Omega^{m-1}(M)$, then there is a unique vector field $X_{\psi^* \alpha} \in \mathfrak{X}(M)$ such that $i(X_{\psi^* \alpha})\eta = \psi^* \alpha$. It follows that the divergence of $X_{\psi^* \alpha}$ is the function $\text{div} X_{\psi^* \alpha} \in C^\infty(M)$ defined as $L(X_{\psi^* \alpha})\omega = (\text{div} X_{\psi^* \alpha}) \Omega$ which satisfies $(\text{div} X_{\psi^* \alpha}) \eta = d\psi^* \alpha$. Therefore, as a consequence of the above property, α is a conserved quantity if, and only if, $\text{div} X_{\psi^* \alpha} = 0$; hence, Stokes theorem assures that in every bounded domain $U \subset M$,

$$\int_{\partial U} \psi^* \alpha = \int_U (\text{div} X_{\psi^* \alpha}) \eta = \int_U d\psi^* \alpha = 0.$$

The form $\psi^* \alpha$ is called the **current** associated with the conserved quantity α . This result associates a **conservation law** on M to every conserved quantity on \mathcal{M} .

The field equations are partial differential equations (PDE's), and symmetries are (local) diffeomorphisms that transform solutions to the field equations into equivalent solutions. In geometric mechanics, this property is stated using the vector fields which generate the symmetries (by their flows) and the dynamical vector fields which solve the equations of motion. For field theories, there is a similar geometric procedure using multivector fields which are solutions to the field equations as follows:

Definition 2.

1. A **symmetry** of the (pre)multisymplectic system (\mathcal{M}, Ω) is a diffeomorphism $\Phi: \mathcal{M} \rightarrow \mathcal{M}$ such that $\Phi_*(\ker^m \Omega) \subset \ker^m \Omega$.
2. An **infinitesimal symmetry** of the (pre)multisymplectic system (\mathcal{M}, Ω) is a vector field $Y \in \mathfrak{X}(\mathcal{M})$ whose local flows are local symmetries or, equivalently, $[Y, \ker^m \Omega] \subset \ker^m \Omega$.

Remark 2. If (\mathcal{M}, Ω) is a premultisymplectic system with final constraint submanifold $J_S: \mathcal{S} \hookrightarrow \mathcal{M}$, then symmetries must be diffeomorphisms $\Phi \in \text{Diff}(\mathcal{M})$ such that $\Phi(\mathcal{S}) = \mathcal{S}$, and infinitesimal symmetries are vector fields $Y \in \mathfrak{X}(\mathcal{M})$ tangent to \mathcal{S} which satisfy the conditions in Definition 2 at least on \mathcal{S} . The set of vector fields on \mathcal{M} which are tangent to \mathcal{S} will be denoted here as

$$\mathfrak{X}(\mathcal{S}) := \{Y \in \mathfrak{X}(\mathcal{M}) \mid \exists Y_S \in \mathfrak{X}(\mathcal{S}) \mid J_{S*}Y_S = Y|_{\mathcal{S}}\}.$$

The elements of $\mathfrak{X}(\mathcal{S})$ are the vector fields $Y \in \mathfrak{X}(\mathcal{M})$ which, for some $Y_S \in \mathfrak{X}(\mathcal{S})$, make the following diagram commutative:

$$\begin{array}{ccc} \mathcal{M} & \xrightarrow{\quad Y \quad} & T\mathcal{M} \\ J_S \uparrow & & \uparrow T J_S \\ \mathcal{S} & \xrightarrow{\quad Y_S \quad} & T\mathcal{S} \end{array}$$

As we have said, symmetries transform solutions to the field equations into solutions, since:

Theorem 1. If $\Phi \in \text{Diff}(\mathcal{M})$ is a symmetry that restricts to a diffeomorphism $\varphi: M \rightarrow M$, then, for every q -transverse integrable multivector field \mathbf{X} , the map Φ transforms integral sections of \mathbf{X} into integral sections of $\Phi_*\mathbf{X}$ and hence $\Phi_*\mathbf{X} \in \ker^m \Omega$ which is also a q -transverse integrable multivector field. Consequently, if $Y \in \mathfrak{X}(\mathcal{M})$ is a π^1 -projectable infinitesimal symmetry and F_t is the local flow of Y , then F_t transforms integral sections of every q -transverse integrable multivector field $\mathbf{X} \in \ker^m \Omega$ into integral sections of $F_{t*}\mathbf{X}$.

If $Y_1, Y_2 \in \mathfrak{X}(\mathcal{M})$ are infinitesimal symmetries, then the Lie bracket $[Y_1, Y_2]$ is also an infinitesimal symmetry. Furthermore, if $\Phi \in \text{Diff}(\mathcal{M})$ is a symmetry and $\alpha \in \Omega^{m-1}(\mathcal{M})$ is a conserved quantity, then $\Phi^*\alpha$ is also a conserved quantity. Consequently, if $Y \in \mathfrak{X}(\mathcal{M})$ is an infinitesimal symmetry, then $L(Y)\alpha$ is also a conserved quantity.

Remark 3. Transformations of the base space M (usually spacetime transformations in field theory) which preserve the (pre)multisymplectic structure of the phase spaces are called **spacetime symmetries**. Therefore, in the bundle $q: \mathcal{M} \rightarrow M$, the corresponding diffeomorphisms $\Phi: \mathcal{M} \rightarrow \mathcal{M}$ must be fiber preserving and thereby restrict to diffeomorphisms $\Phi_M: M \rightarrow M$ which satisfy $\Phi_M \circ q = q \circ \Phi$. For infinitesimal symmetries, this means that the vector fields $Y \in \mathfrak{X}(\mathcal{M})$ must be q -projectable; hence, there exist $Y_M \in \mathfrak{X}(M)$ such that $q_*Y = Y_M$.

Symmetries are generated by the action of a Lie group G on \mathcal{M} :

$$\begin{array}{ccc} \Phi & : & G \rightarrow \text{Diff}(\mathcal{M}) \\ & & g \mapsto \Phi_g \end{array}$$

where $\Phi_g : \mathcal{M} \rightarrow \mathcal{M}$ are the local diffeomorphisms induced by the group action. If \mathfrak{g} is the Lie algebra of G , it follows that every $\xi \in \mathfrak{g}$ induces a vector field $X_\xi \in \mathfrak{X}(\mathcal{M})$ which generates the diffeomorphisms produced by the group action of G on \mathcal{M} . This is achieved by constructing the exponential map for some $g \in G$:

$$\begin{aligned} \exp &: \mathfrak{g} \rightarrow G \\ \xi &\mapsto g = \exp(\lambda\xi) \end{aligned}$$

where $\lambda \in \mathbb{R}$ is a parameter. Then, the induced vector field X_ξ on \mathcal{M} is written in terms of the exponential map as

$$X_\xi = \left. \frac{d}{d\lambda} \right|_{\lambda=0} \Phi_{\exp(\lambda\xi)}^* \in \mathfrak{X}(\mathcal{M}).$$

These vector fields are the *infinitesimal generators* of the symmetries, that is, the infinitesimal symmetries.

2.3. Noether Symmetries

In order to obtain conserved quantities (or conservation laws) associated with symmetries, it is necessary to impose some additional conditions. In general, these conditions are related to the geometric structures underlying the field equations (see, for instance, refs. [24,28] and the references therein, for a complete study in Lagrangian and Hamiltonian mechanics). The most relevant kinds of symmetries are the following:

Definition 3.

1. A **Noether or Cartan symmetry** of the (pre)multisymplectic system (\mathcal{M}, Ω) is a diffeomorphism $\Phi: \mathcal{M} \rightarrow \mathcal{M}$ such that, $\Phi^*\Omega = \Omega$. In the particular case where $\Phi^*\Theta = \Theta$, then Φ is called an **exact Noether or exact Cartan symmetry**.
2. An **infinitesimal Noether or Cartan symmetry** of the (pre)multisymplectic system (\mathcal{M}, Ω) is a vector field $Y \in \mathfrak{X}(\mathcal{M})$ for which $L(Y)\Omega = 0$. In the particular case where $L(Y)\Theta = 0$, Y is called an **infinitesimal exact Noether or infinitesimal exact Cartan symmetry**.

If $Y_1, Y_2 \in \mathfrak{X}(\mathcal{M})$ are infinitesimal Noether symmetries, then so is the Lie bracket $[Y_1, Y_2]$. Furthermore, if $Y \in \mathfrak{X}(\mathcal{M})$ is an infinitesimal Noether symmetry and $X \in \ker^m \Omega$, bearing in mind (A1), we have that

$$i([Y, X])\Omega = L(Y)i(X)\Omega - i(X)L(Y)\Omega = 0 \iff [Y, X] \subset \ker^m \Omega,$$

and hence, infinitesimal Noether symmetries are infinitesimal symmetries (and the same holds for Noether symmetries).

Remark 4. As $L(Y)\Omega_{\mathcal{L}} = d i(Y)\Omega_{\mathcal{L}}$, the condition $L(Y)\Omega = 0$ is equivalent to demanding that $i(Y)\Omega$ is a closed m -form on \mathcal{M} . It follows that $i(Y)\Omega = -d\alpha_Y$ is an open set $U \subset \mathcal{M}$ for some $\alpha_Y \in \Omega^{m-1}(U)$ (where the minus sign is due to physical conventions). Thus, an infinitesimal Noether symmetry is a locally Hamiltonian vector field for the (pre)multisymplectic form Ω and α_Y is the corresponding local Hamiltonian form.

Therefore, **Noether’s theorem** is stated as follows:

Theorem 2 (Noether). Let $Y \in \mathfrak{X}(\mathcal{M})$ be an infinitesimal Noether symmetry with $i(Y)\Omega = -d\alpha_Y$ in an open set $U \subset \mathcal{M}$. Then, for every locally decomposable and q -transverse (integrable) multivector field $X \in \ker^m(\Omega)$, it follows that

$$L(X)\alpha_Y = 0;$$

that is, every Hamiltonian $(m - 1)$ -form α_Y associated with Y is a conserved quantity. For every integral section ψ of X , the form $\psi^*\alpha_Y$ is usually called a **Noether current**.

Observe that the form $L(Y)\Theta$ is closed since

$$L(Y)\Theta = d i(Y)\Theta + i(Y)d\Theta = d i(Y)\Theta - i(Y)\Omega = d(i(Y)\Theta - \alpha_Y) \equiv d\zeta_Y \quad (\text{in } U). \quad (3)$$

In particular, if Y is an exact infinitesimal Noether symmetry, then ζ_Y is closed and $\alpha_Y = -i(Y)\Theta$.

Remark 5. As it was pointed out in Remark 3, we will mainly be interested in Noether symmetries which are fiber-preserving diffeomorphisms for the bundle $q: \mathcal{M} \rightarrow M$ and in infinitesimal Noether symmetries which are q -projectable vector fields.

Furthermore, if (\mathcal{M}, Ω) is a premultisymplectic system with final constraint submanifold $\mathcal{S} \hookrightarrow \mathcal{M}$, then Noether symmetries must be diffeomorphisms $\Phi \in \text{Diff}(\mathcal{M})$ such that $\Phi(\mathcal{S}) = \mathcal{S}$, and infinitesimal Noether symmetries must be vector fields $Y \in \mathfrak{X}(\mathcal{M})$ tangent to \mathcal{S} which satisfy the conditions of Definition 3, at least on \mathcal{S} .

2.4. Gauge Symmetries

The term “gauge” is used in various different contexts throughout the literature. In physics, it is typically used to refer to field transformations which are spacetime dependent. This section discusses some of the different meanings associated with the term “gauge” within the context of classical field theory. In particular, the description of a certain type of gauge symmetries that are the most relevant from both a geometric and physical point of view will be given. See also [15,29] for additional insights.

Let (\mathcal{M}, Ω) be a (pre)multisymplectic system and recall that Noether symmetries preserve the (pre)multisymplectic form Ω . Only fiber-preserving symmetries will be investigated, bearing in mind Remarks 3 and 5 which characterize symmetries in the context of the final constraint submanifold $\mathcal{S} \subset \mathcal{M}$ of the premultisymplectic system under investigation. Furthermore, the particular situation where such symmetries restrict to the identity on M will be the kind which will be generically referred to here as **gauge symmetries**. Then, **infinitesimal gauge symmetries** are vector fields $Y \in \mathfrak{X}(\mathcal{M})$ whose local flows are local gauge symmetries and, consequently, are q -vertical vector fields. In this respect, it should be noted that, the set of q -vertical vector fields are $\mathfrak{X}^{V(q)}(\mathcal{M}) = \ker \omega$.

Furthermore, infinitesimal Noether symmetries are vector fields $Y \in \mathfrak{X}(\mathcal{M})$ characterized by the property that $d i(Y)\Omega = 0$. When $\ker \Omega \neq \{0\}$, then (\mathcal{M}, Ω) is a premultisymplectic system. Then, any nonzero vector field $Y \in \ker \Omega$ (i.e., $i(Y)\Omega = 0$) is a particular type of infinitesimal symmetry that is related to the degeneracy of the premultisymplectic form Ω . Such vector fields will be referred to as **geometric (infinitesimal) gauge symmetries**.

Moreover, when (\mathcal{M}, Ω) is a premultisymplectic system, the field equations, in general, have consistent solutions on a final constraint submanifold $\mathcal{S} \hookrightarrow \mathcal{M}$ (as stated at the end of Section 2.1) which is the manifold where sections representing the physical states take their image. As above, the physically relevant gauge transformations are those which transform physical states into equivalent physical states and hence are automorphisms of the submanifold \mathcal{S} which transform sections of the projection q into themselves. Consequently, these gauge transformations which act along the fibers of the projection q are necessarily generated by q -vertical vector fields.

The concepts above are summarized as follows:

Definition 4. A **geometric infinitesimal gauge symmetry**, or simply a **gauge vector field**, of a (pre)multisymplectic system (\mathcal{M}, Ω) is a vector field $Y \in \mathfrak{X}(\mathcal{M})$ such that:

1. $Y \in \ker \Omega$.
2. It is a q -vertical vector field, $Y \in \mathfrak{X}^{V(q)}(\mathcal{M})$.
3. It is tangent to \mathcal{S} , $Y \in \mathfrak{X}(\mathcal{S})$.

The set of gauge vector fields is denoted \mathcal{G} .

The local diffeomorphisms generated by the flow of gauge vector fields are called **geometric gauge transformations**. Physical states (i.e., sections which are stable solutions to the field equations) related to one another by geometric gauge transformations are called **geometric gauge equivalent states**, and they are physically equivalent (in the sense that they are physically indistinguishable).

Remark 6. In the case where $J_S : \mathcal{S} \hookrightarrow \mathcal{M}$ is strictly a submanifold of \mathcal{M} , besides those of $\ker \Omega$, there are more vector fields related to the non-regularity of the premultiplectic form Ω . Such vector fields belong to the set

$$\ker \Omega_S := \{Z \in \mathfrak{X}(\mathcal{S}) \mid \exists Z_S \in \ker \Omega_S \mid J_{S*}Z_S = Z|_{\mathcal{S}}\},$$

where $\Omega_S = J_S^*\Omega$ which, in general, is also a premultisymplectic form. The vector fields in $\ker \Omega_S$ are those which are tangent to \mathcal{S} and can be identified on the points of \mathcal{S} with the elements of $\ker \Omega_S$. Since $\ker \Omega \cap \mathfrak{X}(\mathcal{S}) \subseteq \ker \Omega_S$, then for every $Z \in \ker \Omega \cap \mathfrak{X}(\mathcal{S})$, it follows that

$$i(Z)\Omega = 0 \implies 0 = J_S^* i(Z)\Omega = i(Z_S)\Omega_S \iff Z_S \in \ker \Omega_S,$$

and, as a consequence,

$$\mathcal{G} \equiv \ker \Omega \cap \mathfrak{X}(\mathcal{S}) \cap \mathfrak{X}^{V(q)}(\mathcal{M}) \subseteq \ker \Omega_S \cap \mathfrak{X}^{V(q)}(\mathcal{M}) \equiv \widehat{\mathcal{G}}.$$

In the case of singular (autonomous) dynamical systems in mechanics, it is proved [30,31] that $\widehat{\mathcal{G}}$ is the complete set of gauge vector fields. Then, the elements of \mathcal{G} are called **primary gauge vector fields**, and those in $\widehat{\mathcal{G}} - \mathcal{G}$ are the **secondary gauge vector fields**. The analogous concept in classical field theories regarding the classification of $\mathcal{G} - \mathcal{G}$ as gauge vector fields (i.e., if they relate geometric gauge equivalent states through their flows) is still an open question.

As usual, if $Z_1, Z_2 \in \mathcal{G}$, then $[Z_1, Z_2] \in \mathcal{G}$ so \mathcal{G} generates an involutive distribution on \mathcal{M} , and the quotient set $\widetilde{\mathcal{S}} = \mathcal{S}/\mathcal{G}$, assumed to be a differentiable manifold, plays the relevant role in the so-called **geometric gauge reduction procedure** [30,31] which consists of taking this reduced manifold as the set of “real physical” states and then removes the unphysical redundancy represented by the geometric gauge equivalent states. Introducing the canonical projection $\tilde{\tau}_S : \mathcal{S} \rightarrow \widetilde{\mathcal{S}}$, gauge equivalent states are sections of this projection. An alternative way to remove the “geometric gauge redundancy” is by performing a **gauge fixing** procedure which consists of fixing sections of this projection. A summarized description of these two procedures may be found in [15]. The most general method for gauge fixing in field theory involves the BRST-BV method (see, for example, [32,33]).

As a last comment, it should be noted that sometimes in physics a broader definition of (infinitesimal) gauge symmetry is handled which pertains to vector fields $Y \in \mathfrak{X}(\mathcal{M})$ such that $d i(Y)\Omega = 0$, which, in the terminology used in this work, are the so-called infinitesimal Noether symmetries.

2.5. Multimomentum Map

Let G be a group of Noether symmetries for a (pre)multisymplectic system (\mathcal{M}, Ω) ; that is, $\Phi_g^*\Omega = \Omega$, for every $g \in G$, and $L(X_\xi)\Omega = 0$, for every $\xi \in \mathfrak{g}$. Therefore, as $d i(X_\xi)\Omega = 0$, there exists a form $J_\xi \in \Omega^{m-1}(U)$ on an open neighbourhood $U \subset \mathcal{M}$ of every point in \mathcal{M} such that,

$$i(X_\xi)\Omega = -dJ_\xi. \tag{4}$$

This form J_ξ is determined up to some exact form $d\beta_\xi$ with $\beta_\xi \in \Omega^{m-2}(U)$ and coincides with the conserved quantity α_Y introduced in Remark 4 and Noether’s Theorem 2. Furthermore,

$$dJ_\xi = -i(X_\xi)\Omega_{\mathcal{S}} = i(X_\xi)d\Theta = L(X_\xi)\Theta - d i(X_\xi)\Theta, \tag{5}$$

and, as X_{ζ} is an infinitesimal Noether symmetry, then (3) holds for X_{ζ} , and $L(X_{\zeta})\Theta = d\zeta_{\zeta}$, for some $\zeta_{\zeta} \in \Omega^{m-1}(F)$. Now, by (5) it follows that

$$J_{\zeta} = -i(X_{\zeta})\Theta + \zeta_{\zeta} + d\beta_{\zeta}.$$

The $(m - 1)$ -form J_{ζ} can be taken to be linear in ζ by taking the generators $\zeta \in \mathfrak{g}$ to be infinitesimal. Therefore, for every point $p \in \mathcal{M}$, define the linear map

$$\begin{aligned} J|_p &: \mathfrak{g}^* \rightarrow \Omega^{m-1}(T_p\mathcal{M}) \\ \zeta &\mapsto J|_p(\zeta) := j_{\zeta}(p). \end{aligned}$$

Then:

Definition 5. *The map*

$$\begin{aligned} J &: \mathcal{M} \rightarrow \mathfrak{g}^* \otimes \Omega^{m-1}(F) \\ p &\mapsto J(p) := J|_p \end{aligned}$$

is called the **multimomentum map** associated with the symmetry group G .

The terminology “multimomentum map” is also used to refer the $(m - 1)$ -form J_{ζ} arising in (4). Furthermore, it is usual to specify the multimomentum map by using the natural pairing between \mathfrak{g}^* and \mathfrak{g} as $J_{\zeta}(p) = \langle J(p), \zeta \rangle$.

Thus, Noether symmetries on \mathcal{M} produce multimomentum maps on \mathcal{M} and the so-called **Noether current** $j = j^{\mu}d^{m-1}x_{\mu} \in \Omega^{m-1}(\mathcal{M})$ of a symmetry is obtained as $j = \psi^*J_{\zeta}$ (see also Remark 1).

3. Lifting Transformations from the Base Space of a Jet Bundle

In classical field theories, some symmetries are associated with diffeomorphisms on the base manifold M (i.e., spacetime transformations in most cases), and in the case of first-order Lagrangian field theories where \mathcal{M} is a first-order jet bundle, $J^1\pi \rightarrow E \rightarrow M$ (see Section 4.1), it is common to obtain the law of transformation in $J^1\pi$ from those diffeomorphisms on M .

This section discusses how to induce jet bundle transformations from diffeomorphisms on the base space. See [3,7] for other details.

3.1. First-Order Jet Bundles

Begin by letting $\pi: E \rightarrow M$ be a fiber bundle over an orientable m -dimensional manifold M with $\dim E = m + n$ and $m > 1$. The *first-order jet bundle* $J^1\pi$ of the projection π is the manifold of the 1-jets (equivalence classes) of local sections of π , denoted as $\phi \in \Gamma(\pi)$. Given a point $x \in M$, the points in $J^1\pi$ are denoted by $\bar{y} \equiv j_x^1\phi$, where $\phi \in \Gamma(\pi)$ is a representative of the equivalence class. The resulting natural projections are

$$\begin{aligned} \pi^1: J^1\pi &\rightarrow E & \bar{\pi}^1: J^1\pi &\rightarrow M \\ j_x^1\phi &\mapsto \phi(x) & j_x^1\phi &\mapsto x \end{aligned}$$

where $\bar{\pi}^1 = \pi \circ \pi^1$. The volume form on M as well as its pull-back $\bar{\pi}^{1*}\omega$ are denoted by ω . Local coordinates on $J^1\pi$ are denoted $(x^{\mu}, y^i, y_{\mu}^i)$, with $0 \leq \mu \leq m - 1, 1 \leq i \leq n$, and (x^{μ}, y^i) are local coordinates on E adapted to the bundle structure so that $\omega = dx^0 \wedge \dots \wedge dx^{m-1} \equiv d^m x$.

The *canonical lift* of a section $\phi \in \Gamma(\pi)$ to $J^1\pi$ is denoted $j^1\phi \in \Gamma(\bar{\pi}^1)$. A section $\psi \in \Gamma(\bar{\pi}^1)$ is said to be **holonomic** if ψ is the canonical lift of a section $\phi = \pi^1 \circ \psi \in \Gamma(\pi)$, and hence, $\psi = j^1(\pi^1 \circ \psi)$.

3.2. Lifting Transformations from M to E

Consider infinitesimal diffeomorphisms $\Phi_M: M \rightarrow M$ produced by the coordinate transformation $x'^\mu = x^\mu + \zeta^\mu(x)$. These transformations are generated by vector fields $Z = -\zeta^\mu(x) \frac{\partial}{\partial x^\mu} \in \mathfrak{X}(M)$ and the π -projectable vector fields generating the corresponding transformations on the configuration manifold E , which are written in full generality as $(x^\mu, y^A) \rightarrow (x^\mu + \zeta^\mu(x), y^i + \zeta^i(x, y))$ with $\zeta^i(x, y) \in C^\infty(E)$, are given as

$$Y \equiv Z_E = -\zeta^\mu(x) \frac{\partial}{\partial x^\mu} - \zeta^i(x, y) \frac{\partial}{\partial y^i} \in \mathfrak{X}(E). \tag{6}$$

The fields $y^i(x)$ which are given as components of the sections $\phi(x) = (x^\mu, y^i(x))$ transform as the Lie derivatives of the fields with respect to the vector field Z on M (denoted by $\delta y^i(x)$) as

$$\delta y^i(x) \equiv L(Z)y^i(x) = y'^i(x) - y^i(x) = -\zeta^\mu(x) \frac{\partial y^i}{\partial x^\mu}(x) + \tilde{\zeta}^i(x). \tag{7}$$

These field transformations are sometimes referred to as the **local variation** of the fields. The functions $\tilde{\zeta}^i(x)$ are not related to the components $\zeta^i(x, y)$ in (6), a priori. The term $-\zeta^\mu(x) \frac{\partial y^i}{\partial x^\mu}$ in (7) is called the **transport term**, and $\tilde{\zeta}^i(x)$ is called the **global variation** of the fields which is given by

$$\tilde{\zeta}^i(x) = y'^i(x') - y^i(x).$$

A geometrical interpretation of the local field variations is given as follows [16,34–36]:

Definition 6. Consider a section $\phi: M \rightarrow E$, and let $Z_E \in \mathfrak{X}(E)$ be a π -projectable vector field which projects to $Z \in \mathfrak{X}(M)$ (and hence has the local expression (6)). The **generalized Lie derivative of the section ϕ by Z** is the map $\mathbb{L}(Z)\phi: M \rightarrow TE$ defined as

$$\mathbb{L}(Z)\phi = T\phi \circ Z - Z_E \circ \phi, \tag{8}$$

which is a vector field along ϕ . The generalized Lie derivative has the form $\mathbb{L}(Z)\phi = (\phi, L(Z)\phi)$, and the section $L(Z)\phi: M \rightarrow V(T\pi)$ is called the **Lie derivative of the section ϕ by Z** and is given as $L(Z)\phi = (x^\mu, \zeta^\nu \frac{\partial \phi^i}{\partial x^\nu} - \zeta^i \circ \phi)$ when the local expression for Z_E is (6).

The definition of this Lie derivative must be in agreement with the Lie derivative of $y^i(x)$ defined by the field transformations $\delta y^i(x)$ in (7). This is achieved by setting $L(Z)\phi^i = L(Z)y^i(x)$ for the Lie derivative of the components ϕ^i in (8); then, it follows that the functions $\tilde{\zeta}^i(x)$ in (7) are given in terms of the component functions $\zeta^i(x, y)$ of Z_E and any local section ϕ as

$$\tilde{\zeta}^i(x) = \zeta^i(x, y) \circ \phi. \tag{9}$$

For theories in which $y^i(x)$ are scalar fields, diffeomorphisms on the base M produce field variations $\delta y^i(x)$ for which $\tilde{\zeta}^i(x) = 0 \implies \zeta^i(x, y) = 0$.

On the other hand, if the fields are, in general, tensor fields on M of type $(k, l) \neq (0, 0)$ (i.e., $T \in \mathfrak{T}^{(k,l)}(TM)$), then the functions $\tilde{\zeta}^i(x)$ are obtained, as usual, from the Jacobian (and its inverse) associated with the coordinate transformation $x^\mu \rightarrow x^\mu + \zeta^\mu(x)$. That is, if the fields $y^i(x)$ are the components of tensor fields $T \equiv (T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x))$, then

$$(T')_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x') = \left(\frac{\partial x'^{\mu_1}}{\partial x^{\alpha_1}} \right) \cdots \left(\frac{\partial x'^{\mu_r}}{\partial x^{\alpha_r}} \right) \left(\frac{\partial x^{\beta_1}}{\partial x'^{\nu_1}} \right) \cdots \left(\frac{\partial x^{\beta_s}}{\partial x'^{\nu_s}} \right) T_{\beta_1, \dots, \beta_s}^{\alpha_1, \dots, \alpha_r}(x). \tag{10}$$

The left-hand side of the expression above can be Taylor expanded around x giving the transport term as

$$(T')_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x') = (T')_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x + \xi) = (T')_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x) + \xi^\alpha \frac{\partial (T')_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x)}{\partial x^\alpha},$$

while performing the same Taylor expansion of the right-hand side of (10) for each Jacobian (and each inverse Jacobian) giving the global variation of the fields $\Delta T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x)$ given as

$$\Delta T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r} = \sum_{\mu=\mu_1}^{\mu_r} \left(\frac{\partial \xi^\mu}{\partial x^\lambda} \right) T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, (\mu \rightarrow \lambda), \dots, \mu_r} - \sum_{\nu=\nu_1}^{\nu_s} \left(\frac{\partial \xi^\lambda}{\partial x^\nu} \right) T_{\nu_1, \dots, (\nu \rightarrow \lambda), \dots, \nu_s}^{\mu_1, \dots, \mu_r}. \tag{11}$$

It follows that the local field variation given by the Lie derivative with respect to $Z = -\xi^\mu(x) \frac{\partial}{\partial x^\mu} \in \mathfrak{X}(M)$ is

$$\begin{aligned} \delta T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x) &= (T')_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x) - T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x) = L(Z)T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x) \\ &= -\xi^\lambda \frac{\partial (T')_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x)}{\partial x^\lambda} + \Delta T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x). \end{aligned}$$

Now, recalling relation (9), it follows that

$$Z_E = -\xi^\mu \frac{\partial}{\partial x^\mu} - \Delta_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x, T) \frac{\partial}{\partial T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}}, \tag{12}$$

where now,

$$\Delta T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x) = \Delta_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x, T) \circ \phi. \tag{13}$$

It is important to note that, when the configuration manifold E is a tensor bundle over M with coordinates $(x^\mu, T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r})$, then (12) is precisely the canonical lift of $Z \in \mathfrak{X}(M)$ to E which is defined as follows:

Definition 7.

1. Let $\Phi_M: M \rightarrow M$ be a diffeomorphism. The **canonical lift of Φ_M to E** is the diffeomorphism $\Phi_E: E \rightarrow E$ defined as follows: for every $(x, T_x) \in E$ where $T_x \in \mathfrak{T}^{(k,l)}(T_x M)$, define $\Phi_E(x, T_x) := (\Phi_M(x), \mathcal{T} \Phi_M(T_x))$, where $\mathcal{T} \Phi_M$ denotes the canonical transformation of tensors on M induced by Φ_M . Thus, $\pi \circ \Phi_E = \Phi_M \circ \pi$.
2. Let $Z \in \mathfrak{X}(M)$ be the vector field induced by local one-parameter groups of diffeomorphisms of M , denoted ϕ_t . The **canonical lift of Z to E** is the vector field $Z_E \in \mathfrak{X}(E)$ induced by local one-parameter groups of diffeomorphisms $(\phi_E)_t$ which are the canonical lifts of ϕ_t to the configuration bundle E .

In conclusion, the definition of the Lie derivative (8) of the local sections ϕ are constructed to always agree with the Lie derivatives of the fields (7), where Z_E in (6) is the canonical lift of Z to E , and hence, the relation (9) is always satisfied.

3.3. Lifting Transformations from E to $J^1\pi$

Let $\Phi: E \rightarrow E$ be a diffeomorphism which induces a diffeomorphism on the base space M as $\Phi_M: M \rightarrow M$ so $\Phi_M \circ \pi = \pi \circ \Phi$. Then, the **canonical lift** of Φ to $J^1\pi$ is the diffeomorphism $j^1\Phi: J^1\pi \rightarrow J^1\pi$ defined as

$$(j^1\Phi)(\bar{y}) := j^1(\Phi \circ \phi \circ \Phi_M^{-1})(\Phi_M(x)); \quad \text{for } \bar{y} \in J^1\pi.$$

Now, it is possible to define the canonical lift of π -projectable vector fields $Y \in \mathfrak{X}(E)$ to $J^1\pi$ for which there exist $Z \in \mathfrak{X}(M)$ such that the local flows of Z and Y are π^1 -related. The **canonical lift** of a π -projectable vector field $Y \in \mathfrak{X}(E)$ to $J^1\pi$ is the vector field

$j^1Y \in \mathfrak{X}(J^1\pi)$ whose local one-parameter groups of diffeomorphisms are the canonical lifts of the local one-parameter groups of diffeomorphisms of Y . If $Y \in \mathfrak{X}(E)$ is the canonical lift of $Z \in \mathfrak{X}(M)$ to E whose expression in local coordinates is given by (6) as $Y = Z_E = -\zeta^\mu(x) \frac{\partial}{\partial x^\mu} - \zeta^i(x, y) \frac{\partial}{\partial y^i}$, then the canonical lift of $Y \in \mathfrak{X}(E)$ to $J^1\pi$ is

$$j^1Z_E = -\zeta^\mu \frac{\partial}{\partial x^\mu} - \zeta^i \frac{\partial}{\partial y^i} - \left(\frac{\partial \zeta^i}{\partial x^\mu} - y^j \frac{\partial \zeta^i}{\partial x^\mu} + y^j \frac{\partial \zeta^i}{\partial y^j} \right) \frac{\partial}{\partial y^i_\mu} \in \mathfrak{X}(J^1\pi). \tag{14}$$

These canonical lifts are characterized by the property that they leave the canonical structures of the jet bundle $J^1\pi$ invariant: in particular, the *contact module* and, consequently, the *canonical endomorphism* under such canonical lifts. This canonical lift can be generalized for vector fields on E that are not π -projectable. If Y is not a π -projectable vector field, its canonical lift $j^1Y \in \mathfrak{X}(J^1\pi)$ is the only vector field that is π^1 -projectable to Y and leaves the canonical structures of $J^1\pi$ invariant (see [3,7] for details). In local coordinates, a vector field $Y = Z_E \in \mathfrak{X}(E)$ is not π -projectable if $\zeta^i \equiv \zeta^i(x, y)$. Then,

$$j^1Z_E = -\zeta^\mu \frac{\partial}{\partial x^\mu} - \zeta^i \frac{\partial}{\partial y^i} - \left(\frac{\partial \zeta^i}{\partial x^\mu} - y^j \left(\frac{\partial \zeta^i}{\partial x^\mu} + y^j \frac{\partial \zeta^i}{\partial y^j} \right) + y^j \frac{\partial \zeta^i}{\partial y^j} \right) \frac{\partial}{\partial y^i_\mu}.$$

In the same way, one can define the canonical lift of any diffeomorphism $\Phi: E \rightarrow E$ to $J^1\pi$.

The variation of the spacetime derivatives of the fields, $\delta y^i_\mu(x) = \delta \left(\frac{\partial y^i(x)}{\partial x^\mu} \right) = L(Z) \frac{\partial y^i(x)}{\partial x^\mu}$, can also be characterized by the generalized Lie derivative $L(Z)j^1\phi$ of the first-jet prolongations $j^1\phi: M \rightarrow J^1\pi: x^\mu \mapsto \left(x^\mu, y^i(x), \frac{\partial y^i(x)}{\partial x^\mu} \right)$ with respect to the vector field $Z = -\zeta^\mu(x) \frac{\partial}{\partial x^\mu} \in \mathfrak{X}(M)$, where now

$$\mathbb{L}(Z)j^1\phi = Tj^1\phi \circ Z - j^1Z \circ j^1\phi, \tag{15}$$

and $\mathbb{L}(Z)j^1\phi = (\phi, L(Z)\phi)$. It thereby follows that

$$L(Z)(j^1\phi)_\mu^i = \delta y^i_\mu(x) = -\zeta^\nu \frac{\partial y^i_\mu}{\partial x^\nu} - \frac{\partial \zeta^\nu}{\partial x^\mu} \frac{\partial y^i}{\partial x^\nu} + \frac{\partial \zeta^i}{\partial x^\mu} + \frac{\partial y^j}{\partial x^\mu} \frac{\partial \zeta^i}{\partial y^j}, \tag{16}$$

where now $\tilde{\zeta}^i(x) = \zeta^i(x, y) \circ j^1\phi$ as above. Furthermore, $\zeta^i = 0$ for scalar fields transformed under spacetime diffeomorphisms, and the equations above simplify it accordingly.

In the case where the fields are tensor fields $T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x)$, the canonical lift j^1Z_E in (14) of Z_E in (12) to $J^1\pi$ is

$$j^1Z_E = -\zeta^\mu \frac{\partial}{\partial x^\mu} - \Delta_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x, T) \frac{\partial}{\partial T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}} - \Gamma_{\alpha \nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r} \frac{\partial}{\partial T_{\alpha \nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}},$$

where the $\Delta_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r}(x, T) \in C^\infty(E)$ are again given by (11) and (13), while

$$\begin{aligned} \Gamma_{\alpha \nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r} &= \sum_{\mu=\mu_1}^{\mu_r} \frac{\partial \zeta^\mu}{\partial x^\lambda} \frac{\partial T_{\nu_1, \dots, \nu_s}^{\mu_1, \dots, (\mu \rightarrow \lambda), \dots, \mu_r}}{\partial x^\alpha} - \sum_{\nu=\nu_1}^{\nu_s} \frac{\partial \zeta^\lambda}{\partial x^\nu} \frac{\partial T_{\nu_1, \dots, (\nu \rightarrow \lambda), \dots, \nu_s}^{\mu_1, \dots, \mu_r}}{\partial x^\alpha} \\ &+ \sum_{\mu=\mu_1}^{\mu_r} \frac{\partial \zeta^\mu}{\partial x^\lambda} T^{\nu_1, \dots, \nu_s} - \sum_{\nu=\nu_1}^{\nu_s} \frac{\partial \zeta^\lambda}{\partial x^\nu} T_{\alpha \nu_1, \dots, (\nu \rightarrow \lambda), \dots, \nu_s}^{\mu_1, \dots, \mu_r} - T_{\beta \nu_1, \dots, \nu_s}^{\mu_1, \dots, \mu_r} \frac{\partial \zeta^\beta}{\partial x^\alpha}. \end{aligned}$$

4. Symmetries for Lagrangian and Hamiltonian Field Theories

The concepts presented in the previous sections will be developed in this section in the Lagrangian and De Donder–Weyl Hamiltonian formalisms of first-order field theories. See [4,9,13–15,17] for other details and considerations.

4.1. First-Order Lagrangian Field Theories

In this section, a comprehensive review of the geometric framework for first-order Lagrangian field theories is given. For more details, see [2–5,7,14,37].

The phase space for the Lagrangian formulation of first-order field theories is the so-called (*first-order*) *multivelocity phase space* which is the first-order jet bundle $J^1\pi$ of a bundle $\pi: E \rightarrow M$ called the *configuration bundle* of the theory, and the field states $y^i(x)$ are given by local sections $\phi: M \rightarrow E: x^\mu \mapsto (x^\mu, y^i(x))$ as in Section 3.

Each particular field theory is usually specified by a *Lagrangian*. However, the definition of the Lagrangian differs between the physics and differential geometry communities. In the differential geometry literature, the Lagrangian is defined as a C^∞ function on $J^1\pi$, while in the physics literature, the Lagrangian is defined as a functional on the space of *jet prolongations* $j^1\phi$ from M to $J^1\pi$. Both definitions can be used equivalently to formulate the variational principle which gives rise to the field equations of the theory under investigation. However, working on $J^1\pi$ has several geometric advantages over working on the space of jet prolongations. One advantage is that $J^1\pi$ is a finite dimensional manifold, while the space of jet prolongations, which is a space of local sections from M to $J^1\pi$, is an infinite-dimensional manifold.

Thus, a first-order field theory is described by a first-order Lagrangian density $\mathcal{L} \in \Omega^m(J^1\pi)$ which is a π^1 -semibasic m -form denoted as $\mathcal{L} = L \Omega \in \Omega^m(J^1\pi)$, and $L \in C^\infty(J^1\pi)$ is called the *Lagrangian function*. The Lagrangian phase space $J^1\pi$ possesses a canonical structure called the *canonical endomorphism* \mathcal{V} which is a $(1, 2)$ -tensor field whose local expression is given as $\mathcal{V} = (dy^i - y^i_\mu dx^\mu) \otimes \frac{\partial}{\partial y^i_\nu} \otimes \frac{\partial}{\partial x^\nu}$. Using this structure, the **Poincaré–Cartan** m and $(m + 1)$ -**forms** associated with \mathcal{L} are defined as $\Theta_{\mathcal{L}} := i(\mathcal{V})d\mathcal{L} + \mathcal{L} \in \Omega^m(J^1\pi)$ and $\Omega_{\mathcal{L}} := -d\Theta_{\mathcal{L}} \in \Omega^{m+1}(J^1\pi)$, respectively, and they have the following coordinate expressions:

$$\begin{aligned} \Theta_{\mathcal{L}} &= \frac{\partial L}{\partial y^i_\mu} dy^i \wedge d^{m-1}x_\mu - \left(\frac{\partial L}{\partial y^i_\mu} y^i_\mu - L \right) d^m x \equiv \frac{\partial L}{\partial y^i_\mu} dy^i \wedge d^{m-1}x_\mu - E_{\mathcal{L}} d^m x, \\ \Omega_{\mathcal{L}} &= -\frac{\partial^2 L}{\partial y^j_\nu \partial y^i_\mu} dy^j_\nu \wedge dy^i \wedge d^{m-1}x_\mu - \frac{\partial^2 L}{\partial y^j \partial y^i_\mu} dy^j \wedge dy^i \wedge d^{m-1}x_\mu \\ &\quad + \frac{\partial^2 L}{\partial y^j_\nu \partial y^i_\mu} y^i_\nu dy^j_\nu \wedge d^m x + \left(\frac{\partial^2 L}{\partial y^j \partial y^i_\mu} y^i_\mu - \frac{\partial L}{\partial y^j} + \frac{\partial^2 L}{\partial x^\mu \partial y^j_\mu} \right) dy^j \wedge d^m x. \end{aligned}$$

The couple $(J^1\pi, \Omega_{\mathcal{L}})$ is called a *first-order Lagrangian system*, and it is said to be *regular* if $\Omega_{\mathcal{L}}$ is 1-nondegenerate (that is, a *multisymplectic form*) and *singular* otherwise (i.e., $\Omega_{\mathcal{L}}$ is *premultisymplectic*). In the terminology of multisymplectic geometry, $\Theta_{\mathcal{L}}$ is said to be a (*pre*)*multisymplectic potential* of $\Omega_{\mathcal{L}}$. The regularity condition is locally equivalent to demanding that the generalized Hessian matrix $H^{i\mu}_{j\nu} = \frac{\partial^2 L}{\partial y^i_\mu \partial y^j_\nu}$ be non-singular everywhere on $J^1\pi$.

The solutions to the Lagrangian variational problem stated for a Lagrangian \mathcal{L} are holonomic sections $j^1\phi: M \rightarrow J^1\pi$ such that

$$(j^1\phi)^* i(X)\Omega_{\mathcal{L}} = 0, \quad \text{for every } X \in \mathfrak{X}(J^1\pi). \tag{17}$$

Equivalently, $j^1\phi$ are the integral sections of a class of locally decomposable and holonomic multivector fields $\{\mathbf{X}_{\mathcal{L}}\} \subset \mathfrak{X}^m(J^1\pi)$ which satisfy

$$i(\mathbf{X}_{\mathcal{L}})\Omega_{\mathcal{L}} = 0. \tag{18}$$

Furthermore, the holonomic multivector fields must be $\bar{\pi}^1$ -transverse,

$$i(\mathbf{X}_{\mathcal{L}})\omega \neq 0. \tag{19}$$

This condition can be fixed by taking a representative in the class $\{\mathbf{X}_{\mathcal{L}}\}$ such that $i(\mathbf{X}_{\mathcal{L}})\omega = 1$. The holonomic sections which solve the field equations (18) represent the physical states of the theory. In local coordinates, the field equations for sections are given as

$$\frac{\partial L}{\partial y^i} \circ j^1\phi - \frac{\partial}{\partial x^\mu} \left(\frac{\partial L}{\partial y_\mu^i} \circ j^1\phi \right) = 0,$$

which are the Euler–Lagrange equations of the system.

Remark 7. Equation (18) has a multiplicity of solutions (even in the regular case) [13,37]. This means that there is no unique class of integrable multivector fields (i.e., a unique distribution) which solve the field equations (18) on $J^1\pi$. Instead, there is a multiplicity of integral sections passing through every point in $J^1\pi$ which solve the field equations. If the Lagrangian is singular, there is another arbitrariness which comes from the degeneracy of the form $\Omega_{\mathcal{L}}$ and is related to the existence of geometric gauge symmetries. In general, when $(J^1\pi, \Omega_{\mathcal{L}})$ is a solvable singular Lagrangian system, the field equations admit stable solutions given by multivector fields which are locally decomposable and $\bar{\pi}^1$ -transverse on some $\bar{\pi}^1$ -transverse submanifold $S_f \subseteq J^1\pi$. Furthermore, these multivector fields are not always integrable (even for regular field theories). In addition, the multivector field solutions are not necessarily holonomic on all of S_f , but might be instead only on another $\bar{\pi}^1$ -transverse submanifold $\mathcal{S}_f \subseteq S_f$ to which these holonomic multivector fields solution to field equations must be tangent. This means that the image of the holonomic sections which solve the field equations (17) are on \mathcal{S}_f which is called the final constraint submanifold on which the field equations are said to have “consistent solutions”. The constraint algorithm needed to find \mathcal{S}_f is overviewed in Section 4.4. See [27,38] for a deeper analysis of these features.

4.2. De Donder–Weyl Hamiltonian Formalism

This section provides a review of the (pre)multisymplectic De Donder–Weyl formalism for classical field theories. See [6,39–41] and references therein for more details.

The De Donder–Weyl Hamiltonian formalism for regular Lagrangian (multisymplectic) field theories is performed on the so-called *multimomentum bundle* $J^{1*}\pi$ which is constructed as follows: consider the *extended multimomentum bundle* $\Lambda_2^m T^*E$ which is the bundle of m -forms on E vanishing by contraction with two π -vertical vector fields; then $J^{1*}\pi \equiv \Lambda_2^m T^*E / \Lambda_1^m T^*E$ (where $\Lambda_1^m T^*E$ is the bundle of π -semibasic m -forms on E). Natural coordinates on $\Lambda_2^m T^*E$ adapted to the bundle $\pi: E \rightarrow M$ are (x^ν, y^i, p_i^ν, p) (so $\dim \Lambda_2^m T^*E = nm + n + m + 1$), and natural coordinates on $J^{1*}\pi$ are (x^μ, y^i, p_i^μ) (so $\dim J^{1*}\pi = nm + n + m$). The natural projections for these bundles are

$$\bar{\tau}: J^{1*}\pi \rightarrow M, \quad \tau: J^{1*}\pi \rightarrow E, \quad \mathfrak{p}: \Lambda_2^m T^*E \rightarrow J^{1*}\pi.$$

The Legendre map $\mathcal{FL}: J^1\pi \rightarrow J^{1*}\pi$ associated with a Lagrangian function $L \in C^\infty(J^1\pi)$ is locally given by

$$\mathcal{FL}^*x^\nu = x^\nu, \quad \mathcal{FL}^*y^i = y^i, \quad \mathcal{FL}^*p_i^\nu = \frac{\partial L}{\partial y_i^\nu}.$$

The Lagrangian L is regular if, and only if, \mathcal{FL} is a local diffeomorphism; \mathcal{FL} is hyperregular when \mathcal{FL} is a global diffeomorphism. Then, there exist $\Theta_h \in \Omega^m(J^{1*}\pi)$ and

$\Omega_h := -d\tilde{\Theta}_h \in \Omega^{m+1}(J^{1*}\pi)$ such that $\mathcal{F}\mathcal{L}^*\Theta_h = \Theta_{\mathcal{L}}$ and $\mathcal{F}\mathcal{L}^*\Omega_h = \Omega_{\mathcal{L}}$ which are called the **Hamilton–Cartan m and $(m + 1)$ -forms**; Ω_h is the multisymplectic form on $J^{1*}\pi$, and the couple $(J^{1*}\pi, \Omega_h)$ is the *Hamiltonian system* associated with the (hyper)regular Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$. The local expressions for Θ_h and Ω_h are

$$\Theta_h = p_i^\mu dy^i \wedge d^{m-1}x_\mu - H d^m x, \quad \Omega_h = -dp_i^\mu \wedge dy^i \wedge d^{m-1}x_\mu + dH \wedge d^m x,$$

where

$$H = (\mathcal{F}\mathcal{L}^{-1})^*E_{\mathcal{L}} = p_i^\mu (\mathcal{F}\mathcal{L}^{-1})^*y_\mu^i - (\mathcal{F}\mathcal{L}^{-1})^*L \in C^\infty(J^{1*}\pi),$$

is the **De Donder–Weyl Hamiltonian function**. The field equations can be obtained from the so-called *Hamilton–Jacobi variational principle*, and their solutions are sections $\psi: M \rightarrow J^{1*}\pi$ which satisfy

$$\psi^* i(X)\Omega_h = 0, \quad \text{for every } X \in \mathfrak{X}(J^{1*}\pi).$$

Equivalently, such sections are integral sections of a class of integrable and $\bar{\tau}$ -transverse multivector fields $\{\mathbf{X}_h\} \subset \mathfrak{X}^m(J^{1*}\pi)$ satisfying

$$i(\mathbf{X}_h)\Omega_h = 0, \quad \text{for every } \mathbf{X}_h \in \{\mathbf{X}_h\}.$$

The $\bar{\tau}$ -transversality condition is fixed (as in the Lagrangian setting) by taking a representative in the class $\{\mathbf{X}_h\}$ such that $i(\mathbf{X}_h)w = 1$. Working with the natural local coordinates on $J^{1*}\pi$, the local sections are written as $\psi = (x^\mu, y^i(x^\nu), p_i^\mu(x^\nu))$ and satisfy

$$\frac{\partial(y^i \circ \psi)}{\partial x^\mu} = \frac{\partial H}{\partial p_i^\mu} \circ \psi, \quad \frac{\partial(p_i^\mu \circ \psi)}{\partial x^\mu} = -\frac{\partial H}{\partial y^i} \circ \psi.$$

which are the *Hamilton–De Donder–Weyl equations* of the system.

For singular Lagrangians, some minimal conditions must be imposed in order to ensure the existence of a De Donder–Weyl Hamiltonian description. In particular, one considers the so-called *almost regular Lagrangians* $L \in C^\infty(J^1\pi)$ which are those such that: (i) $P_0 := \mathcal{F}\mathcal{L}(J^1\pi)$ is a closed submanifold of $J^{1*}\pi$ which is called the *primary constraint submanifold*, (ii) $\mathcal{F}\mathcal{L}$ is a submersion onto its image, and (iii) for every $\bar{y} \in J^1\pi$, the fibers $\mathcal{F}\mathcal{L}^{-1}(\mathcal{F}\mathcal{L}(\bar{y}))$ are connected submanifolds of $J^1\pi$. Observe that, as $P_0 := \mathcal{F}\mathcal{L}(J^1\pi)$, it follows that $P_0 \rightarrow E \rightarrow M$. Then, denoting by $j_0: P_0 \hookrightarrow J^{1*}\pi$ the natural embedding of the primary constraint submanifold, the restriction $\mathcal{F}\mathcal{L}_0: J^1\pi \rightarrow P_0$ is the map defined by $\mathcal{F}\mathcal{L} = j_0 \circ \mathcal{F}\mathcal{L}_0$. Furthermore, there exist Hamilton–Cartan forms $\Theta_h^0 \in \Omega^m(P_0)$ and $\Omega_h^0 = -d\Theta_h^0 \in \Omega^{m+1}(\mathcal{P})$ such that $\Theta_{\mathcal{L}} = \mathcal{F}\mathcal{L}_0^* \Theta_h^0$ and $\Omega_{\mathcal{L}} = \mathcal{F}\mathcal{L}_0^* \Omega_h^0$ and also a *De Donder–Weyl Hamiltonian function* $H_0 \in C^\infty(P_0)$ such that $E_{\mathcal{L}} = \mathcal{F}\mathcal{L}_0^* H_0$. Therefore, the coordinate expression of the Hamilton–Cartan forms are

$$\Theta_h^0 = j_0^*(p_i^\mu dy^i \wedge d^{m-1}x_\mu) - H_0 d^m x, \quad \Omega_h^0 = j_0^*(-dp_i^\mu \wedge dy^i \wedge d^{m-1}x_\mu) + dH_0 \wedge d^m x.$$

In general, Ω_h^0 is a premultisymplectic form, and (P_0, Ω_h^0) is the *Hamiltonian system* associated with the almost regular Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$. The Hamilton–de Donder–Weyl equations are stated as in the regular case. When the non-regular Hamiltonian system admits stable solutions, such solutions to the Hamiltonian field equations exist only on a submanifold $P_f \subseteq P_0$ which is obtained by implementing the corresponding constraint algorithm described later in Section 4.4. The submanifold P_f is the *final constraint submanifold* in the De Donder–Weyl Hamiltonian formalism and which satisfies $\mathcal{F}\mathcal{L}(\mathcal{S}_f) = \mathcal{F}\mathcal{L}(S_f) = P_f \subset J^{1*}\pi$ since $\mathcal{S}_f \hookrightarrow S_f$ is defined on S_f by non- $\mathcal{F}\mathcal{L}$ -projectable constraints (see Section 4.4).

4.3. Symmetries, Conserved Quantities, and Multimomentum Maps

Recall that the Lagrangian formalism takes place on the bundle $J^1\pi \rightarrow E \rightarrow M$ both in the regular and singular cases. Then, all the definitions and results from Section 2 regarding

conserved quantities and symmetries apply directly. The concepts of *Lagrangian conserved quantities* (or *conservation laws*), *Lagrangian (infinitesimal) symmetries*, *Lagrangian (infinitesimal) Noether symmetries*, *Lagrangian gauge symmetries*, and *Lagrangian momentum maps* of a Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$ are also well-defined.

Nevertheless, in this situation, there are some new insights to be taken into account due to the fact that now, sections, diffeomorphisms and vector fields in $J^1\pi$ can be canonical lifts of sections, diffeomorphisms and vector fields in E , respectively. Therefore, we define:

Definition 8.

1. A (Noether) symmetry $\Phi: J^1\pi \rightarrow J^1\pi$ of a Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$ is said to be **natural** if Φ is a canonical lift; i.e., $\Phi = j^1\varphi$ for a diffeomorphism $\varphi: E \rightarrow E$.
2. An infinitesimal (Noether) symmetry $X \in \mathfrak{X}(J^1\pi)$ of a Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$ is said to be **natural** if X is a canonical lift; i.e., $X = j^1Z_E$ for some $Z_E \in \mathfrak{X}(E)$.

If $j^1\phi: M \rightarrow J^1\pi$ is a holonomic solution to the field equations (17) and $\Phi = j^1\varphi \in \text{Diff}(J^1\pi)$ is a natural Noether symmetry, then

$$\begin{aligned} (j^1(\varphi \circ \phi))^* i(X)\Omega_{\mathcal{L}} &= (j^k\phi)^*(j^k\varphi)^* i(X)\Omega_{\mathcal{L}} \\ &= (j^1\phi)^* i((j^1\varphi)_*^{-1}X)((j^1\varphi)^*\Omega_{\mathcal{L}}) \\ &= (j^1\phi)^* i(X')\Omega_{\mathcal{L}} = 0, \end{aligned} \tag{20}$$

since $X' = \Phi_*^{-1}X \in \mathfrak{X}(J^1\pi)$, $\Phi^*\Omega_{\mathcal{L}} = \Omega_{\mathcal{L}}$, and $\Phi^*\Omega_{\mathcal{L}} = 0$. Therefore, $j^1(\varphi \circ \phi)$ is also a holonomic solution to (17), and thus we have proved that:

Proposition 1. Every natural (infinitesimal) Noether symmetry transforms holonomic solutions to the Lagrangian field equations into equivalent holonomic solutions.

As stated for the general case, when $(J^1\pi, \Omega_{\mathcal{L}})$ is a singular Lagrangian system with final constraint submanifold $\mathcal{S}_f \hookrightarrow J^1\pi$, then (Noether) symmetries must be diffeomorphisms leaving \mathcal{S}_f invariant, and infinitesimal (Noether) symmetries are vector fields tangent to \mathcal{S}_f and satisfying the conditions of Definition 3 at least on \mathcal{S}_f .

Furthermore, as stated in Remarks 3 and 5, we will be especially interested in the case where (Noether) symmetries are fiber-preserving diffeomorphisms for the bundle $\pi^1: J^1\pi \rightarrow M$, and infinitesimal (Noether) symmetries are π^1 -projectable vector fields.

Similarly, for Lagrangian gauge symmetries we can define:

Definition 9. A geometric infinitesimal Lagrangian gauge symmetry (i.e., a Lagrangian gauge vector field) $X \in \mathfrak{X}(J^1\pi)$ of a Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$ is said to be **natural** if $X = j^1Z_E$ for some vector field $Z_E \in \mathfrak{X}(E)$.

As a consequence of Proposition 1, if $X \in \mathfrak{X}(J^1\pi)$ is a natural Lagrangian gauge vector field, then it transforms holonomic sections of the projection π^1 which are solutions to the Lagrangian field equations onto other holonomic sections solutions, all of which represent gauge equivalent physical states. Nevertheless, it should be noted that a holonomic section could be gauge equivalent to a non-holonomic one (both of them solutions to the field equations) when they are related by gauge transformations generated by gauge vector fields which are not natural.

In local coordinates, if X is a natural geometric gauge vector field, then the corresponding vector field $Z_E \in \mathfrak{X}(E)$ only has the component $Z_E = -\zeta^i(x, y) \frac{\partial}{\partial y^i} \in \mathfrak{X}(E)$, and it follows that the gauge vector field $X = j^1Z$ is given by

$$X = -\zeta^i \frac{\partial}{\partial y^i} - \left(\frac{\partial \zeta^i}{\partial x^\mu} + y^\mu \frac{\partial \zeta^i}{\partial y^j} \right) \frac{\partial}{\partial y^\mu} \in \mathfrak{X}(J^1\pi).$$

A particular type of symmetries in the Lagrangian formalism are the following:

Definition 10.

1. A **Lagrangian symmetry** of a Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$ is a diffeomorphism $\Phi: J^1\pi \rightarrow J^1\pi$ that leaves \mathcal{L} invariant: $\Phi^*\mathcal{L} = \mathcal{L}$.
If $\Phi = j^1\varphi$ for some fiber-preserving diffeomorphism $\varphi: E \rightarrow E$, then the Lagrangian symmetry is said to be **natural**.
2. An **infinitesimal Lagrangian symmetry** of a Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$ is a vector field $X \in \mathfrak{X}(J^1\pi)$ that leaves \mathcal{L} invariant.
If $X = j^1Z_E$, for some π -projectable vector field $Z_E \in \mathfrak{X}(E)$, then the infinitesimal Lagrangian symmetry is said to be **natural**.

Observe that given a diffeomorphism $\varphi: E \rightarrow E$ and $Z_E \in \mathfrak{X}(E)$, then

$$\begin{aligned} (j^1\varphi)^*\mathcal{L} = \mathcal{L} &\iff (j^1\varphi)^*\Theta_{\mathcal{L}} = \Theta_{\mathcal{L}}, \\ L(j^1Z_E)\mathcal{L} = 0 &\iff L(j^1Z_E)\Theta_{\mathcal{L}} = 0, \end{aligned}$$

and hence, (infinitesimal) Lagrangian symmetries are (infinitesimal) exact Noether symmetries.

Nevertheless, a generic (infinitesimal) Lagrangian symmetry does not necessarily leave the Poincaré–Cartan form $\Omega_{\mathcal{L}}$ invariant unless it is a natural Lagrangian symmetry. Likewise, as $\Omega_{\mathcal{L}}$ is not canonical (since it depends on the choice of the Lagrangian density \mathcal{L}), it is not invariant under canonical lifts of fiber-preserving diffeomorphisms and π -projectable vector fields unless some additional condition is assumed, such as the invariance of the Lagrangian density. In the spirit of this discussion, consider the following type of symmetries:

Definition 11.

1. A **geometric Lagrangian symmetry** of a Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$ is a diffeomorphism $\Phi: J^1\pi \rightarrow J^1\pi$ such that:
 - (a) $\Phi^*\mathcal{L} = \mathcal{L}$.
 - (b) The canonical geometric structures of $J^1\pi$ are invariant by Φ .
2. An **infinitesimal geometric Lagrangian symmetry** of a Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$ is a vector field $X \in \mathfrak{X}(J^1\pi)$ such that:
 - (a) $L(X)\mathcal{L} = 0$.
 - (b) The canonical geometric structures of $J^1\pi$ are invariant under the action of X .

It follows that (infinitesimal) natural Lagrangian symmetries are (infinitesimal) geometric Lagrangian symmetries, and (infinitesimal) geometric Lagrangian symmetries are (infinitesimal) exact Noether symmetries.

It is also worth noting that demanding the invariance of \mathcal{L} under some Lie group action on $J^1\pi$ is an exceedingly strong condition that is not necessary in order to obtain the same field equations from the variational principle. There are Lie group actions on $J^1\pi$ which produce so-called **gauge equivalent Lagrangians** which give rise to the same Euler–Lagrange equations (see, for instance, [12,42]). Gauge equivalent Lagrangian densities differ by an exact differential form: $\Phi^*\mathcal{L} = \mathcal{L} + d\beta$. Recall that in the physics literature, the analysis of field theories occurs on the jet prolongations $j^1\phi: M \rightarrow J^1\pi$ so that $\tilde{\mathcal{L}} = \tilde{L}\Omega = (j^1\phi)^*\mathcal{L} \in C^\infty(M)$. In this setting, gauge equivalent Lagrangians differ by a total derivative $\delta\tilde{L} = \partial_\mu\tilde{K}^\mu$ where $\tilde{K}^\mu = (j^1\phi)^*K^\mu$ for some K^μ . Furthermore, recall that when the field variations are produced by diffeomorphisms of M generated by some vector

field $\zeta \in \mathfrak{X}(M)$, the variation of the Lagrangian function \tilde{L} on $j^1\phi$ can be written as a Lie derivative of the local sections $j^1\phi$ with respect to ζ so that

$$\delta\tilde{L} = \frac{\partial\tilde{L}}{\partial y^i(x)}\delta y^i(x) + \frac{\partial\tilde{L}}{\partial y^i_\mu(x)}\delta y^i_\mu(x) \in C^\infty(M),$$

using Equations (8) and (16). Then, the corresponding Lie group action Φ on $J^1\pi$ is generated by the canonical lift $X_\zeta \in \mathfrak{X}(J^1\pi)$ of $\zeta \in \mathfrak{X}(M)$ to $J^1\pi$ given by (14). It follows that for $\Phi^*\mathcal{L} = \mathcal{L} + d\beta$,

$$d\beta = L(X_\zeta)\mathcal{L} = -d(K^\mu + L\zeta^\mu) \wedge d^{m-1}x_\mu \in \Omega^m(J^1\pi).$$

Similarly, when the field variations are gauge transformations, $\delta y^i(x) = \tilde{\zeta}^i(x) = \phi^*\zeta^i(x, y)$, it follows that

$$d\beta = L(X_\zeta)\mathcal{L} = -dK^\mu \wedge d^{m-1}x_\mu \in \Omega^m(J^1\pi),$$

where now X_ζ is given by (14) with $\zeta^\mu = 0$.

Finally, let G be a group of Noether symmetries for the Lagrangian system $(J^1\pi, \Omega_{\mathcal{L}})$, and let $J_\zeta \in \Omega^{m-1}(U)$, for $U \subset J^1\pi$ be the corresponding Lagrangian multimomentum map. Now, consider the situation in which the Noether symmetries are exact, and they are associated with diffeomorphisms on the base M , as described in Section 3.2. Then, denoting $X_\zeta \equiv j^1Z_E$, we have $L(j^1Z_E)\Theta_{\mathcal{L}} = 0$, and (modulo exact differential forms) the **Lagrangian multimomentum map** is given as

$$J_{\mathcal{L}}(X_\zeta) = -i(X_\zeta)\Theta_{\mathcal{L}} = \frac{\partial L}{\partial y^i_\mu}(\zeta^i d^{m-1}x_\mu - \zeta^\nu y^i_\mu d^{m-1}x_\nu - \zeta^\nu dy^i \wedge d^{m-2}x_{\mu\nu}) + L\zeta^\nu d^{m-1}x_\nu.$$

The corresponding Noether current is given by

$$\begin{aligned} j_\zeta^\mu d^{m-1}x_\mu &= (j^1\phi)^*[-i(X_\zeta)\Theta_{\mathcal{L}}] \\ &= \left[\left(\frac{\partial L}{\partial y^i_\mu} \circ j^1\phi \right) \left(\tilde{\zeta}^i - \zeta^\nu \frac{\partial y^i(x)}{\partial x^\nu} \right) + \zeta^\mu (L \circ j^1\phi) \eta^\mu_\nu \right] d^{m-1}x_\mu. \end{aligned}$$

The part of the Noether current shown above which corresponds to the infinitesimal spacetime transformations is linear in ζ , and the **canonical energy-momentum tensor**, denoted as usual as T^{μ}_ν , is defined from the terms contracted with ζ^ν above:

$$-\zeta^\nu \left[\left(\frac{\partial L}{\partial y^i_\mu} \circ j^1\phi \right) \frac{\partial y^i(x)}{\partial x^\nu} - (L \circ j^1\phi) \eta^\mu_\nu \right] d^{m-1}x_\mu \equiv -\zeta^\nu T^{\mu}_\nu d^{m-1}x_\mu.$$

Similarly to the Lagrangian formalism, all the definitions and results introduced in Section 2 apply straightforwardly to the De Donder–Weyl Hamiltonian setting by taking $\mathcal{M} \equiv J^{1*}\pi \rightarrow E \rightarrow M$ for regular field theories or $\mathcal{M} \equiv P_0 \rightarrow E \rightarrow M$ for almost regular field theories as discussed in the previous section. The concepts of *Hamiltonian conserved quantities* (or *conservation laws*), *Hamiltonian (infinitesimal) symmetries*, *Hamiltonian (infinitesimal) Noether symmetries*, *Hamiltonian gauge symmetries*, and *Hamiltonian multimomentum maps* are well-defined for regular and almost regular Hamiltonian systems $(J^{1*}\pi, \Omega_h)$ and (P_0, Ω_h^0) .

In particular, the **Hamiltonian multimomentum map** can be obtained by applying the push-forward of the Legendre map $\mathcal{F}L_*$ to $J_{\mathcal{L}}(X_\zeta)$ or, equivalently, using the $\mathcal{F}L$ -projection of X_ζ and contracting it with the corresponding (pre)multisymplectic form on the image of the Legendre map. When the field theory under investigation is regular, the calculation is straightforward, and the Hamiltonian multimomentum map is given as

$$\begin{aligned} \mathcal{F}\mathcal{L}_*(J_{\mathcal{L}}(X_{\xi})) &= J_h(\mathcal{F}\mathcal{L}_*X_{\xi}) = -i(\mathcal{F}\mathcal{L}_*X_{\xi})\Omega_h \\ &= p_i^{\mu}(\xi^i d^{m-1}x_{\mu} - \xi^{\nu} dy^i \wedge d^{m-2}x_{\mu\nu}) - H\xi^{\mu} d^{m-1}x_{\mu}, \end{aligned}$$

where H is the De Donder–Weyl Hamiltonian function. However, when the field theory is singular, important subtleties (which are detailed below) arise when projecting vector fields via $\mathcal{F}\mathcal{L}$.

4.4. Symmetries in the Presence of Constraints

The projection of Lagrangian symmetries to the Hamiltonian framework is straightforward when the Lagrangian is regular; that is, when the Legendre map $\mathcal{F}\mathcal{L}$ is a (local) diffeomorphism. Given a vector field $X \in \mathfrak{X}(J^1\pi)$ which generates a Lagrangian Noether symmetry on $J^1\pi$; that is, $L(X)\Omega_{\mathcal{L}} = 0$, the corresponding Hamiltonian symmetry on $J^{1*}\pi$ is generated by the vector field obtained from the push-forward by the Legendre map, $Y = \mathcal{F}\mathcal{L}_*X \in \mathfrak{X}(J^{1*}\pi)$; in fact,

$$0 = L(X)\Omega_{\mathcal{L}} = L(X)(\mathcal{F}\mathcal{L}^*\Omega_h) = \mathcal{F}\mathcal{L}_*[L(Y)\Omega_h] \iff L(Y)\Omega_h = 0,$$

where the push-forward by the Legendre map $\mathcal{F}\mathcal{L}_*$ is given by the matrix

$$\Gamma_{\mathcal{F}\mathcal{L}} \equiv \begin{pmatrix} (\text{Id})_{m \times m} & (0)_{m \times n} & (0)_{m \times nm} \\ (0)_{n \times m} & (\text{Id})_{n \times n} & (0)_{n \times nm} \\ (0)_{nm \times m} & \begin{pmatrix} \partial^2 \mathcal{L} \\ \partial y^B \partial y_{\mu}^A \end{pmatrix} & \begin{pmatrix} \partial^2 \mathcal{L} \\ \partial y^B \partial y_{\mu}^A \end{pmatrix} \end{pmatrix}.$$

When the Lagrangian is singular (in particular, almost regular) the primary constraint submanifold $P_0 = \text{Im}\mathcal{F}\mathcal{L} \subset J^{1*}\pi$, is defined by some independent functions $\varphi_I \in C^{\infty}(J^{1*}\pi)$. It follows that the null vectors of the Hessian matrix are given by [38]:

$$\left(\gamma_{\mu}^i\right)_I = \mathcal{F}\mathcal{L}_* \frac{\partial \varphi_I}{\partial p_{\mu}^i},$$

and since $\ker \mathcal{F}\mathcal{L}_* \subset \mathfrak{X}^{V(\pi)}(J^1\pi)$, the vector fields $\Gamma_I \in \ker \mathcal{F}\mathcal{L}_*$ can be written using a local basis for $\ker \mathcal{F}\mathcal{L}_*$ in natural coordinates on $J^1\pi$ as

$$\Gamma_I = \left(\gamma_{\mu}^i\right)_I \frac{\partial}{\partial y_{\mu}^i}.$$

Furthermore, since

$$\ker \mathcal{F}\mathcal{L}_* = \ker \Omega_{\mathcal{L}} \cap \mathfrak{X}^{V(\pi_1)}(J^1E) \subset \ker \Omega_{\mathcal{L}} \cap \mathfrak{X}^{V(\tilde{\pi}_1)}(J^1E) = \ker \Omega_{\mathcal{L}} \cap \ker \omega,$$

it follows that $\Gamma_I \in \ker \Omega_{\mathcal{L}} \cap \ker \omega$ and thereby generate geometric gauge symmetries.

In general, the Lagrangian and the Hamiltonian field equations have consistent solutions on the final constraint submanifolds $\mathcal{S}_f \subseteq J^1\pi$ and $P_f \subseteq P_0 \subset J^{1*}\pi$, respectively, where $\mathcal{F}\mathcal{L}(\mathcal{S}_f) = P_f$. Moreover, vector fields $X \in \mathfrak{X}(J^1\pi)$ which generate Noether symmetries on $J^1\pi$ must be tangent to \mathcal{S}_f in order to preserve the full constraint structure of the field theory under investigation; the tangency of X to \mathcal{S}_f is guaranteed by ensuring $L(X)\Omega_{\mathcal{L}} = 0$ (at least on \mathcal{S}_f).

If X is $\mathcal{F}\mathcal{L}$ -projectable from $J^1\pi$, then, proceeding similarly as in the regular case, the vector fields which generate Hamiltonian Noether symmetries on P_0 are given by $Y_0 = (\mathcal{F}\mathcal{L}_0)_*X \in \mathfrak{X}(P_0)$, and it follows that,

$$0 = L(X)\Omega_{\mathcal{L}} = L(X)(\mathcal{F}\mathcal{L}_0^*\Omega_h^0) = \mathcal{F}\mathcal{L}_0^*[L(Y_0)\Omega_h^0] \iff L(Y_0)\Omega_h^0 = 0,$$

this is the case, then the vector field produced by the push-forward of the Legendre map is a vector field on the corresponding Hamiltonian constraint submanifold P_i . Furthermore, a Noether symmetry may exist only on one of the Lagrangian constraint submanifolds S_i (or \mathcal{S}_i). If this is the case, then it follows that on the Hamiltonian side, the Noether symmetry exists only on the corresponding constraint submanifold P_i . The collection of different scenarios of how symmetries behave in the presence of premultisymplectic constraints are listed as follows:

1. There exists some $X \in \mathfrak{X}(J^1\pi)$ such that $L(X)\Omega_{\mathcal{L}} = 0$ and is \mathcal{FL} -projectable only from the constraint submanifold $\mathcal{S}_f \hookrightarrow J^1\pi$. Then, on the corresponding Hamiltonian constraint submanifold $P_f \subset P_0$, there exists the vector field $Y = \mathcal{FL}_*X|_{\mathcal{S}_f} \in \mathfrak{X}(P_f)$ such that $L(Y)\Omega_h^0|_{P_f} = 0$. Furthermore, the vector field $X \in \mathfrak{X}(J^1\pi)$ may or may not be the local extension of some $Z|_{\mathcal{S}_f}$ to $J^1\pi$ for some $Z \in \mathfrak{X}(J^1\pi)$.
2. There exists some $X \in \mathfrak{X}(\mathcal{S}_f)$ such that $L(X)\Omega_{\mathcal{L}}|_{\mathcal{S}_f} = 0$, and X is \mathcal{FL} -projectable only from \mathcal{S}_f . Then, on the corresponding Hamiltonian constraint submanifold $P_f \subset P_0$, there exists the vector field $Y|_{P_f} = \mathcal{FL}_*X|_{\mathcal{S}_f} \in \mathfrak{X}(P_f)|_{P_f}$ such that $L(Y)\Omega_h^0|_{P_f} = 0$. Furthermore, it is possible to construct a local extension of Y to P_0 , denoted as $\tilde{Y} \in \mathfrak{X}(P_0)$, but $X \in \mathfrak{X}(J^1\pi)$ may or may not be the local extension of some $Z|_{\mathcal{S}_f}$ to $J^1\pi$ for some $Z \in \mathfrak{X}(J^1\pi)$.

In the following sections, various field theories are worked out as examples whose symmetry structures in the presence of constraints exhibit special cases of the scenarios described above.

5. Some Examples

In the next sections, we summarize the main results about gauge and Noether symmetries and their associated conserved quantities (given by the corresponding multimomentum maps) of some different and known classical field theories in theoretical physics. For the details and calculations of the systems studied in Sections 5.1, 5.4 and 5.5 see [38].

5.1. Bosonic String Theories

Now spacetime M is a smooth $(d + 1)$ -dimensional manifold endowed with a spacetime metric $G_{\mu\nu}$ with signature $(- + \dots +)$. The string worldsheet Σ is a smooth two-dimensional manifold and has local coordinates σ^a with $a = 0, 1$. The fields $x^\mu(\sigma)$ are the scalar fields on Σ given by the embedding maps $X : \Sigma \rightarrow M : \sigma^a \mapsto x^\mu(\sigma)$ (see [44] for a thorough presentation and discussion on String Theory); so the configuration bundle E over Σ is $E = \Sigma \times M$, with natural projection $\pi : E \rightarrow \Sigma$ whose sections are $\phi : \Sigma \rightarrow \Sigma \times M : \sigma^a \mapsto (\sigma^a, x^\mu(\sigma))$. The Lagrangian and Hamiltonian phase bundles $J^1\pi$ and $J^1\pi^*$ have local coordinates $(\sigma^a, x^\mu, x_a^\mu)$ and $(\sigma^a, x^\mu, p_a^\mu)$, respectively, and the jet prolongations are $j^1\phi : \Sigma \times M \rightarrow J^1\pi : \sigma^a \mapsto (\sigma^a, x^\mu(\sigma), \frac{\partial x^\mu}{\partial \sigma^a}(\sigma))$.

The bosonic string theory is described by the standard *Nambu-Goto Lagrangian density*

$$\mathcal{L} = L(\sigma^a, x^\mu, x_a^\mu) d^2\sigma = -T \sqrt{-\det g} d^2\sigma = -T \sqrt{-\det(G_{\mu\nu} x_a^\mu x_b^\nu)} d^2\sigma,$$

where T is called the *string tension*. This Lagrangian is regular as can be seen directly from the regularity of the multi-Hessian matrix

$$\frac{\partial^2 L}{\partial x_a^\mu \partial x_b^\nu} = -T \sqrt{-\det g} \left[G_{\mu\nu} g^{ba} - G_{\mu\alpha} G_{\rho\nu} x_c^\alpha x_i^\rho (g^{ba} g^{ci} + g^{cb} g^{ai} - g^{ca} g^{bi}) \right], \tag{21}$$

where $g^{ba} \equiv (g^{-1})^{ba} = \frac{1}{\det g} \epsilon^{bc} \epsilon^{ad} g_{dc}$. The Lagrangian energy function and the Poincaré–Cartan forms on $J^1\pi$ are

$$\begin{aligned}
 E_{\mathcal{L}} &= \frac{\partial L}{\partial x_a^\mu} x_a^\mu - L = -T \sqrt{-\det g} (g^{ba} g_{ab} - 1) = -T \sqrt{-\det g}, \\
 \Theta_{\mathcal{L}} &= \frac{\partial L}{\partial x_a^\mu} dx^\mu \wedge d^1\sigma_a - E_{\mathcal{L}} \wedge d^2\sigma = -T \sqrt{-\det g} [G_{\mu\nu} g^{ba} x_b^\nu dx^\mu \wedge d^1\sigma_a - d^2\sigma], \\
 \Omega_{\mathcal{L}} &= T \left\{ \sqrt{-\det g} [G_{\mu\nu} g^{ba} - G_{\mu\alpha} G_{\beta\nu} x_c^\alpha x_i^\beta (g^{ba} g^{ci} + g^{cb} g^{ai} - g^{ca} g^{bi})] dx_b^\nu \wedge dx^\mu \wedge d^1\sigma_a \right. \\
 &\quad + \frac{\partial}{\partial x^\rho} (\sqrt{-\det g} G_{\mu\nu} g^{ba} x_b^\nu) dx^\rho \wedge dx^\mu \wedge d^1\sigma_a \\
 &\quad - \sqrt{-\det g} [G_{\mu\nu} g^{ba} - G_{\mu\alpha} G_{\beta\nu} x_c^\alpha x_i^\beta (g^{ba} g^{ci} + g^{cb} g^{ai} - g^{ca} g^{bi})] x_a^\mu dx_b^\nu \wedge d^2\sigma \\
 &\quad \left. - \left[\frac{\partial}{\partial \sigma^a} (\sqrt{-\det g} G_{\mu\nu} g^{ba} x_b^\nu) + \frac{1}{2} \sqrt{-\det g} g^{ba} x_a^\alpha x_b^\beta \partial_\mu G_{\alpha\beta} \right] dx^\mu \wedge d^2\sigma \right\}.
 \end{aligned}$$

Writing $h_{ab} = (j^1\phi)^* g_{ab}$, the resulting Euler–Lagrange equations are

$$\begin{aligned}
 0 &= -\sqrt{-\det h} h^{ba} \frac{\partial x^\alpha}{\partial \sigma^a} \frac{\partial x^\beta}{\partial \sigma^b} \frac{\partial G_{\alpha\beta}}{\partial x^\mu} + \frac{\partial}{\partial \sigma^a} \left(\sqrt{-\det h} G_{\mu\nu} h^{ba} \frac{\partial x^\nu}{\partial \sigma^b} \right) \\
 &\quad + \frac{\partial}{\partial x^\rho} \left(\sqrt{-\det h} G_{\mu\nu} h^{ba} \frac{\partial x^\nu}{\partial \sigma^b} \right) \frac{\partial x^\rho}{\partial \sigma^a} \\
 &\quad + \sqrt{-\det h} \left[G_{\mu\nu} h^{ba} - G_{\mu\alpha} G_{\beta\nu} \frac{\partial x^\alpha}{\partial \sigma^c} \frac{\partial x^\beta}{\partial \sigma^i} (h^{ba} h^{ci} + h^{cb} h^{ai} - h^{ca} h^{bi}) \right] \frac{\partial^2 x^\nu}{\partial \sigma^a \partial \sigma^b}.
 \end{aligned}$$

The Legendre map $\mathcal{FL} : J^1\pi \rightarrow J^1\pi^*$ gives

$$\mathcal{FL}^* \sigma^a = \sigma^a, \quad \mathcal{FL}^* x^\mu = x^\mu, \quad \mathcal{FL}^* p_\mu^a = -T \sqrt{-\det g} G_{\mu\nu} g^{ba} x_b^\nu,$$

which is invertible due to the regularity of the generalized Hessian (21). Using that

$$\begin{aligned}
 \Pi^{ab} &\equiv G^{\mu\nu} p_\mu^a p_\nu^b \Rightarrow \mathcal{FL}^* \Pi^{ab} = -T^2 \det g g^{ba} \\
 &\iff \\
 \mathcal{FL}^* \det \Pi &= (-T^2 \det g)^2 \det(g^{-1}) = T^4 \det g,
 \end{aligned}$$

and that $\Pi_{ab} = \frac{1}{\det \Pi} \epsilon_{cb} \epsilon_{da} \Pi^{cd}$, it follows that

$$(\mathcal{FL}^{-1})^* x_b^\nu = -\frac{1}{T} \sqrt{-\det \Pi} G^{\mu\nu} \Pi_{ab} p_\mu^a,$$

and therefore, the De Donder–Weyl Hamiltonian function is

$$H(\sigma^a, x^\mu, p_\mu^a) \equiv p_\mu^a (\mathcal{FL}^{-1})^* x_a^\mu - (\mathcal{FL}^{-1})^* L = -\frac{1}{T} \sqrt{-\det \Pi} \in C^\infty(J^1\pi^*).$$

Then, the Hamilton–Cartan forms on $J^{1*}\pi$ are given as

$$\begin{aligned}
 \Theta_h &= p_\mu^a dx^\mu \wedge d^1\sigma_a - H \wedge d^2\sigma = p_\mu^a dx^\mu \wedge d^1\sigma_a + \frac{1}{T} \sqrt{-\det \Pi} d^2\sigma, \\
 \Omega_h &= -dp_\mu^a \wedge dx^\mu \wedge d^1\sigma_a + dH \wedge d^2\sigma \\
 &= -dp_\mu^a \wedge dx^\mu \wedge d^1\sigma_a - \frac{\sqrt{-\det \Pi}}{T} \Pi_{ba} \left(\frac{1}{2} \partial_\mu G^{\rho\sigma} p_\rho^a p_\sigma^b dx^\mu + G^{\mu\nu} p_\nu^b dp_\mu^a \right) \wedge d^2\sigma.
 \end{aligned}$$

The field equations are the Hamilton–De Donder–Weyl equations given by

$$\frac{\partial p_\mu^a}{\partial \sigma^a} - \frac{\sqrt{-\det \Pi}}{2T} \Pi_{ba} \partial_\mu G^{\rho\sigma} p_\rho^a p_\sigma^b = 0, \quad \frac{\partial x^\mu}{\partial \sigma^a} + \frac{\sqrt{-\det \Pi}}{T} \Pi_{ba} G^{\mu\nu} p_\nu^b = 0.$$

The worldsheet diffeomorphisms are produced by transformations $\sigma^a + \zeta^a(\sigma)$; so the vector field $\zeta \in \mathfrak{X}(\Sigma)$ which generates the worldsheet diffeomorphisms is given by

$$\zeta = -\zeta^a \frac{\partial}{\partial \sigma^a} \in \mathfrak{X}(\Sigma).$$

Then, $\zeta \in \mathfrak{X}(\Sigma)$ can be lifted from Σ to the trivial bundle $E = \Sigma \times M$ and from E to $J^1\pi$ as

$$\zeta_E = -\zeta^a \frac{\partial}{\partial \sigma^a} \in \mathfrak{X}(E), \quad X_\zeta = -\zeta^a \frac{\partial}{\partial \sigma^a} + x_b^\mu \frac{\partial \zeta^b}{\partial \sigma^a} \frac{\partial}{\partial x_a^\mu} \in \mathfrak{X}(J^1\pi).$$

It follows that the field variation given by the generalized Lie derivative (8) of the local sections $\phi : \Sigma \rightarrow E$ with respect to ζ is $\delta X^\mu(\sigma) = -\zeta^a \frac{\partial X^\mu}{\partial \sigma^a}$. Furthermore,

$$L(X_\zeta)\mathcal{L} = 0 \Rightarrow L(X_\zeta)\Theta_{\mathcal{L}} = 0,$$

so the multimomentum map on $J^1\pi$ is

$$J_{\mathcal{L}}(X_\zeta) = -i(X_\zeta)\Theta_{\mathcal{L}} = -T\sqrt{-\det g} \left(\epsilon_{ac} \zeta^c G_{\mu\nu} g^{ba} x_b^\nu dx^\mu + \zeta^a d^1\sigma_a \right) \in \Omega^1(J^1\pi).$$

The vector field $Y_\zeta = \mathcal{F}\mathcal{L}_* X_\zeta \in \mathfrak{X}(J^1\pi^*)$ which generates the worldsheet diffeomorphisms on $J^{1*}\pi$ takes the form

$$Y_\zeta = -\zeta^a \frac{\partial}{\partial \sigma^a} - \frac{1}{T} \sqrt{-\det \Pi} G^{\mu\nu} \Pi_{bc} p_\nu^b \frac{\partial \zeta^c}{\partial \sigma_a} \frac{\partial}{\partial p_a^\mu},$$

and the corresponding multimomentum map is

$$J_h(Y_\zeta) = -i(Y_\zeta)\Theta_h = \epsilon_{ac} \zeta^c p_\mu^a dx^\mu - \frac{1}{T} \sqrt{-\det \Pi} \zeta^a d^1\sigma_a \in \Omega^1(J^{1*}\pi).$$

On the other hand, spacetime diffeomorphisms are generated by the vector field

$$\zeta = -\zeta^\mu \frac{\partial}{\partial x^\mu} \in (M).$$

whose canonical lift from M to E and from E to $J^1\pi$ are

$$\zeta_E = -\zeta^\mu \frac{\partial}{\partial x^\mu} \in \mathfrak{X}(E), \quad X_\zeta = -\zeta^\mu \frac{\partial}{\partial x^\mu} - x_a^\nu \partial_\nu \zeta^\mu \frac{\partial}{\partial x_a^\mu} \in \mathfrak{X}(J^1\pi).$$

Then, when ζ is a Killing vector field on M , it follows that

$$L(X_\zeta)\mathcal{L} = \frac{T}{2} \sqrt{-\det g} g^{ba} x_b^\nu x_a^\mu L(\zeta) G_{\mu\nu} d^2\sigma \zeta_E = -\zeta^\mu \frac{\partial}{\partial x^\mu} \in E = 0 \Rightarrow L(X_\zeta)\Theta_{\mathcal{L}} = 0,$$

and this means that spacetime isometries produce exact multisymplectomorphisms. So, the multimomentum map on $J^1\pi$ is

$$J_{\mathcal{L}}(X_\zeta) = -i(X_\zeta)\Theta_{\mathcal{L}} = T\sqrt{-\det g} G_{\mu\nu} G^{ba} x_b^\nu \zeta^\mu d^1\sigma_a \in \Omega^1(J^1\pi).$$

Furthermore, the exact multisymplectomorphisms of $J^{1*}\pi$ are generated by

$$Y_\zeta = \mathcal{F}\mathcal{L}^*X_\zeta = -\zeta^\mu \frac{\partial}{\partial x^\mu} + \frac{1}{T}\sqrt{-\det\Pi}G^{\mu\nu}\Pi_{ba}p_\mu^b \frac{\partial\zeta_\nu}{\partial x^\nu} \frac{\partial}{\partial p_\mu^a} \in \mathfrak{X}(J^{1*}\pi),$$

and the corresponding multimomentum map is given by

$$J_h(Y_\zeta) = -i(Y_\zeta)\Theta_h = -p_\mu^a \zeta^\mu d^1\sigma_a \in \Omega^1(J^{1*}\pi).$$

5.2. Yang–Mills Theory

The Yang–Mills theory with non-Abelian gauge group $SU(N)$ takes place on a four-dimensional Minkowski spacetime manifold M endowed with the Minkowski metric $\eta_{\mu\nu}$ of signature $(-+++)$. The gauge connection is denoted as $A_\mu = A_\mu^a t_a$, where t_a are the generators (in some matrix representation) of the corresponding Lie algebra $\mathfrak{g} = \mathfrak{su}(N)$. These generators satisfy $[t_b, t_c] = f_{bc}^a t_a$ ($a = 1, \dots, \dim\mathfrak{g}$), where f_{bc}^a are the structure constants. Then, $J^1\pi$ has local coordinates $(x^\mu, A_\nu^a, A_{\mu\nu}^a)$, while on $J^{1*}\pi$, they are $(x^\mu, A_\nu^a, \pi_a^{\mu\nu})$. The Lagrangian density is

$$\mathcal{L} = L d^4x = -\frac{1}{4}F_a^{\mu\nu}F_{\mu\nu}^a d^4x \in \Omega^4(J^1\pi),$$

where $F_{\mu\nu}^a = A_{\mu\nu}^a - A_{\nu\mu}^a - f_{bc}^a A_\mu^b A_\nu^c$ give the non-Abelian field strength curvature tensor when pulled back by $j^1\phi$. The Yang–Mills Lagrangian is singular as the multi-Hessian is singular:

$$\frac{\partial^2 L}{\partial A_{\mu\nu}^a \partial A_{\rho\sigma}^b} = \delta_{ab}(\eta^{\mu\sigma}\eta^{\nu\rho} - \eta^{\mu\rho}\eta^{\nu\sigma}).$$

Using the notation in which rank-2 tensor components split up into their symmetric part $V_{(\mu\nu)}$ and antisymmetric part $V_{[\mu\nu]}$ as $V_{\mu\nu} = V_{(\mu\nu)} + V_{[\mu\nu]}$, it is evident that the relevant null vectors of the Hessian matrix above form the set of symmetric spacetime matrices $\{V_{\mu\nu} = V_{(\mu\nu)}\}$.

The Lagrangian energy function $E_{\mathcal{L}} \in C^\infty(J^1\pi)$ and the Poincaré–Cartan forms on $J^1\pi$ are

$$\begin{aligned} E_{\mathcal{L}} &= -F_a^{\mu\nu}A_{\mu\nu}^a + \frac{1}{4}F_a^{\mu\nu}F_{\mu\nu}^a, \\ \Theta_{\mathcal{L}} &= -F_a^{\mu\nu}dA_\nu^a \wedge d^3x_\mu + \left(F_a^{\mu\nu}A_{\mu\nu}^a - \frac{1}{4}F_a^{\mu\nu}F_{\mu\nu}^a\right)d^4x, \\ \Omega_{\mathcal{L}} &= -(\eta^{\mu\sigma}\eta^{\nu\rho} - \eta^{\mu\rho}\eta^{\nu\sigma})\left(dA_{a\rho\sigma} + f_{abc}A_\sigma^b dA_\rho^c\right) \wedge dA_\nu^a \wedge d^3x_\mu - f_{bc}^a F_a^{\mu\nu}A_\mu^b dA_\nu^c \wedge d^4x \\ &\quad + (\eta^{\mu\sigma}\eta^{\nu\rho} - \eta^{\mu\rho}\eta^{\nu\sigma})A_a^{\rho\sigma}\left(dA_{\mu\nu}^a \wedge d^4x + f_{bc}^a A_\mu^b dA_\nu^c \wedge d^4x\right). \end{aligned}$$

The resulting field equations are the Euler–Lagrange equations,

$$D_{\mu b}^a F_a^{\mu\nu} = 0,$$

where $D_{\mu b}^a = \left(\frac{\partial}{\partial x^\mu} - f_{cb}^a A_\mu^c\right)$.

The Legendre map $\mathcal{F}\mathcal{L} : J^1\pi \rightarrow J^{1*}\pi$ is almost regular, and its image is the primary constraint submanifold $P_0 \subset J^{1*}\pi$ defined by

$$\mathcal{F}\mathcal{L}^*\pi_a^{\mu\nu} = -F_a^{\mu\nu}. \tag{23}$$

It follows that, as $F_a^{\mu\nu} = F_a^{[\mu\nu]}$, the image of $\mathcal{F}\mathcal{L}$ gives $\pi^{(\mu\nu)} = 0$. The Hamilton–Cartan forms are

$$\begin{aligned} \Theta_h^0 &= \pi_a^{[\mu\nu]} dA_\nu^a \wedge d^3x_\mu - \left(\frac{1}{2} f_{bc}^a \pi_a^{[\mu\nu]} A_\mu^b A_\nu^c - \frac{1}{4} \pi_a^{[\mu\nu]} \pi_a^{[\mu\nu]} \right) d^4x, \\ \Omega_h^0 &= -d\pi_a^{[\mu\nu]} \wedge dA_\nu^a \wedge d^3x_\mu + \left[\frac{1}{2} \left(f_{bc}^a A_\mu^b A_\nu^c - \pi_a^{[\mu\nu]} \right) d\pi_a^{[\mu\nu]} + f_{bc}^a \pi_a^{[\mu\nu]} A_\mu^b dA_\nu^c \right] \wedge d^4x. \end{aligned}$$

There are no Hamiltonian constraints other than the primary constraints (23), hence $P_f = P_0$. The (Hamilton)–De Donder–Weyl field equations for this theory are

$$D_{\mu b}^a \pi_a^{[\mu\nu]} = 0, \quad \frac{\partial A_\nu^a}{\partial x^\mu} + \frac{1}{2} \left(\pi_{\mu\nu}^a - f_{bc}^a A_\mu^b A_\nu^c \right) = 0.$$

The $SU(N)$ gauge fields transform as

$$\delta A_\mu = \frac{\partial \chi}{\partial x^\mu} + [A_\mu, \chi] \equiv D_\mu \chi, \quad \delta A_\mu^a = \delta_b^a \frac{\partial \chi^b}{\partial x^\mu} - f_{cb}^a \chi^c A_\mu^b \equiv D_{\mu b}^a \chi^b,$$

The vector field which generates the $SU(N)$ transformations on E is

$$X_\chi^E = -D_{\mu b}^a \chi^b \frac{\partial}{\partial A_\mu^a} \in \mathfrak{X}(E),$$

the canonical lift of X_χ^E to $J^1\pi$ is given by

$$X_\chi = j^1 X_\chi^E = -D_{\mu b}^a \chi^b \frac{\partial}{\partial A_\mu^a} - \left(\frac{\partial D_{\nu b}^a \chi^b}{\partial x^\mu} - f_{cb}^a A_{\mu\nu}^c \chi^b \right) \frac{\partial}{\partial A_{\mu\nu}^a} \in \mathfrak{X}(J^1\pi),$$

and the $\mathcal{F}\mathcal{L}$ -projection onto $P_0 \subset J^{1*}\pi$, $Y_\chi = \mathcal{F}\mathcal{L}_* X_\chi \in \mathfrak{X}(P_0)$, is

$$Y_\chi = -D_{\mu b}^a \chi^b \frac{\partial}{\partial A_\mu^a} - \left[\frac{\partial D_{\nu b}^a \chi^b}{\partial x^\mu} + \frac{1}{2} f_{cb}^a \chi^c \left(\pi_{[\mu\nu]}^c - f_{de}^c A_\mu^d A_\nu^e \right) \right] \left(\delta^{\mu\beta} \delta^{\nu\alpha} - \delta^{\mu\alpha} \delta^{\nu\beta} \right) \frac{\partial}{\partial \pi_a^{[\alpha\beta]}}.$$

The corresponding multimomentum maps are

$$\begin{aligned} J_{\mathcal{L}}(X_\chi) &= -i(X_\chi)\Theta_{\mathcal{L}} = -F_a^{\mu\nu} D_{\nu b}^a \chi^b d^3x_\mu \in \Omega^3(J^1\pi), \\ J_h^0(Y_\chi) &= -i(Y_\chi)\Theta_h^0 = \pi_a^{[\mu\nu]} D_{\nu b}^a \chi^b d^3x_\mu \in \Omega^3(P_0). \end{aligned}$$

Yang–Mills also exhibits spacetime isometries as a Noether symmetry. The infinitesimal version of this symmetry begins by lifting Killing vectors, in this case of the Minkowski metric, $\zeta = -\zeta^\mu(x) \frac{\partial}{\partial x^\mu} \in \mathfrak{X}(M)$, to the configuration bundle,

$$\zeta_E = -\zeta^\mu(x) \frac{\partial}{\partial x^\mu} + A_\nu^a \frac{\partial \zeta^\nu}{\partial x^\mu} \frac{\partial}{\partial A_\mu^a} \in \mathfrak{X}(E).$$

It follows that the canonical lift to the multivelocity phase space $J^1\pi$ is given as

$$X_\zeta = j^1 \zeta_E = -\zeta^\mu(x) \frac{\partial}{\partial x^\mu} + A_\nu^a \frac{\partial \zeta^\nu}{\partial x^\mu} \frac{\partial}{\partial A_\mu^a} + \left(A_{\nu\mu}^a \frac{\partial \zeta^\nu}{\partial x^\rho} + A_{\rho\nu}^a \frac{\partial \zeta^\nu}{\partial x^\mu} \right) \frac{\partial}{\partial A_{\rho\mu}^a},$$

and the corresponding multimomentum map on $J^1\pi$ is

$$\begin{aligned} J_{\mathcal{L}}(X_\zeta) &= -i(X_\zeta)\Theta_{\mathcal{L}} \\ &= F_a^{\mu\nu} A_\rho^a \frac{\partial \zeta^\rho}{\partial x^\nu} d^3x_\mu + F_a^{\mu\nu} \zeta^\rho dA_\nu^a \wedge d^2x_{\mu\rho} + \left(F_a^{\mu\nu} A_{\mu\nu}^a - \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu} \right) \zeta^\rho d^3x_\rho. \end{aligned}$$

The \mathcal{FL} projection of X_{ξ} is

$$Y_{\xi} = \mathcal{FL}_* X_{\xi} = -\xi^{\mu}(x) \frac{\partial}{\partial x^{\mu}} + A_{\nu}^a \frac{\partial \xi^{\nu}}{\partial x^{\mu}} \frac{\partial}{\partial A_{\mu}^a} + \eta^{\alpha\mu} \eta^{\beta\rho} \left[\left(\pi_{b[\mu\nu]} - f_{bcd} A_{\mu}^c A_{\nu}^d \right) \frac{\partial \xi^{\nu}}{\partial x^{\rho}} + \left(\pi_{b[\nu\rho]} - f_{bcd} A_{\nu}^c A_{\rho}^d \right) \frac{\partial \xi^{\nu}}{\partial x^{\mu}} \right] \frac{\partial}{\partial \pi_b^{[\alpha\beta]}}$$

and the resulting multimomentum map on $P_0 \subset J^1\pi^*$ is

$$J_{\mu}^0(Y_{\xi}) = -\pi_a^{[\mu\nu]} A_{\rho}^a \frac{\partial \xi^{\rho}}{\partial x^{\nu}} d^3 x_{\mu} - \pi_a^{[\mu\nu]} \xi^{\rho} dA_{\nu}^a \wedge d^2 x_{\mu\rho} - \left(\pi_a^{[\mu\nu]} A_{\mu\nu}^a + \frac{1}{4} \pi_{[\mu\nu]}^a \pi_a^{[\mu\nu]} \right) \xi^{\rho} d^3 x_{\rho}.$$

5.3. Chern–Simons Theory

For the multisymplectic treatment of the Chern–Simons theory with the Abelian gauge group $U(1)$, see [16]. Consider Chern–Simons theory in $2 + 1$ dimensions with gauge group $G = SU(N)$, $\dim G = N^2 - 1$. The gauge connection is denoted as $A_{\mu} = A_{\mu}^a t_a$, where t_a are the generators (in some matrix representation) of the corresponding Lie algebra $\mathfrak{g} = \mathfrak{su}(N)$. The generators satisfy $[t_b, t_c] = f_{bc}^a t_a$, $a = 1, \dots, \dim G$, where f_{bc}^a are structure constants. Then, $J^1\pi$ has local coordinates $(x^{\mu}, A_{\nu}^a, A_{\mu\nu}^b)$. Now, let $A = A_{\mu} dx^{\mu} \in \Omega^1(J^1\pi)$ and $F = \frac{1}{2} F_{\mu\nu} dx^{\mu} \wedge dx^{\nu} \in \Omega^2(J^1\pi)$, with $F_{\mu\nu} \equiv 2A_{[\mu\nu]} + \frac{1}{2}[A_{\mu}, A_{\nu}]$; so the Chern–Simons Lagrangian density is

$$\begin{aligned} \mathcal{L} &= L(x^{\mu}, A_{\nu}, A_{\mu\nu}) d^3 x = \text{Tr} \left[F \wedge A - \frac{1}{3} A \wedge A \wedge A \right] = \epsilon^{\mu\nu\rho} \text{Tr} \left[A_{\mu\nu} A_{\rho} + \frac{2}{3} A_{\mu} A_{\nu} A_{\rho} \right] d^3 x \\ &= \epsilon^{\mu\nu\rho} g_{ab} \left(-\frac{1}{4} F_{\mu\nu}^a A_{\rho}^b + \frac{1}{6} f_{cd}^b A_{\mu}^c A_{\nu}^d A_{\rho}^a \right) d^3 x = -\frac{1}{2} \epsilon^{\mu\nu\rho} g_{ab} \left(A_{\mu\nu}^a A_{\rho}^b + \frac{2}{3} f_{cd}^b A_{\mu}^c A_{\nu}^d A_{\rho}^a \right) d^3 x, \end{aligned}$$

where the trace shown above is ad-invariant on the Lie algebra and can thereby be taken using the Cartan–Killing metric $g_{ab} = -2\text{Tr}(t_a t_b)$ on \mathfrak{g} . This theory is singular as the Hessian is

$$\frac{\partial^2 L}{\partial A_{\mu\nu}^a \partial A_{\rho\sigma}^b} = 0.$$

The Lagrangian energy and the Poincaré–Cartan forms are given by

$$\begin{aligned} E_{\mathcal{L}} &\equiv \frac{\partial L}{\partial A_{\mu\nu}^a} A_{\mu\nu}^a - L = \frac{1}{3} \epsilon^{\mu\nu\rho} g_{ab} f_{cd}^b A_{\mu}^c A_{\nu}^d A_{\rho}^a \in C^{\infty}(J^1\pi), \\ \Theta_{\mathcal{L}} &= -\epsilon^{\mu\nu\rho} g_{ab} \left[\frac{1}{2} A_{\rho}^b dA_{\nu}^a \wedge d^2 x_{\mu} + \frac{1}{3} f_{cd}^b A_{\mu}^c A_{\nu}^d A_{\rho}^a d^3 x \right], \\ \Omega_{\mathcal{L}} &= \epsilon^{\mu\nu\rho} g_{ab} \left[\frac{1}{2} dA_{\rho}^b \wedge dA_{\nu}^a \wedge d^2 x_{\mu} + f_{cd}^b A_{\mu}^c A_{\nu}^d dA_{\rho}^a \wedge d^3 x \right]. \end{aligned}$$

The Lagrangian constraint in this theory arises from imposing the SOPDE condition on the multivector field solutions to the Lagrangian field equations, giving the following constraint

$$\epsilon^{\mu\nu\rho} \left(g_{ab} A_{\mu\nu}^b + g_{db} f_{ca}^b A_{\mu}^c A_{\nu}^d \right) = 0,$$

which defines the submanifold $\mathcal{S}_f \subset J^1\pi$. The tangency condition on this submanifold gives no new constraints.

The multimomentum phase space $J^1\pi^*$ has local coordinates $(x^{\mu}, A_{\nu}^a, \pi_a^{\mu\nu})$, and the Legendre map $\mathcal{FL} : J^1\pi \rightarrow J^1\pi^* : (x^{\mu}, A_{\nu}^a, A_{\mu\nu}^b) \mapsto (x^{\mu}, A_{\nu}^a, \pi_a^{\mu\nu})$ satisfies

$$\mathcal{FL}^* \pi_a^{\mu\nu} = \frac{\partial \mathcal{L}}{\partial A_{\mu\nu}^a} = -\frac{1}{2} \epsilon^{\mu\nu\rho} g_{ab} A_{\rho}^b,$$

which means that the image of \mathcal{FL} is the submanifold $P_0 \subset J^{1*}\pi$ defined by the constraints

$$\pi_a^{\mu\nu} + \frac{1}{2}\epsilon^{\mu\nu\rho}g_{ab}A_\rho^b = 0.$$

Thus, local coordinates on P_0 are (x^μ, A_ν^a) . The De Donder–Weyl Hamiltonian function is

$$H_0(x^\mu, A_\nu) = \frac{1}{3}\epsilon^{\mu\nu\rho}g_{ab}f_{cd}^b A_\mu^a A_\nu^c A_\rho^d \in C^\infty(P_0),$$

and the Hamilton–Cartan forms on P_0 are

$$\begin{aligned} \Theta_h^0 &= -\epsilon^{\mu\nu\rho}g_{ab} \left[\frac{1}{2}A_\rho^b dA_\nu^a \wedge d^2x_\mu + \frac{1}{3}f_{cd}^b A_\mu^a A_\nu^c A_\rho^d d^3x \right], \\ \Omega_h^0 &= \epsilon^{\mu\nu\rho}g_{ab} \left[\frac{1}{2}dA_\rho^b \wedge dA_\nu^a \wedge d^2x_\mu + f_{cd}^b A_\mu^a A_\nu^c dA_\rho^d \wedge d^3x \right]. \end{aligned}$$

There are no further Hamiltonian constraints other than the primary constraints and $\mathcal{FL}(\mathcal{S}_f) = P_0$. The field equations in both the Langrangian and Hamiltonian settings are

$$\epsilon^{\mu\nu\rho}g_{ab} \left(\partial_\mu A_\nu^b + f_{dc}^b A_\mu^d A_\nu^c \right) = 0 \Rightarrow F_{\mu\nu}^b = 0 \Rightarrow F_{\mu\nu} = 0.$$

The gauge symmetry group $SU(N)$ of the Chern–Simons theory produces infinitesimal transformations

$$\delta A_\mu = D_\mu \chi = \partial_\mu \chi + [A_\mu, \chi] \quad , \quad \delta A_\mu^a = D_{\mu b}^a \chi^b = \delta_{cb}^a \partial_\mu \chi^b - f_{cb}^a \chi^b A_\mu^c,$$

where $\chi = \chi^a t_a \in C^\infty(M)$. The vector field which generates the $SU(N)$ transformations on E is

$$X_\chi^E = -D_{\mu b}^a \chi^b \frac{\partial}{\partial A_\mu^a} \in \mathfrak{X}(E),$$

whose canonical lift to $J^1\pi$ is

$$X_\chi = j^1 X_\chi^E = -D_{\mu b}^a \chi^b \frac{\partial}{\partial A_\mu^a} - \left(\frac{\partial D_{\nu b}^a \chi^b}{\partial x^\mu} - f_{cb}^a A_{\mu\nu}^c \chi^b \right) \frac{\partial}{\partial A_{\mu\nu}^a} \in \mathfrak{X}(J^1\pi),$$

and its \mathcal{FL} -projection onto $P_0 \subset J^{1*}\pi$ is

$$Y_\chi = \mathcal{FL}_* X_\chi = -D_{\mu b}^a \chi^b \frac{\partial}{\partial A_\mu^a} \in \mathfrak{X}(P_0).$$

Finally, the corresponding multimomentum maps are

$$\begin{aligned} J_{\mathcal{L}}(X_\chi) &= -i(X_\chi)\Theta_{\mathcal{L}} = -\frac{1}{2}\epsilon^{\mu\nu\rho}g_{ab}A_\rho^b D_{\nu b}^a \chi^b d^2x_\mu \in \Omega^2(J^1\pi), \\ J_h^0(Y_\chi) &= -i(Y_\chi)\Theta_h^0 = -\frac{1}{2}\epsilon^{\mu\nu\rho}g_{ab}A_\rho^b D_{\nu b}^a \chi^b d^2x_\mu \in \Omega^2(J^1\pi) \in \Omega^2(P_0). \end{aligned}$$

Exactly as in the Yang–Mills theory, spacetime diffeomorphisms which are isometries are a Noether symmetry generated by the Killing vector field $\zeta = -\zeta^\mu(x) \frac{\partial}{\partial x^\mu} \in \mathfrak{X}(M)$. The canonical lift to the configuration bundle E is

$$\zeta_E = -\zeta^\mu(x) \frac{\partial}{\partial x^\mu} + A_\nu^a \frac{\partial \zeta^\nu}{\partial x^\mu} \frac{\partial}{\partial A_\nu^a} \in \mathfrak{X}(E),$$

which can be lifted again to $J^1\pi$, giving

$$X_{\xi} = j^1\xi_E = -\xi^\mu(x) \frac{\partial}{\partial x^\mu} + A_\nu^a \frac{\partial \xi^\nu}{\partial x^\mu} \frac{\partial}{\partial A_\mu^a} + \left(A_{\nu\mu}^a \frac{\partial \xi^\nu}{\partial x^\rho} + A_{\rho\nu}^a \frac{\partial \xi^\nu}{\partial x^\mu} \right) \frac{\partial}{\partial A_{\rho\mu}^a} \in \mathfrak{X}(J^1\pi),$$

as before. The corresponding multimomentum map $J_{\mathcal{L}}(X_{\xi}) \in \Omega^2(J^1\pi)$ is

$$\begin{aligned} J_{\mathcal{L}}(X_{\xi}) &= -i(X_{\xi})\Theta_{\mathcal{L}} \\ &= \epsilon^{\mu\nu\rho} g_{ab} \left[\frac{1}{2} A_\rho^b \left(A_\sigma^a \frac{\partial \xi^\sigma}{\partial x^\nu} d^2x_\mu + \xi^\sigma dA_\nu^a \wedge d^1x_{\mu\sigma} \right) - \frac{1}{3} f_{cd}^b \xi^\sigma A_\mu^a A_\nu^c A_\rho^d d^2x_\sigma \right]. \end{aligned}$$

Furthermore, the projection of $X_{\xi} \in \mathfrak{X}(J^1\pi)$ to $P_0 \subset J^1\pi^*$ is

$$Y_{\xi} = \mathcal{F}\mathcal{L}_* X_{\xi} = -\xi^\mu(x) \frac{\partial}{\partial x^\mu} + A_\nu^a \frac{\partial \xi^\nu}{\partial x^\mu} \frac{\partial}{\partial A_\mu^a} \in \mathfrak{X}(P_0).$$

The resulting multimomentum map $J_h^0(Y_{\xi}) \in \Omega^2(P_0)$ is

$$\begin{aligned} J_h^0(Y_{\xi}) &= -i(Y_{\xi})\Theta_h^0 \\ &= \epsilon^{\mu\nu\rho} g_{ab} \left[\frac{1}{2} A_\rho^b \left(A_\sigma^a \frac{\partial \xi^\sigma}{\partial x^\nu} d^2x_\mu + \xi^\sigma dA_\nu^a \wedge d^1x_{\mu\sigma} \right) - \frac{1}{3} f_{cd}^b \xi^\sigma A_\mu^a A_\nu^c A_\rho^d d^2x_\sigma \right]. \end{aligned}$$

5.4. Electric Carrollian Scalar Field Theory

Consider two scalar fields $\phi(x)$ and $\pi(x)$ which are given as the local sections $\phi : M \rightarrow E : x^\mu \mapsto (x^\mu, \phi(x), \pi(x))$ of a configuration bundle $\pi : E \rightarrow M : (\phi, \pi) \mapsto x^\mu$, where M is an m -dimensional spacetime manifold with Minkowski metric $\eta_{\mu\nu}$. Now, take the electric Carrollian contraction of the following Lagrangian,

$$L = \pi\phi_0 - \frac{1}{2}\pi^2 - \frac{1}{2}\phi_i\phi^i \in C^\infty(J^1\pi), \tag{27}$$

by making the field redefinition $\phi(x) \rightarrow c\phi(x)$, $\pi(x) \rightarrow \frac{1}{c}\pi(x)$, and taking the limit $c \rightarrow 0$ for the speed of light. It follows that the Minkowski metric becomes degenerate:

$$ds^2 = \eta_{\mu\nu} dx^\mu \otimes dx^\nu = -c^2 dx^0 \otimes dx^0 + \delta_{ij} dx^i \otimes dx^j \longrightarrow \delta_{ij} dx^i \otimes dx^j. \tag{28}$$

The Lagrangian function $L \in C^\infty(J^1\pi)$ which is obtained from the electric Carrollian contraction of (27) is

$$L = \pi\phi_0 - \frac{1}{2}\pi^2,$$

which is the electric Carrollian scalar field Lagrangian studied in this section. This Lagrangian is singular since $\frac{\partial^2 L}{\partial \phi_\mu \partial \phi_\nu} = \frac{\partial^2 L}{\partial \pi_\mu \partial \pi_\nu} = \frac{\partial^2 L}{\partial \pi_\mu \partial \phi_\nu} = 0$. The Lagrangian energy function and the Poincaré–Cartan forms are

$$\begin{aligned} E_{\mathcal{L}} &= \frac{1}{2}\pi^2 \in C^\infty(J^1\pi), \\ \Theta_{\mathcal{L}} &= \pi d\phi \wedge d^{m-1}x_0 - \frac{1}{2}\pi^2 d^m x \in \Omega^m(J^1\pi), \\ \Omega_{\mathcal{L}} &= d\phi \wedge d\pi \wedge d^{n-1}x_0 + \pi d\pi \wedge d^m x \in \Omega^{m+1}(J^1\pi). \end{aligned}$$

Again, the compatibility of the Lagrangian field equations produce two SOPDE constraints

$$\pi_0 = 0, \quad \phi_0 - \pi = 0,$$

which define the constraint submanifold $\mathcal{S}_f \subset J^1\pi$, and the tangency condition on this submanifold gives no new constraints.

The Legendre map is

$$\mathcal{FL}^* p_\phi^0 \equiv \frac{\partial L}{\partial \phi_0} = \pi \quad , \quad \mathcal{FL}^* p_\phi^i = \frac{\partial L}{\partial \phi_i} = 0 \quad , \quad \mathcal{FL}^* p_\pi^\mu = \frac{\partial L}{\partial \pi_\mu} = 0 \quad ,$$

and gives the primary constraints

$$p_\phi^0 - \pi = 0, \quad p_\phi^i = 0 \quad , \quad p_\pi^\mu = 0 \quad ,$$

which define the submanifold $P_0 \subset J^1\pi^*$ with local coordinates (x^μ, ϕ, π) . The Lagrangian function L is almost regular, and the De Donder–Weyl Hamiltonian is

$$H_0 = \frac{1}{2}\pi^2 \in C^\infty(P_0) \quad ,$$

and the Hamilton-Cartan forms are

$$\begin{aligned} \Theta_h^0 &= \pi d\phi \wedge d^{m-1}x_0 - \frac{1}{2}\pi^2 d^m x \in \Omega^m(P_0) \quad , \\ \Omega_h^0 &= d\phi \wedge d\pi \wedge d^{m-1}x_0 + \pi d\pi \wedge d^m x. \end{aligned}$$

The spacetime symmetries are the Carroll transformations [26]

$$x'^0 = x^0 + b_k x^k \quad , \quad x'^i = x^i + \epsilon^i_j x^j \quad \Rightarrow \quad x'^\mu = C^\mu_\nu x^\nu \quad , \tag{29}$$

where $C^0_0 = 1, C^0_k = b_k, C^k_0 = 0$ and $(C^i_j) \in O(d)$. These transformations are represented infinitesimally by

$$C^\mu_\nu = \delta^\mu_\nu + \epsilon^\mu_\nu \quad , \tag{30}$$

so that now $b_k = \epsilon^0_k$ is infinitesimal. Up to linear terms in $\phi'(x') = \phi(x)$ and $\pi'(x') = \pi(x)$, the infinitesimal transformations of the fields are

$$\begin{aligned} \delta\phi(x) &\equiv \phi'(x) - \phi(x) = -\epsilon^\alpha_\beta x^\beta \frac{\partial\phi}{\partial x^\alpha} = -b_k x^k \frac{\partial\phi}{\partial x^0} - \epsilon^i_j x^j \frac{\partial\phi}{\partial x^i} \quad , \\ \delta\pi(x) &\equiv \pi'(x) - \pi(x) = -\epsilon^\alpha_\beta x^\beta \frac{\partial\pi}{\partial x^\alpha} = -b_k x^k \frac{\partial\pi}{\partial x^0} - \epsilon^i_j x^j \frac{\partial\pi}{\partial x^i}. \end{aligned}$$

Furthermore,

$$\phi'_\mu(x') = \frac{\partial x^\nu}{\partial x'^\mu} \phi_\nu(x) \iff \phi'_\mu(C \cdot x) = (C^{-1})^\nu_\mu \phi_\nu(x) \quad ,$$

where $C \equiv (C^\mu_\nu)$ and $C \cdot x \equiv C^\mu_\nu x^\nu$; hence,

$$\begin{aligned} \delta\phi_\mu(x) &\equiv \phi'_\mu(x) - \phi_\mu(x) = -\epsilon^v_\mu \phi_\nu - \epsilon^\alpha_\beta x^\beta \frac{\partial\phi_\mu}{\partial x^\alpha} \quad , \\ \delta\pi_\mu(x) &\equiv \pi'_\mu(x) - \pi_\mu = -\epsilon^v_\mu \pi_\nu - \epsilon^\alpha_\beta x^\beta \frac{\partial\pi_\mu}{\partial x^\alpha}. \end{aligned}$$

Component-wise,

$$\begin{aligned} \delta\phi_0(x) &= -\epsilon^\alpha_\beta x^\beta \frac{\partial\phi_0}{\partial x^\alpha} = -b_k x^k \frac{\partial\phi_0}{\partial x^0} - \epsilon^k_j x^j \frac{\partial\phi_0}{\partial x^k} \quad , \\ \delta\phi_i(x) &= -\epsilon^v_i \phi_\nu - \epsilon^\alpha_\beta x^\beta \frac{\partial\phi_i}{\partial x^\alpha} = -b_i \phi_0 - \epsilon^k_i \phi_k - b_k x^k \frac{\partial\phi_i}{\partial x^0} - \epsilon^k_j x^j \frac{\partial\phi_i}{\partial x^k} \quad , \end{aligned}$$

and similarly for the components of $\delta\pi_\mu(x)$. Under the transformations presented above, the Lagrangian transforms as $\delta\mathcal{L}(x) = -\partial_\alpha(\epsilon^\alpha_\beta x^\beta \mathcal{L}(x))$. These transformations are produced by the Lie derivatives of the local sections $\phi : M \rightarrow E$ and $j^1\phi : M \rightarrow J^1\pi$ by a

vector field $\zeta = -\zeta^\mu(x) \frac{\partial}{\partial x^\mu} \in \mathfrak{X}(M)$ which generates the Carroll transformations; where now $\zeta^\mu(x) = \epsilon^\mu_\nu x^\nu$ is given by (29) and (30). The vector field which generates the Carroll transformations on the configuration bundle E is

$$\zeta_E = -\epsilon^\mu_\nu x^\nu \frac{\partial}{\partial x^\mu} \in \mathfrak{X}(E),$$

while the canonical lift of ζ_E to $J^1\pi$ is given by

$$\begin{aligned} X &= -\epsilon^\mu_\nu x^\nu \frac{\partial}{\partial x^\mu} + \epsilon^\nu_\mu \phi_\nu \frac{\partial}{\partial \phi_\mu} + \epsilon^\nu_\mu \pi_\nu \frac{\partial}{\partial \pi_\mu} \\ &= -b_i x^i \frac{\partial}{\partial x^0} - \epsilon^i_j x^j \frac{\partial}{\partial x^i} + (b_i \phi_0 + \epsilon^j_i \phi_j) \frac{\partial}{\partial \phi_i} + (b_i \pi_0 + \epsilon^j_i \pi_j) \frac{\partial}{\partial \pi_i} \in \mathfrak{X}(J^1\pi). \end{aligned}$$

Evidently, X is tangent to the constraint submanifold \mathcal{S}_f since $L(X)(\pi - \phi_0) = 0$ and $L(X)\pi_0 = 0$. Finally, since $L(X)\mathcal{L} = 0 \Rightarrow L(X)\Theta_{\mathcal{L}} = 0$, the momentum map $J_{\mathcal{L}}(X)$ corresponding to the Carroll spacetime transformations in the Lagrangian setting is given by

$$\begin{aligned} J_{\mathcal{L}}(X) &= -i(X)\Theta_{\mathcal{L}} \\ &= -\epsilon^i_j x^j \pi d\phi \wedge d^{m-2}x_{0i} - \frac{1}{2}\pi^2 (b_i x^i d^{m-1}x_0 + \epsilon^i_j x^j d^{m-1}x_i) \in \Omega^{m-1}(J^1\pi). \end{aligned}$$

In the Hamiltonian setting, which takes place on $P_0 \subset J^1\pi^*$, the momentum map $J_h^0(Y) \in \Omega^{m-1}(P_0)$ is constructed using the vector field

$$Y = \mathcal{F}\mathcal{L}_{0*}X = -\epsilon^\mu_\nu x^\nu \frac{\partial}{\partial x^\mu} = -b_i x^i \frac{\partial}{\partial x^0} - \epsilon^i_j x^j \frac{\partial}{\partial x^i} \in \mathfrak{X}(P_0).$$

As in the Lagrangian setting, $L(Y)\Theta_h^0 = 0$ and $J_h^0(Y)$ is given by

$$J_h^0(Y) = -i(Y)\Theta_h^0 = -\epsilon^i_j x^j \pi d\phi \wedge d^{m-2}x_{0i} - \frac{1}{2}\pi^2 (b_i x^i d^{m-1}x_0 + \epsilon^i_j x^j d^{m-1}x_i).$$

5.5. Magnetic Carrollian Scalar Field Theory

The magnetic Carrollian contraction [26] of the canonical Klein–Gordon Lagrangian is performed by reinserting the factors of c (speed of light) into the Lagrangian (27) and taking the limit $c \rightarrow 0$. The Minkowski metric (28) becomes degenerate as in the electric Carrollian scalar field theory. The Lagrangian function $L \in C^\infty(J^1\pi)$ obtained from taking the magnetic Carrollian contraction of the canonical Klein–Gordon Lagrangian (27) is

$$L = \pi\phi_0 - \frac{1}{2}\phi_i\phi^i.$$

This Lagrangian is singular since the components of the Hessian matrix (with respect to the multivelocities) are

$$\frac{\partial^2 L}{\partial \phi_0 \partial \phi_0} = 0, \quad \frac{\partial^2 L}{\partial \phi_0 \partial \phi_i} = 0, \quad \frac{\partial^2 L}{\partial \phi_i \partial \phi_j} = -\delta^{ij}, \quad \frac{\partial^2 L}{\partial \pi_\mu \partial \pi_\nu} = 0, \quad \frac{\partial^2 L}{\partial \pi_\mu \partial \phi_\nu} = 0.$$

The Lagrangian energy function and the Poincaré–Cartan forms are now

$$\begin{aligned} E_{\mathcal{L}} &= -\frac{1}{2}\phi_i\phi^i \in C^\infty(J^1\pi), \\ \Theta_{\mathcal{L}} &= \pi d\phi \wedge d^{m-1}x_0 - \phi^i d\phi \wedge d^{m-1}x_i + \frac{1}{2}\phi_i\phi^i d^m x \in \Omega^m(J^1\pi), \\ \Omega_{\mathcal{L}} &= d\phi \wedge d\pi \wedge d^{m-1}x_0 + d\phi^i \wedge d\phi \wedge d^{m-1}x_i + \phi^i d\phi_i \wedge d^m x \in \Omega^{m+1}(J^1\pi), \end{aligned}$$

and $\Omega_{\mathcal{L}}$ is a premultisymplectic form. Imposing the SOPDE condition,

$$C_i^i - \pi_0 = 0 \quad , \quad \phi_0 = 0.$$

The first equation gives a relation for the coefficients C_i^i , and the second one is a SOPDE-constraint which defines the constraint submanifold $S_1 \hookrightarrow J^1\pi$, and no more constraints arise from the constraint algorithm.

The Legendre map $\mathcal{FL} : J^1\pi \rightarrow J^1\pi^*$ gives the multimomenta

$$\mathcal{FL}^* p_\phi^0 = \frac{\partial \mathcal{L}}{\partial \phi_0} = \pi, \quad \mathcal{FL}^* p_\phi^i = \frac{\partial \mathcal{L}}{\partial \phi_i} = -\phi^i, \quad \mathcal{FL}^* p_\pi^\mu = \frac{\partial \mathcal{L}}{\partial \pi_\mu} = 0.$$

It follows that $p_\phi^0 - \pi = 0$ and $p_\pi^\mu = 0$ are primary constraints which define the primary constraint submanifold $P_0 \hookrightarrow J^1\pi^*$ with local coordinates $(x^\mu, \phi, \pi, p_\phi^i)$, and the Lagrangian is again almost regular. The De Donder–Weyl Hamiltonian $H_0 \in C^\infty(P_0)$ is given by

$$H_0 = -\frac{1}{2} p_{\phi i} p_\phi^i,$$

and the Hamilton–Cartan forms on P_0 are

$$\begin{aligned} \Theta_h^0 &= \pi d\phi \wedge d^{m-1}x_0 + p_\phi^i d\phi \wedge d^{m-1}x_i + \frac{1}{2} p_{\phi i} p_\phi^i d^m x \in \Omega^m(P_0), \\ \Omega_h^0 &= d\phi \wedge d\pi \wedge d^{m-1}x_0 + d\phi \wedge dp_\phi^i \wedge d^{m-1}x_i - p_{\phi i} dp_\phi^i \wedge d^m x \in \Omega^{m+1}(P_0). \end{aligned}$$

The Carroll transformations which are symmetries for the magnetic scalar field theory are written slightly differently than for the electric scalar field and are

$$\begin{aligned} \delta\phi(x) &= -\epsilon^\alpha_\beta x^\beta \frac{\partial\phi}{\partial x^\alpha} = -b_k x^k \frac{\partial\phi}{\partial x^0} - \epsilon^i_j x^j \frac{\partial\phi}{\partial x^i}, \\ \delta\pi(x) &= -\epsilon^\alpha_\beta x^\beta \frac{\partial\pi}{\partial x^\alpha} - b_i \phi^i = -b_k x^k \frac{\partial\pi}{\partial x^0} - \epsilon^i_j x^j \frac{\partial\pi}{\partial x^i} - b_i \phi^i, \\ \delta\phi_0(x) &= -\epsilon^\alpha_\beta x^\beta \frac{\partial\phi_0}{\partial x^\alpha} = -b_k x^k \frac{\partial\phi_0}{\partial x^0} - \epsilon^k_j x^j \frac{\partial\phi_0}{\partial x^k}, \\ \delta\phi_i(x) &= -\epsilon^\nu_i \phi_\nu - \epsilon^\alpha_\beta x^\beta \frac{\partial\phi_i}{\partial x^\alpha} = -b_i \phi_0 - \epsilon^k_i \phi_k - b_k x^k \frac{\partial\phi_i}{\partial x^0} - \epsilon^k_j x^j \frac{\partial\phi_i}{\partial x^k}. \end{aligned}$$

Under these transformations, the Lagrangian transforms as $\delta L(x) = -\frac{\partial}{\partial x^\alpha} \left(\epsilon^\alpha_\beta x^\beta L(x) \right)$. Notice that the field $\pi(x)$ no longer transforms as a Carrollian scalar; instead, $\pi(x)$ transforms under a Carroll spacetime transformation $-\epsilon^\alpha_\beta x^\beta \frac{\partial\pi}{\partial x^\alpha}$ plus a term $-b_i \phi^i(x)$ which cannot be represented on E . The field transformations behave geometrically as the Lie derivatives of the local sections $j^1\phi : M \rightarrow J^1\pi$. The vector field generating the Carroll transformations on $J^1\pi$ is

$$\begin{aligned} X &= -\epsilon^\mu_\nu x^\nu \frac{\partial}{\partial x^\mu} + b_i \phi^i \frac{\partial}{\partial \pi} + \epsilon^\nu_\mu \phi_\nu \frac{\partial}{\partial \phi_\mu} + \epsilon^\nu_\mu \pi_\nu \frac{\partial}{\partial \pi_\mu} \\ &= -b_i x^i \frac{\partial}{\partial x^0} - \epsilon^i_j x^j \frac{\partial}{\partial x^i} + b_i \phi^i \frac{\partial}{\partial \pi} + \left(b_i \phi_0 + \epsilon^j_i \phi_j \right) \frac{\partial}{\partial \phi_i} + \epsilon^\nu_\mu \pi_\nu \frac{\partial}{\partial \pi_\mu} \in \mathfrak{X}(J^1\pi), \end{aligned}$$

which is tangent to the constraint submanifold $S_1 \subset J^1\pi$, given by the constraint $\phi_0 = 0$, since $L(X)\phi_0 = 0$. Furthermore, although the vector field X leaves the Lagrangian density invariant ($L(X)\mathcal{L} = 0$), it does not produce an exact Noether symmetry as

$$L(X)\Theta_{\mathcal{L}} = b^i \phi_0 (\phi_i d^m x - d\phi \wedge d^{m-1}x_i).$$

Moreover, the vector field X as written above on all of $J^1\pi$ is not projectable onto $P_0 \subset J^1\pi^*$. Instead, X on S_1 given by

$$X|_{S_1} = -b_i x^i \frac{\partial}{\partial x^0} - \epsilon^i_j x^j \frac{\partial}{\partial x^i} + b_i \phi^i \frac{\partial}{\partial \pi} + \epsilon^i_j \phi_j \frac{\partial}{\partial \phi_i} + \epsilon^\nu_\mu \pi_\nu \frac{\partial}{\partial \pi_\mu}, \quad (31)$$

is \mathcal{FL}_0 -projectable onto P_0 , giving

$$Y = \mathcal{FL}_{0*} X|_{S_1} = -b_i x^i \frac{\partial}{\partial x^0} - \epsilon^i_j x^j \frac{\partial}{\partial x^i} - b_i p_\phi^i \frac{\partial}{\partial \pi} + \epsilon^i_j p_\phi^j \frac{\partial}{\partial p_\phi^i} \in \mathfrak{X}(P_0).$$

Now, letting \tilde{X} be the local extension of $X|_{S_1}$ to $J^1\pi$ whose coordinate expression given by (31), it follows that $L(\tilde{X})\Theta_{\mathcal{L}} = 0$ on $J^1\pi$. It is thereby possible to define the momentum maps, giving

$$\begin{aligned} J_{\mathcal{L}}(\tilde{X}) = -i(\tilde{X})\Theta_{\mathcal{L}} &= -\left(\epsilon^i_j x^j \pi + b_j x^j \phi^i\right) d\phi \wedge d^{m-2}x_{0i} + \phi^i \epsilon^j_k x^k d\phi \wedge d^{m-2}x_{ij} \\ &\quad + \frac{1}{2} \phi_i \phi^i \left(b_k x^k d^{m-1}x_0 + \epsilon^k_j x^j d^{m-1}x_k\right) \in \Omega^{m-1}(J^1\pi), \\ J_h^0(Y) = -i(Y)\Theta_h^0 &= -\left(\epsilon^i_j x^j \pi - b_j x^j p_\phi^i\right) d\phi \wedge d^{m-2}x_{0i} - p_\phi^i \epsilon^j_k x^k d\phi \wedge d^{m-2}x_{ij} \\ &\quad + \frac{1}{2} p_{\phi i} p_\phi^i \left(b_k x^k d^{m-1}x_0 + \epsilon^k_j x^j d^{m-1}x_k\right) \in \Omega^{m-1}(P_0). \end{aligned}$$

6. Conclusions

One of the primary objectives achieved in this work is the precise connection of how classical field theories are studied in the standard physics literature and how they are treated in the more modern works of (pre)multiplectic geometry. This work provides a general overview of Cartan (Noether) symmetries present in (pre)multiplectic geometry and how they are used to describe Noether symmetries in classical field theories both in the Lagrangian and DeDonder–Weyl Hamiltonian formalisms. In the literature of differential geometry, the treatment of Cartan symmetries is well-known [9,14,16,17]. However, subtleties arise in the treatment of Noether symmetries of field theories whose Lagrangians are singular; the constraints of such singular field theories are obtained from the premultiplectic forms of the corresponding multiphase spaces [38]. In particular, the analysis of Noether symmetries in the presence of the constraints is described in this paper.

More specifically, the Noether symmetries of classical field theories developed on the relevant (pre)multiplectic multi-phase spaces are encoded by their conserved quantities, called multimomentum maps (as described by Noether's theorem). The construction of the relevant multimomentum maps is carried out explicitly for various singular field theories that are important in theoretical physics. These multimomentum maps give the standard Noether currents of classical field theories when pulled back by local sections which map from the base space of the bundle structure to the multi-phase spaces. Furthermore, the multimomentum maps associated with gauge symmetries are essential for carrying out a desired symmetry reduction procedure (or gauge fixing), e.g., [19]; the formal development of such symmetry reduction procedures in premultiplectic field theories continues to be an open area of research (see, for instance, [18,45,46]).

Author Contributions: Investigation, A.G.IV and N.R.-R. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by *Ministerio de Ciencia, Innovación y Universidades* (Spain), project PID2021-125515NB-C21, and the financial support for research groups AGRUPS-2022 of the *Universitat Politècnica de Catalunya* (UPC).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data sharing not applicable.

Acknowledgments: We want to thank Joaquim Gomis and Jordi Gaset for helpful and clarifying discussions. We are indebted to Victor Tapia for having drawn our attention to an error in the first calculation of Equation (21). We also thank the referees for their constructive comments.

Conflicts of Interest: The authors declare no conflict of interest.

Appendix A. Multivector Fields on Manifolds and Fiber Bundles

(See Refs. [37,47,48]). Let \mathcal{M} be an N -dimensional differentiable manifold. The m -**multivector fields** in \mathcal{M} ($m \leq N$) are the sections of the m -multitangent bundle $\bigwedge^m T\mathcal{M} := \overbrace{T\mathcal{M} \wedge \dots \wedge T\mathcal{M}}^m$; that is, the skew-symmetric contravariant tensor fields of order m in \mathcal{M} ; the set of which is denoted $\mathfrak{X}^m(\mathcal{M})$. Then, if $\mathbf{X} \in \mathfrak{X}^m(\mathcal{M})$, for every point $\bar{y} \in \mathcal{M}$, there is an open neighbourhood $U \subset \mathcal{M}$ and $X_1, \dots, X_r \in \mathfrak{X}(U)$ such that, for $m \leq r \leq \dim \mathcal{M}$,

$$\mathbf{X}|_U = \sum_{1 \leq i_1 < \dots < i_m \leq r} f^{i_1 \dots i_m} X_{i_1} \wedge \dots \wedge X_{i_m},$$

with $f^{i_1 \dots i_m} \in C^\infty(U)$. In particular, $\mathbf{X} \in \mathfrak{X}^m(\mathcal{M})$ is said to be a **locally decomposable multivector field** if there exist $X_1, \dots, X_m \in \mathfrak{X}(U)$ such that $\mathbf{X}|_U = X_1 \wedge \dots \wedge X_m$. The locally decomposable m -multivector fields are locally associated with m -dimensional distributions $D \subset T\mathcal{M}$, and this splits $\mathfrak{X}^m(\mathcal{M})$ into *equivalence classes* $\{\mathbf{X}\} \subset \mathfrak{X}^m(\mathcal{M})$ which are made of the locally decomposable multivector fields associated with the same distribution. If $\mathbf{X}, \mathbf{X}' \in \{\mathbf{X}\}$ then, for every $U \subset \mathcal{M}$, there exists a non-vanishing function $f \in C^\infty(U)$ such that $\mathbf{X}' = f\mathbf{X}$ on U .

If $\Omega \in \Omega^p(\mathcal{M})$ and $\mathbf{X} \in \mathfrak{X}^m(\mathcal{M})$, the **contraction** between \mathbf{X} and Ω is the natural contraction between tensor fields; in particular, it gives zero when $p < m$ and, if $p \geq m$,

$$i(\mathbf{X})\Omega|_U := \sum_{1 \leq i_1 < \dots < i_m \leq r} f^{i_1 \dots i_m} i(X_{i_1} \wedge \dots \wedge X_{i_m})\Omega = \sum_{1 \leq i_1 < \dots < i_m \leq r} f^{i_1 \dots i_m} i(X_{i_m}) \dots i(X_{i_1})\Omega.$$

The **Lie derivative** of Ω with respect to \mathbf{X} is the graded bracket (of degree $m - 1$)

$$L(\mathbf{X})\Omega := [d, i(\mathbf{X})]\Omega = (d i(\mathbf{X}) - (-1)^m i(\mathbf{X})d)\Omega.$$

If $\mathbf{X} \in \mathfrak{X}^i(\mathcal{M})$ and $\mathbf{Y} \in \mathfrak{X}^j(\mathcal{M})$, the **Schouten-Nijenhuis bracket** of \mathbf{X}, \mathbf{Y} is the bilinear map $\mathbf{X}, \mathbf{Y} \mapsto [\mathbf{X}, \mathbf{Y}]$, where $[\mathbf{X}, \mathbf{Y}]$ is a $(i + j - 1)$ -multivector field obtained as the graded commutator of $L(\mathbf{X})$ and $L(\mathbf{Y})$; that is,

$$L([\mathbf{X}, \mathbf{Y}])\Omega := [L(\mathbf{X}), L(\mathbf{Y})]\Omega.$$

This is an operation of degree $i + j - 2$ which is also called the **Lie derivative** of \mathbf{Y} with respect to \mathbf{X} , and is denoted $L(\mathbf{X})\mathbf{Y} := [\mathbf{X}, \mathbf{Y}]$. Furthermore, if $Y \in \mathfrak{X}(\mathcal{M})$ and $\mathbf{X} \in \mathfrak{X}^m(\mathcal{M})$, then

$$i([Y, \mathbf{X}])\Omega = L(Y) i(\mathbf{X})\Omega - i(\mathbf{X}) L(Y)\Omega. \tag{A1}$$

Now, let $q: \mathcal{M} \rightarrow M$ a fiber bundle. A multivector field $\mathbf{X} \in \mathfrak{X}^m(\mathcal{M})$ is q -*transverse* if, for every $\beta \in \Omega^m(M)$ such that $\beta_{\rho(p)} \neq 0$, at every point $p \in \mathcal{M}$, we have that $(i(\mathbf{X})(\bar{\pi}^{k*}\beta))_p \neq 0$. We are interested in the **integrable multivector fields**, which are those locally decomposable multivector fields whose associated distribution is integrable. Then, if $\mathbf{X} \in \mathfrak{X}^m(\mathcal{M})$ is integrable and q -transverse, its integral manifolds are local sections of the projection $q: \mathcal{M} \rightarrow M$.

In the particular case where $\mathcal{M} = J^1\pi$ and we have the jet bundle $\bar{\pi}^1: J^1\pi \rightarrow M$; an integrable multivector field $\mathbf{X} \in \mathfrak{X}^m(\mathcal{M})$ is said to be **holonomic** if its integral sections are holonomic sections of $\bar{\pi}^1$.

References

1. Kosmann-Schwarzbach, Y. *The Noether Theorems. Invariance and Conservation Laws in the Twentieth Century*; Springer: New York, NY, USA, 2011. [CrossRef]
2. Aldaya, V.; de Azcarraga, J.A. Variational Principles on r th order jets of fibre bundles in Field Theory. *J. Math. Phys.* **1978**, *19*, 1869–1875. [CrossRef]
3. Echeverría-Enríquez, A.; Muñoz-Lecanda, M.C.; Román-Roy, N. Geometry of Lagrangian first-order classical field theories. *Forts Phys.* **1996**, *44*, 235–280. [CrossRef]
4. García, P.L. The Poincaré–Cartan invariant in the calculus of variations. *Symp. Math.* **1973**, *14*, 219–246.
5. Goldschmidt, H.; Sternberg, S. The Hamilton–Cartan formalism in the calculus of variations. *Ann. Inst. Fourier Grenoble* **1973**, *23*, 203–267. Available online: <http://eudml.org/doc/74112> (accessed on 27 January 2023). [CrossRef]
6. Hélein, F.; Kouneither, J. Covariant Hamiltonian formalism for the calculus of variations with several variables: Lepage–Dedecker versus De Donder–Weyl. *Adv. Theor. Math. Phys.* **2004**, *8*, 565–601. [CrossRef]
7. Saunders, D.J. *The Geometry of Jet Bundles*; London Mathematical Society, Lect. Notes Ser. 142; Cambridge University Press: Cambridge, MA, USA; New York, NY, USA, 1989. [CrossRef]
8. Dedecker, P. On the Generalization of Symplectic Geometry to Multiple Integrals in the Calculus of Variations. In *Differential Geometrical Methods in Mathematical Physics (Proc. Sympos., Univ. Bonn, Bonn, 1975)*; Lecture Notes in Math.; Springer: Berlin, Germany, 1977; Volume 570, pp. 395–456. [CrossRef]
9. de León, M.; Martín de Diego, D.; Santamaría-Merino, A. Symmetries in classical field theory. *Int. J. Geom. Meths. Mod. Phys.* **2004**, *1*, 651–710. [CrossRef]
10. Forger, M.; Soares, B.L. Local symmetries in gauge theories in a finite-dimensional setting. *J. Geom. Phys.* **2012**, *62*, 1925–1938. [CrossRef]
11. Giachetta, G.; Mangiarotti, L.; Sardanashvily, G. *New Lagrangian and Hamiltonian Methods in Field Theory*; World Scientific Publishing Co., Inc.: River Edge, NJ, USA, 1997. [CrossRef]
12. Krupka, D. *Introduction to Global Variational Geometry*; Atlantis Studies in Variational Geometry; Atlantis Press: Paris, France, 2015. Available online: <https://link.springer.com/book/10.2991/978-94-6239-073-7> (accessed on 27 January 2023).
13. Echeverría-Enríquez, A.; Muñoz-Lecanda, M.C.; Román-Roy, N. Multivector field formulation of Hamiltonian field theories: Equations and symmetries. *J. Phys. A Math. Gen.* **1999**, *32*, 8461–8484. [CrossRef]
14. Gaset, J.; Prieto-Martínez, P.D.; Román-Roy, N. Variational principles and symmetries on fibered multisymplectic manifolds. *Comm. Math.* **2016**, *24*, 137–152. [CrossRef]
15. Gaset, J.; Román-Roy, N. Symmetries and gauge symmetries in multisymplectic first and second-order Lagrangian field theories: Electromagnetic and gravitational fields. *Rev. Acad. Cienc. Canar. Math. arXiv* **2021**, arXiv:2107.08846.
16. Gotay, M.J.; Isenberg, J.; Marsden, J.E.; Montgomery, R. Momentum maps and classical relativistic fields. I. Covariant theory. *arXiv* **2004**, arXiv:physics/9801019.
17. Ryvkin, L.; Wurzbacher, T.; Zambon, M. Conserved quantities on multisymplectic manifolds. *J. Austral Math. Soc.* **2020**, *108*, 120–144. [CrossRef]
18. Blacker, C. Reduction of multisymplectic manifolds. *Lett. Math. Phys.* **2021**, *111*, 64. [CrossRef]
19. Echeverría-Enríquez, A.; Muñoz-Lecanda, M.C.; Román-Roy, N. Remarks on multisymplectic reduction. *Rep. Math. Phys.* **2018**, *81*, 415–424. [CrossRef]
20. Madsen, T.B.; Swann, A. Multimoment maps. *Adv. Math.* **2012**, *229*, 2287–2309. [CrossRef]
21. Sniatycki, J. Multisymplectic reduction for proper actions. *Can. J. Math.* **2004**, *56*, 638–654. [CrossRef]
22. Arnold, V.I. *Mathematical Methods of Classical Mechanics*; Graduate Texts in Mathematics 60; Springer: New York, NY, USA, 1989. [CrossRef]
23. Mangiarotti, L.; Sardanashvily, G. *Gauge Mechanics*; World Scientific: Singapore, 1998. [CrossRef]
24. Sarlet, W.; Cantrijn, F. Higher-order Noether symmetries and constants of the motion. *J. Phys. A Math. Gen.* **1981**, *14*, 479–492. [CrossRef]
25. Bleecker, D. *Gauge Theory and Variational Principles*; Dover Publications: New York, NY, USA, 1981; ISBN: 0486445461.
26. Henneaux, M.; Salgado-Rebolledo, P. Carroll contractions of Lorentz-invariant theories. *J. High Energy Phys.* **2021**, *2021*, 180. [CrossRef]
27. de León, M.; Marín-Solano, J.M.; Marrero, J.C.; Muñoz-Lecanda, M.C.; Román-Roy, N. Premultisymplectic constraint algorithm for field theories. *Int. J. Geom. Meth. Mod. Phys.* **2005**, *2*, 839–871. [CrossRef]
28. Román-Roy, N. A summary on symmetries and conserved quantities of autonomous Hamiltonian systems. *J. Geom. Mech.* **2020**, *12*, 541–551. [CrossRef]
29. Gaset, J. Geometric Gauge Freedom in Multisymplectic Field Theories. *arXiv* **2022**, arXiv:2209.11212.
30. Bergvelt, M.J.; de Kerf, E.A. The Hamiltonian structure of Yang–Mills theories and instantons (Part I). *Physica* **1986**, *139A*, 101–124. [CrossRef]
31. Gotay, M.J.; Nester, J.M. Presymplectic Hamilton and Lagrange systems, gauge transformations and the Dirac theory of constraints. In *Group Theoretical Methods in Physics*; Beigelbock, W., Böhm, A., Takasugi, E., Eds.; Lect. Notes in Phys.; Springer: Berlin, Germany, 1979; Volume 94, pp. 272–279. [CrossRef]

32. Batlle, C.; Gomis, J.; París, J.; Roca, J. Field-antifield formalism and Hamiltonian BRST approach. *Nucl. Phys. B* **1990**, *329*, 139–154. [[CrossRef](#)]
33. Gomis, J.; París, J.; Samuel, S. Antibracket, antifields and gauge-theory quantization. *Phys. Rep.* **1995**, *259*, 1–145. [[CrossRef](#)]
34. Gràcia, X.; Pons, J.M. Symmetries and infinitesimal symmetries of singular differential equations. *J. Phys. A Math. Gen.* **2002**, *35*, 5059. [[CrossRef](#)]
35. Kolár, I.; Slovák, J.; Michor, P.W. *Natural Operations in Differential Geometry*; Springer: Berlin/Heidelberg, Germany, 1993. [[CrossRef](#)]
36. Michor, P.W. *Topics in Differential Geometry*; Grad. Stud. in Math. 93; American Mathematical Soc.: Wien, Austria, 2008; ISBN: 0821820036.
37. Echeverría-Enríquez, A.; Muñoz-Lecanda, M.C.; Román-Roy, N. Multivector fields and connections: Setting Lagrangian equations in field theories. *J. Math. Phys.* **1998**, *39*, 4578–4603. [[CrossRef](#)]
38. Gomis, J.; Guerra, A., IV; Román-Roy, N. Multisymplectic Constraint Analysis of Scalar Field Theories, Chern-Simons Gravity, and Bosonic String Theory. *Nucl. Phys. B* **2023**, *987*, 116069. [[CrossRef](#)]
39. Cariñena, J.F.; Crampin, M.; Ibort, L.A. On the multisymplectic formalism for first order field theories. *Diff. Geom. Appl.* **1991**, *1*, 345–374. [[CrossRef](#)]
40. de León, M.; Marín-Solano, J.M.; Marrero, J.C. A Geometrical approach to Classical Field Theories: A constraint algorithm for singular theories. In *New Developments in Differential Geometry*; Tamassi, L., Szenthe, J., Eds.; Kluwer Acad. Press: Amsterdam, The Netherlands, 1996; pp. 291–312. [[CrossRef](#)]
41. Román-Roy, N. Multisymplectic Lagrangian and Hamiltonian formalisms of classical field theories. *Symm. Integ. Geom. Meth. Appl. (SIGMA)* **2009**, *5*, 100. [[CrossRef](#)]
42. Castrillón, M.; Muñoz-Masqué, J.; Rosado, E. First-order equivalent to Einstein–Hilbert Lagrangian. *J. Math. Phys.* **2014**, *55*, 082501. [[CrossRef](#)]
43. Adame-Carrillo, D.; Gaset, J.; Román-Roy, N. The second-order problem for k-presymplectic Lagrangian field theories: Application to the Einstein–Palatini model. *RACSAM* **2022**, *116*, 20. [[CrossRef](#)]
44. Becker, K.; Becker, M.; Schwarz, J.H. *String Theory and M-Theory. A Modern Introduction*; Cambridge University Press: Cambridge, MA, USA, 2006. [[CrossRef](#)]
45. Blacker, C. Reduction of L_∞ -algebras of observables on multisymplectic manifolds. *arXiv* **2022**, arXiv:92206.03137.
46. Ciaglia, F.M.; Cosmo, F.D.; Ibort, A.; Marmo, G.; Schiavone, L. Covariant reduction of classical Hamiltonian Field Theories: From D’Alembert to Klein–Gordon and Schrödinger. *Mod. Phys. Lett. A* **2020**, *35*, 2050214. [[CrossRef](#)]
47. Cantrijn, F.; Ibort, L.A.; de León, M. Hamiltonian structures on multisymplectic manifolds. *Rend. Sem. Mat. Univ. Pol. Torino* **1996**, *54*, 225–236. Available online: <https://www.researchgate.net/publication/233917958> (accessed on 27 January 2023).
48. Cantrijn, F.; Ibort, L.A.; de León, M. On the geometry of multisymplectic manifolds. *J. Aust. Math. Soc.* **1999**, *66*, 303–330. Available online: <https://www.researchgate.net/publication/231926129> (accessed on 27 January 2023). [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.