

Proceeding Paper

# The Four Principal Megabiases in the Known Fossil Record: Taphonomy, Rock Preservation, Fossil Discovery and Fossil Study <sup>†</sup>

Adrian P. Hunt <sup>1,\*</sup> and Spencer G. Lucas <sup>2</sup> 

<sup>1</sup> Flying Heritage and Combat Armor Museum, 3407 109th St. SW, Everett, WA 98204, USA

<sup>2</sup> New Mexico Museum of Natural History, 1801 Mountain Road N. W., Albuquerque, NM 87104, USA; spencer.lucas@dca.nm.gov

\* Correspondence: ahunt@flyingheritage.org

† Presented at the 4th International Electronic Conference on Geosciences, 1–15 December 2022; Available online: <https://sciforum.net/event/IECG2022>.

**Abstract:** The Known Fossil Record represents museum collections and the published literature, and it is subject to multiple large-scale megabiases grouped into four major categories: (1) taphonomy; (2) rock preservation; (3) fossil discovery; and (4) fossil study. Taphonomic megabiases are largescale patterns in the quality of the fossil record that affect paleobiologic analysis at provincial to global levels and at timescales usually exceeding ten million years. Taphonomic megabiases are intrinsic (form and behavior) and extrinsic (biotic and abiotic controls on preservation). Other megabiases are the preservation and exposure of rock strata, kyreonomy (discovery) and concipionomy (study). Kyreonomy megabiases include location of fossil sites, mineral evaluation, mineral extraction and colonialism. Concipionomy megabiases include the Taxophile Effect, language and development and distribution of technology.

**Keywords:** fossil record; megabiases; taphonomy; kyreonomy; concipionomy



**Citation:** Hunt, A.P.; Lucas, S.G. The Four Principal Megabiases in the Known Fossil Record: Taphonomy, Rock Preservation, Fossil Discovery and Fossil Study. *Proceedings* **2023**, *87*, 13. <https://doi.org/10.3390/IECG2022-13956>

Academic Editor: Angelos G. Maravelis

Published: 9 January 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The Known Fossil Record represents museum collections and the published literature [1], and it is subject to multiple large-scale biases. The purpose of this paper is to demonstrate that megabiases in the fossil record can be grouped into four major categories: (1) taphonomic megabiases; (2) rock preservation; (3) fossil discovery; and (4) fossil study.

## 2. Taphonomic Megabiases

Behrensmeyer et al. [2] introduced the term megabiases into taphonomy for largescale patterns in the quality of the fossil record that affect paleobiologic analysis at provincial to global levels and at timescales usually exceeding ten million years. Taphonomic megabiases are intrinsic (form and behavior) and extrinsic (biotic and abiotic controls on preservation). Examples of intrinsic megabiases in the vertebrate-fossil record include body size (larger organisms are better preserved), robusticity of skeleton (e.g., fewer bird and pterosaur fossils), presence of armor (dense osteoderms preserve well such as in the nonmarine Late Triassic) and behavior (e.g., semiaquatic or terrestrial).

One significant extrinsic megabiases involves the development of vascular plants and related land surface evolution. Schumm [3,4] first speculated that plant evolution caused changes in fluvial style, and Cotter [5] documented the relationship in the Paleozoic. Essentially, increasing plant cover in the Paleozoic led to increased stabilization of channels and floodplains [6–8]. This had profound effects on the taphonomy of plants, trace fossils and body fossils [7,9–11]. The later evolution of land plants, notably grasses, presumably had additional taphonomic impacts. Thus, Hunt et al. (Ref. [11] predicted four distinct

temporal phases of vertebrate track preservation: (1) Devonian—few tracks, because terrestrial tetrapods are rare, and lack of plant ground cover resulted in frequent reworking of terrestrial surfaces; (2) Carboniferous-Triassic—many tracks because terrestrial tetrapods are common, and increased ground cover reduced the reworking of terrestrial surfaces; (3) Jurassic-Cretaceous—tracks will be numerous and preserved in more diverse sedimentary environments because terrestrial animals are very large, even though ground cover is increased; (4) Cenozoic—increased ground cover, especially after the diversification of grasses, resulted in less unvegetated areas where tracks can be preserved (with a few notable exceptions such as lacustrine margins).

Another extrinsic example is the evolution of dentition. Pre-mammalian vertebrates generally lack the dental morphology for fine occlusion. Thus, for example, dentalites are rarer on dinosaur bones than on Cenozoic mammal bones because non-avian theropods lacked the dentition or jaw mechanics to manipulate and modify bones in a similar manner [9,10,12,13]. Fiorillo [13] validated this hypothesis by demonstrating that dinosaur faunas exhibited 4% or less of bones with dentalites, whereas in the mammal faunas he studied, the percentages varied from 13.1 to 37.5% (however, see [14] for a notable exception).

Other examples of extrinsic megabiases include digestive evolution (e.g., preservational effects of GI tract acidity [9,10,15] and Lagerstätten and Megalagerstätten (e.g., Upper Cretaceous of Western Interior of North America [16])).

### 3. Rock Preservation

Raup [17] persuasively argued that aspects of the rock record have resulted in systematic biases in the fossil record, notably exposed rock area, available rock volume and intensity of subsequent metamorphism and erosion. Subsequent workers have discussed aspects of this topic with regard to diversity through time (e.g., [18–22]). Sheehan [19] recognized Paleontologic Interest Units as a measure of the effort devoted to acquiring knowledge concerning fossils and concluded that eight times as many paleontologists (per million years) work on Cenozoic fossils as on Cambrian fossils, reflecting the relative exposure of these ages of rocks [19].

### 4. Fossil Discovery

The history of the discovery of fossils has been heavily influenced by the prospecting for, and extraction of, mineral resources. For example, the major difference between Moscovian and older/younger Carboniferous tetrapod records has its primary basis in coal mining [22,23]. Thus, the larger Middle Pennsylvanian tetrapod record is biased because almost all of the Moscovian tetrapod assemblages are associated with coal beds [23–26]. There is an abrupt decrease in mineable coals across the Middle–Late Pennsylvanian boundary, due to climate change driven by sea-level drop, the drifting northward of Euramerica and changing topography and drainage patterns due to Variscan tectonism (e.g., [27–29]). The tetrapod fossil record diminishes with these changes because of the megabiases associated with the coal interval.

We apply the term *kyreonomy* (from the Greek *kyreo* to find) to address biases caused by discovery. Other *kyreonomic* megabiases include location of fossil sites (related to human geography, climate and geological context), mineral evaluation (e.g., exploration of Cretaceous coalfields of western USA), mineral extraction (e.g., Late Jurassic–Early Cretaceous lithographic limestones) and colonialism (e.g., Tanzanian dinosaurs, Karoo tetrapods, North African dinosaurs).

### 5. Fossil Study

There are markedly distinct levels of interest, and hence study, of different fossil groups, which Hunt et al. [30] termed the Taxophile Effect (e.g., [17]). This is clearly evident, for example, among dentalites where a single tooth mark on a dinosaur bone warrants a published paper, whereas the numerous occurrences on Cenozoic bones are barely noted

until the Quaternary [15]. There is consistent elevated interest in certain groups evident from popular culture to rock shop sales to the scientific literature. Dinosaurs and ammonites are the clear winners among vertebrates and invertebrates, respectively. Intrinsic interest is not the only driver to increased study. Extrinsic factors include the relative abundance of exposure of strata in that “geologic systems with more rock contain more species and this leads to more species being described” ([31], p. 328) and [19]. Other extrinsic factors include employment possibilities such as the decrease in the number of petroleum company micropaleontologists and the decline in traditional systematists among academic faculty. The burgeoning of paleobiology has benefitted the understanding patterns of the fossil record but has had a detrimental effect on the prevalence of study of the building blocks of the record. Other biases include language (non-English literature is under cited) and the development and availability of technology (e.g., SEM, CT scanning). We propose the term concipionomy (from the Latin *concupio* to comprehend) for the biases introduced by study of the fossil record.

## 6. Conclusions

The Known Fossil Record is subject to multiple large-scale biases. These megabiases can be grouped into four major categories: (1) taphonomy; (2) rock preservation; (3) kyreonomy; and (4) concipionomy.

**Author Contributions:** Conceptualization, A.P.H. and S.G.L.; investigation, A.P.H. and S.G.L.; writing—original draft preparation, A.P.H.; writing—review and editing, A.P.H. and S.G.L. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Benton, M.J.; Dunhill, A.M.; Lloyd, G.T.; Marx, F.G. Assessing the quality of the fossil record: Insights from vertebrates. *Geol. Soc. Lond. Spec. Publ.* **2011**, *358*, 63–94. [[CrossRef](#)]
2. Behrensmeyer, A.K.; Kidwell, S.M.; Gastaldo, R.A. Taphonomy and paleobiology. *Paleobiology* **2000**, *26*, 103–147. [[CrossRef](#)]
3. Schumm, S.A. Paleohydrology: Applications of modern hydrologic data to problems of the ancient past. In Proceedings of the International Hydrology Symposium, Fort Collins, CO, USA, 6–7 September 1967; Volume 1, pp. 185–193.
4. Schumm, S.A. Speculations concerning paleohydrologic controls of terrestrial sedimentation. *Geol. Soc. Am. Bull.* **1968**, *79*, 1573–1588. [[CrossRef](#)]
5. Cotter, E. The evolution of fluvial style, with special reference to the central Appalachian Paleozoic. *Can. Soc. Pet. Geol. Mem.* **1978**, *5*, 361–383.
6. Davies, N.S.; Gibling, M.R. Cambrian to Devonian evolution of alluvial systems: The sedimentological impact of the earliest land plants. *Earth Sci. Rev.* **2010**, *98*, 171–200. [[CrossRef](#)]
7. Davies, N.S.; Gibling, M.R. The sedimentary record of Carboniferous rivers: Continuing influence of land plant evolution on alluvial processes and Palaeozoic ecosystems. *Earth Sci. Rev.* **2013**, *120*, 40–79. [[CrossRef](#)]
8. Gibling, M.R.; Davies, N.S. Palaeozoic landscapes shaped by plant evolution. *Natl. Geogr.* **2012**, *5*, 99–105. [[CrossRef](#)]
9. Hunt, A.P. Fluvial vertebrate taphonomy: Historical perspectives. *New Mex. J. Sci.* **1984**, *24*, 26–27.
10. Hunt, A.P. Phanerozoic trends in nonmarine taphonomy: Implications for Mesozoic vertebrate taphonomy and paleoecology. *Geol. Soc. Am. Abstr. Programs* **1987**, *19*, 171.
11. Hunt, A.P.; Santucci, V.L.; Lucas, S.G. *Vertebrate Trace Fossils from Arizona with Special Reference to Tracks Preserved in National Park Service Units and Notes on the Phanerozoic Distribution of Fossil Footprints*; New Mexico Museum of Natural History and Science Bulletin: Albuquerque, NM, USA, 2005; Volume 29, pp. 159–167.
12. Farlow, J.O. A consideration of the trophic dynamics of a Late Cretaceous large-dinosaur community (Oldman Formation). *Ecology* **1976**, *57*, 841–857. [[CrossRef](#)]
13. Fiorillo, A.R. Prey bone utilization by predatory dinosaurs. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **1991**, *88*, 157–166. [[CrossRef](#)]

14. Drumheller, S.K.; McHugh, J.B.; Kane, M.; Riedel, A.; D'Amore, D.C. High frequencies of theropod bite marks provide evidence for feeding, scavenging, and possible cannibalism in a stressed Late Jurassic ecosystem. *PLoS ONE* **2020**, *15*, e0233115. [[CrossRef](#)] [[PubMed](#)]
15. Hunt, A.P.; Lucas, S.G. *The Ichnology of Vertebrate Consumption: Dentalites, Gastroliths and Bromalites*; New Mexico Museum of Natural History and Science Bulletin: Albuquerque, NM, USA, 2021; Volume 87, pp. 1–215.
16. Hunt, A.P.; Lucas, S.G. The Upper Cretaceous fossil record of the Western Interior Basin of North America is a Megalagerstätte. *Geol. Soc. Am. Abstr. Programs* **2022**, *54*. [[CrossRef](#)]
17. Raup, D.M. Taxonomic diversity during the Phanerozoic. *Science* **1972**, *177*, 1065–1071. [[CrossRef](#)]
18. Raup, D.M. Species diversity in the Phanerozoic: A tabulation. *Paleobiology* **1976**, *2*, 279–288. [[CrossRef](#)]
19. Sheehan, P.M. Species diversity in the Phanerozoic. A reflection of labor by systematists? *Paleobiology* **1977**, *2*, 325–328. [[CrossRef](#)]
20. Signor, P.W., III. Species richness in the Phanerozoic: Compensating for sampling bias. *Geology* **1982**, *10*, 625–628. [[CrossRef](#)]
21. Signor, P.W., III. Real and apparent trends in species richness through time. In *Phanerozoic Diversity Patterns: Profiles in Macroevolution*; Valentine, J.W., Ed.; Princeton University Press: Princeton, NJ, USA, 1985; pp. 129–150.
22. Smith, A.B.; McGowan, A.J. The shape of the Phanerozoic marine palaeodiversity curve: How much can be predicted from the sedimentary rock record of Western Europe? *Palaeontology* **2007**, *50*, 765–774. [[CrossRef](#)]
23. Lucas, S.G. Carboniferous tetrapod biostratigraphy, biochronology and evolutionary events. *Geol. Soc. Lond. Spec. Publ.* **2022**, *512*, 965–1001. [[CrossRef](#)]
24. Lucas, S.G. Middle to Late Pennsylvanian tetrapod evolution: The Kasimovian bottleneck. *Geol. Soc. Lond. Spec. Publ.* **2023**, *535*. [[CrossRef](#)]
25. Milner, A.R. The Westphalian tetrapod fauna; some aspects of its geography and ecology. *J. Geol. Soc.* **1987**, *144*, 495–506. [[CrossRef](#)]
26. Clack, J.A.; Milner, A.R. Basal Tetrapoda. In *Handbook of Paleoherpetology 3A1*; Verlag Dr. Friedrich Pfeil: München, Germany, 2015; pp. 1–90.
27. Schutter, S.R.; Heckel, P.H. Missourian (early late Pennsylvanian) climate in midcontinent North America. *Int. J. Coal Geol.* **1985**, *5*, 111–140. [[CrossRef](#)]
28. Cleal, C.J.; Oplustil, S.; Thomas, B.A.; Tenchov, V. Late Moscovian terrestrial biotas and palaeoenvironments of Variscan Euramerica. *Neth. J. Geosci.* **2009**, *88*, 181–278. [[CrossRef](#)]
29. Boucot, A.J.; Xu, C.; Scotese, C.R.; Morley, R.J. *Phanerozoic Paleoclimate: An Atlas of Lithologic Indicators of Climate*; SEPM (Society for Sedimentary Geology): Tulsa, OK, USA, 2013; Volume 11.
30. Hunt, A.P.; Lucas, S.G.; Klein, H. Late Triassic nonmarine vertebrate and invertebrate trace fossils and the pattern of the Phanerozoic record of vertebrate trace fossils. In *The Late Triassic World, Topics in Geobiology 46*; Tanner, L.H., Ed.; Springer: New York, NY, USA, 2018; pp. 447–543.
31. Raup, D.M. Systematists follow the fossils. *Paleobiology* **1977**, *3*, 328–329. [[CrossRef](#)]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.