

Supplemental Information

Removal of Phosphorus and Cadmium from Wastewaters by Periphytic Biofilm

Jin Zhang ^{1,2}, Yawei Liu ^{1,2}, Jiajia Liu ³, Yu Shen ^{1,2}, Hui Huang ^{1,2}, Yongli Zhu ^{1,2}, Jiangang Han ^{1,2,*}
and Haiying Lu ^{1,2,*}

¹ Co-Innovation Center for Sustainable Forestry in Southern China, College of Biology and the Environment, Nanjing Forestry University, Nanjing 210037, China; jinzhang@njfu.edu.cn (J.Z.); ywliu@njfu.edu.cn (Y.L.); shenyu@njfu.edu.cn (Y.S.); huanghui@njfu.edu.cn (H.H.); yonglizhu@njfu.edu.cn (Y.Z.)

² National Positioning Observation Station of Hung-tse Lake Wetland Ecosystem in Jiangsu Province, Nanjing 210037, China

³ Environmental Engineering College, Nanjing Polytechnic Institute, Nanjing 210044, China

* Correspondence: jianganghan@njfu.edu.cn (J.H.); luhaiying@njfu.edu.cn (H.L.)



Figure S1. Experimental device in the experiment (Part). Note, one group for the electricity addition device for the periphyton (A); and the detailed device description image (B).




Figure S2. Concentration of $\text{NH}_4^+\text{-N}$ in solution under different treatments with (A) daily change and (B) monthly change in periphyton. Note, Different letters indicate significant differences between treatments ($p < 0.05$).