

Article

Synthesis of a Novel D-Glucose-Conjugated 15-Crown-5 Ether with a Spiro Ketal Structure

Takashi Yamanoi *, Yoshiki Oda, Hitomi Muraishi and Sho Matsuda

The Noguchi Institute, 1-8-1 Kaga, Itabashi-ku, Tokyo 173-0003, Japan; E-mails: odayoshiki@noguchi.or.jp (Yoshiki Oda), hitommy-heart@bridge.ocn.ne.jp (Hitomi Muraishi), s.matsuda@noguchi.or.jp (Sho Matsuda)

* Author to whom correspondence should be addressed. E-mail: tyama@noguchi.or.jp; Fax: (+81) 3 5944 3213.

Received: 18 July 2008; in revised form: 12 August 2008 / Accepted: 19 August 2008 / Published: 22 August 2008

Abstract: This paper describes a synthetic approach to a novel D-glucose-conjugated 15-crown-5 ether having a spiroketal structure starting from a 1-C-vinylated glucose derivative. The approach consists of the glycosylation of the vinylated glucose derivative to give an ethyleneoxy spacer derivative using bismuth(III) triflate, the conversion of the 1-C-vinyl group of the glucoside produced into a carboxylic acid group, and the intramolecular condensation between the carboxyl group and the terminal hydroxyl group in the ethyleneoxy spacer. A D-glucose-conjugated 15-crown-5 ether having a unique spiroketal structure was thus successfully synthesized.

Keywords: Crown ether; Spiroketal; 1-C-Vinylated glucose; Glycosylation

Introduction

Crown ether molecules with saccharide moieties are interesting as chiral phase-transfer catalysts [1-2]. An enzymatic approach for synthesizing these types of crown ethers provides the cyclofructan family (cyclo β (2 $\rightarrow$ 1)-D-fructooligosaccharides) via the digestion of inulin. The cyclofructan contains a structurally interesting crown ether framework in its central core [3-4]. It is noteworthy that this is the

first example of saccharide-based crown ethers which have spiroketal structures. Many saccharide-based crown ether molecules have also been synthesized by chemical procedures [5-7]. As these chemical methods bind the original hydroxyl groups of the saccharide with an ethyleneoxy spacer, they cannot produce however crown ether compounds having spiroketal structures.

Sugar derivatives (1-*C*-vinylated sugars) having a vinyl group at the anomeric center, which are readily prepared by the addition of organometallic reagents, such as vinylMgX, to a suitably protected sugar lactone, are a synthetically useful tool in carbohydrate chemistry [8-11]. Our recent studies have shown that these 1-*C*-vinylated sugar derivatives were good precursors for preparing some functionalized *exo*-glycal derivatives [12] and naturally occurring anhydroketopyranoses [13]. For the purpose of further exploring the utility of the 1-*C*-vinylated sugars, we investigated the synthesis of a novel crown ether molecule from a 1-*C*-vinylated D-glucose derivative **1**. The D-glucose-conjugated 15-crown-5 ether **2** that we designed is a dicyclic compound with a unique spiroketal structure derived from the structural characteristic of **1**, *i.e.*, its spiro carbon atom corresponds to the anomeric carbon atom. This paper describes our synthetic approach to a novel 15-crown-5 ether **2** having a spiroketal structure from a 1-*C*-vinylated glucose derivative (**1**).

Results and Discussion


The synthetic approach to compound **2** from **1** is shown in Scheme 1. It consists of the following reaction steps: 1) introduction of the ethyleneoxy spacer, tetraethyleneglycol monobenzoate (**3**) onto the vinylated D-glucopyranose derivative **1** by the glycosylation reaction; 2) conversion of the vinyl group at the anomeric center of **4** to a carboxyl group, and 3), intramolecular condensation between the carboxyl group and the terminal hydroxyl group in the ethyleneoxy spacer to produce the desired **2**.

The glycosylation of **1** to **3** (1.3 equiv.) using bismuth(III) triflate (Bi(OTf)₃) (0.05 equiv.) in the presence of anhydrous CaSO₄ in dichloromethane at 0 °C for 24 h afforded the desired glucoside **4** [14], which was purified by preparative TLC (ethyl acetate/hexane = 1/2) in 81% yield. The glycosylation proceeded with an α -stereoselectivity. The high α -stereoselectivity of the glycosylation using **1** was in agreement with our previously reported observation [15]. The α -anomeric configuration of **4** was determined by the NOE interaction between the H-2 and the H-1'.

The ozone oxidation of **4** in dichloromethane at -78 °C for 5 h and treatment with triphenylphosphine (3.4 equiv.) at room temperature for 19 h gave the crude aldehyde product. The subsequent oxidation using NaClO₂ (12 equiv.)-NaH₂PO₄ (3 equiv.) in *t*-butyl alcohol-H₂O (4/1) produced the carboxylic acid derivative **5**, which was purified by preparative TLC (CHCl₃/MeOH = 5/1) in 85% yield.

Deprotection of the benzoyl group of **5** was performed using 0.5 M NaOH/THF to afford **6** in 83% yield. The cyclization of **6** using (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP) (2.5 equiv.) and DIEA (1.8 equiv.) in dichloromethane for 24 h afforded the desired **2**, which was purified by preparative TLC (CHCl₃/MeOH = 20/1) in 84% yield.

In conclusion, we have demonstrated the synthesis of a novel 15-crown-5 ether **2** having a spiroketal structure from a 1-*C*-vinylated D-glucose derivative. This compound **2** is expected to function as a chiral phase-transfer catalyst.

Scheme 1. Synthetic approach to **2**.

Reagents and conditions: i) $\text{Bi}(\text{OTf})_3$ (0.05 equiv.), $\text{Bz}(\text{OCH}_2\text{CH}_2)_4\text{OH}$ **3** (1.3 equiv.), CH_2Cl_2 , 0 °C, 24 h, 81%; ii) (a) O_3 , -78 °C, 5 h, CH_2Cl_2 , then Ph_3P (3.4 equiv.), rt, 19 h. (b) $\text{Me}_2\text{C}=\text{CHMe}$ (4.5 equiv.), NaClO_2 (12 equiv.), NaH_2PO_4 (3 equiv.), $t\text{-BuOH-H}_2\text{O}$, rt, 24 h, 85%; iii) 0.5 M NaOH (13 equiv.), THF, rt, 3 h, 83%; iv) PyBOP (2.5 equiv.), DIEA (1.8 equiv.), CH_2Cl_2 , 24 h, 84%.

Experimental

General

^1H -NMR (600 MHz) and ^{13}C -NMR (150 MHz) spectra were recorded using a JEOL ECA-600 spectrometer in CDCl_3 with TMS as the internal standard. The optical rotations were recorded by a JASCO DIP-360 digital polarimeter. The HRMS were obtained using a Mariner spectrometer (PerSeptive Biosystems Inc.). Preparative TLC was performed using Merck silica gel 60GF254. Column chromatography was conducted using silica gel 60 N (40–50 μm , Kanto Chemical Co., Inc.). $\text{Bi}(\text{OTf})_3$ was purchased from Sigma-Aldrich. All anhydrous solvents were purified according to standard methods.

11-Benzoyloxy-3,6,9-trioxaundecyl 2,3,4,6-tetra-O-benzyl-1-C-vinyl- α -D-glucopyranoside (4**):** To a stirred solution of $\text{Bi}(\text{OTf})_3$ (15 mg, 0.023 mmol) in CH_2Cl_2 (3.5 mL) were added tetraethyleneglycol monobenzoate (**3**) (165 mg, 0.55 mmol) and 2,3,4,6-tetra-O-benzyl-1-C-vinyl- α -D-glucopyranose (**1**) (257 mg, 0.42 mmol) in the presence of anhydrous CaSO_4 (280 mg) under an Ar atmosphere. The resulting mixture was stirred at 0 °C for 24 h. The reaction was then quenched by the addition of a sat.

NaHCO₃ solution (5 mL). The reaction mixture was extracted with CH₂Cl₂ (three times), and the organic layer was washed with water and a sat. NaCl solution. After the organic layer was dried over Na₂SO₄, the solvent was evaporated under reduced pressure. The crude product was purified by preparative silica gel TLC (ethyl acetate/hexane = 1/2) to give **4** (311 mg, 81% yield) as a colorless oil. $[\alpha]_D^{25} = +3^\circ$ (*c* 4.7, CHCl₃); ¹H-NMR: δ 3.34 (d, 1H, *J* = 9.6 Hz, H-2), 3.48–3.70 (m, 14H, H-4, H_a-6, CH₂CH₂), 3.76–3.79 (m, 3H, H_b-6, CH₂CH₂), 3.85 (m, 1H, H-5), 4.10 (t, 1H, *J* = 9.7 Hz, H-3), 4.43–4.91 (m, 8H, CH₂Ph), 5.27 (dd, 1H, *J* = 2.0 Hz, *J* = 11.0 Hz, CH=CH_aH_b), 5.54 (dd, 1H, *J* = 2.1 Hz, *J* = 17.9 Hz, CH=CH_aH_b), 5.99 (m, 1H, CH=CH₂), 7.19–7.54, 8.04–8.05 (m, 25H, Ph); ¹³C-NMR: δ 61.1 (CH₂CH₂), 64.1 (CH₂CH₂), 68.8 (C-6), 69.2 (CH₂CH₂), 70.0 (CH₂CH₂), 70.6 (CH₂CH₂), 70.64 (CH₂CH₂), 70.7 (CH₂CH₂), 71.5 (C-5), 73.4 (CH₂Ph), 75.0 (CH₂Ph), 75.5 (CH₂Ph), 75.8 (CH₂Ph), 78.5 (C-4), 83.0 (C-3), 84.3 (C-2), 99.5 (C-1), 118.8 (CH=CH₂), 127.5–133.0 (Ph), 135.3 (CH=CH₂), 138.1–138.4 (Ph), 166.5 (C=O); HRMS (ESI) *m/z* calcd for C₅₁H₅₈NaO₁₁ 869.3871 [M+Na]⁺, found 869.3865.

(11-Benzoyloxy-3,6,9-trioxaundecyl 3,4,5,7-tetra-*O*-benzyl- α -D-glucopyranosid)onic acid (**5**): Ozone was bubbled through a stirred solution of **4** (224 mg, 0.26 mmol) in CH₂Cl₂ (15 mL) at -78 °C for 5 h. After triphenylphosphine (230 mg, 0.88 mmol) was added at -78 °C and the reaction temperature was raised to room temperature, the reaction mixture was stirred for 19 h. The solvent was then evaporated under reduced pressure. To a solution of the crude product in *t*-butyl alcohol (4 mL)-H₂O (1 mL) were added NaClO₂ (277 mg, 3.1 mmol), NaH₂PO₄ (124 mg, 0.8 mmol) and 2-methyl-2-butene (123 μ L, 1.2 mmol). After the reaction mixture was stirred for 24 h, the reaction was quenched by adding 2 M HCl (1 mL) and water (5 mL). The reaction mixture was then extracted with CH₂Cl₂ (three times), and the combined organic solvent was dried over anhydrous Na₂SO₄. The organic solvent was filtered and evaporated under reduced pressure. The crude product was purified by preparative silica gel TLC (CHCl₃/MeOH = 5/1) to afford **5** (194 mg, 85% yield) as a colorless oil. $[\alpha]_D^{25} = +21^\circ$ (*c* 3.9, CHCl₃); ¹H-NMR: δ 3.54–4.08 (m, 20H, H-3, H-4, H-5, H-6, H-7, CH₂CH₂), 4.41–4.49 (m, 2H, CH₂CH₂), 4.51–5.28 (m, 8H, CH₂Ph), 7.03–7.53 (m, 23H, Ph), 8.03–7.53 (d, 2H, *J* = 6.8 Hz, Ph); ¹³C-NMR: δ 64.0 (CH₂CH₂), 69.1 (C-7), 69.9 (CH₂CH₂), 70.2 (CH₂CH₂), 70.3 (CH₂CH₂), 70.4 (CH₂CH₂), 70.5 (CH₂CH₂), 70.6 (CH₂CH₂), 72.6 (CH₂CH₂), 75.19 (CH₂Ph), 75.20 (CH₂Ph), 75.4 (CH₂Ph), 75.5 (CH₂Ph), 77.6 (C-6), 80.9 (C-5), 82.7 (C-3, C-4), 99.3 (C-2), 126.0–139.2 (Ph), 166.5 (C(O)Ph), 177.7 (C-1); HRMS (ESI) *m/z* calcd for C₅₀H₅₆NaO₁₃ 887.3613 [M+Na]⁺, found 887.3653.

(11-Hydroxy-3,6,9-trioxaundecyl 3,4,5,7-tetra-*O*-benzyl- α -D-glucopyranosid)onic acid (**6**): A 0.5 M NaOH solution (4 mL, 2 mmol) was added to a solution of **5** (142 mg, 0.16 mmol) in THF (4 mL). After the reaction mixture was stirred for 3 h at room temperature, the reaction was quenched by adding 2 M HCl (1 mL) and water (5 mL). After the reaction mixture was extracted with CH₂Cl₂ (three times), the combined organic solvent was dried over anhydrous Na₂SO₄. The organic solvent was filtered and evaporated under reduced pressure. The crude product was purified by preparative silica gel TLC (CHCl₃/MeOH = 5/1) to afford **6** (103 mg, 83% yield) as a colorless oil. $[\alpha]_D^{25} = +25^\circ$ (*c* 1.8, CHCl₃); ¹H-NMR: δ 3.37–4.00 (m, 22H, H-3, H-4, H-5, H-6, H-7, CH₂CH₂), 4.44–4.84 (m, 8H, CH₂Ph), 6.94–7.43 (m, 20H, Ph); ¹³C-NMR: δ 60.4 (CH₂CH₂), 62.6 (CH₂CH₂), 68.5–70.4 (CH₂CH₂, C-7), 72.4 (CH₂Ph), 73.4 (CH₂Ph), 74.9 (CH₂Ph), 75.3 (CH₂Ph), 78.1 (C-6), 82.4 (C-5), 82.9 (C-3, C-4),

99.7 (C-2), 127.3-128.4, 137.8-138.9 (Ph), 172.3 (C-1); HRMS (ESI) m/z calcd for $C_{43}H_{52}NaO_{12}$ 783.3351 $[M+Na]^+$, found 783.3396.

(1*R*)-2,3,4,6-Tetra-*O*-benzylspiro[1,5-anhydro-D-glucitol-1,2'-[3,6,9,12]tetraoxatetradecan]-14'-olide (2): To a solution of **6** (20 mg, 0.027 mmol) in CH_2Cl_2 (3 mL) were added 4-dimethylaminopyridine (5.9 mg, 0.048 mmol) and PyBOP (35 mg, 0.067 mmol). After the reaction mixture was stirred for 24 h. The reaction was then quenched by the addition of a sat. citric acid solution (5 mL). The reaction mixture was extracted with EtOAc and the organic layer was washed with water and a sat. NaCl solution. After the organic layer was dried over Na_2SO_4 , the solvent was evaporated under reduced pressure. The crude product was purified by preparative silica gel TLC ($CHCl_3/MeOH = 20/1$) to give **2** (17 mg, 84% yield) as a colorless oil. $[\alpha]_D^{25} = +11^\circ$ (c 0.15, $CHCl_3$); 1H -NMR: δ 3.48-3.71 (m, 15H, H-4, H-5, H-6, CH_2CH_2), 3.73-3.75 (m, 1H, CH_2CH_2), 3.81 (d, 1H, $J = 9.6$ Hz, H-2), 3.87-3.92 (m, 1H, CH_2CH_2), 3.94-3.98 (m, 1H, CH_2CH_2), 4.00-4.05 (m, 1H, $CH_aH_bCH_2$), 4.06-4.07 (m, 1H, H-3), 4.31-4.34 (m, 1H, $CH_aH_bCH_2$), 4.54-4.65 (m, 4H, CH_2Ph), 4.79-4.89 (m, 4H, CH_2Ph), 7.16-7.35 (m, 20H, Ph); ^{13}C -NMR: δ 63.6 (CH_2CH_2), 65.2 (CH_2CH_2), 68.3 (CH_2CH_2), 68.4 (C-6), 69.6 (CH_2CH_2), 70.1 (CH_2CH_2), 70.4 (CH_2CH_2), 70.9 (CH_2CH_2), 71.2 (CH_2CH_2), 73.41 (C-5), 73.44 (CH_2Ph), 75.1 (CH_2Ph), 75.2 (CH_2Ph), 75.6 (CH_2Ph), 78.0 (C-4), 82.2 (C-2), 82.7 (C-3), 99.7 (C-1), 127.5-128.4 (Ph), 137.9-138.5 (Ph), 173.5 (C-1'); HRMS (ESI) m/z calcd for $C_{43}H_{50}NaO_{11}$ 765.3245 $[M+Na]^+$, found 765.3247.

References and Notes

1. Mako, A.; Szollosy, A.; Keglevich, G.; Menyhard, D. K.; Bako, P.; Toke, L. Synthesis of methyl- α -D-glucopyranoside-based azacrown ethers and their application in enantioselective reactions. *Monatsh. Chem.* **2008**, *139*, 525-535. Their other references cited therein.
2. Itoh, T.; Shirakami, S. Synthesis of chiral crown ethers derived from α -D-glucose and their catalytic properties on the asymmetric michael addition. *Heterocycles* **2001**, *55*, 37-43.
3. Takai, Y.; Okumura, Y.; Tanaka, T.; Sawada, M.; Takahashi, S.; Shiro, M.; Kawamura, M.; Uchiyama, T. Binding characteristics of a new host family of cyclic oligosaccharides from inulin: Permethylated cycloinulohexaose and cycloinulohepaose. *J. Org. Chem.* **1994**, *59*, 2967-2975. Their other references cited therein.
4. It has also been reported the cyclo(2 $\rightarrow$ 1)-D-fructodisaccharides have spiro ketal structures. Louis, F.; Garcia-Moreno, M. I.; Balbuena, P.; Mellet, C. O.; Garcia-Fernandez, J. M. Stereoselective synthesis of nonsymmetrical difructose dianhydrides from xylylene-tethered D-fructose precursors. *Tetrahedron* **2008**, *64*, 2792-2800.
5. Mani, N. S.; Kanakamma, P. P. Synthesis of novel chiral macrocycles: Crown ethers derived from D-glucose. *Tetrahedron Lett.* **1994**, *35*, 3629-3632.
6. Faltin, F.; Fehring, V.; Miethchen, R. Chiral crown ethers based on galactopyranosides. *Synthesis* **2002**, 1851-1856 and the numerous references cited therein.
7. Dumont-Hornebeck, B.; Joly, J.P.; Coulon, J.; Chapleur, Y. Synthesis of ethoxy-linked pseudo-disaccharides incorporating a crown ether macrocycles and lectin recognition. *Carbohydr. Res.* **1999**, *321*, 214-227.

8. Tomooka, K.; Nakamura, Y.; Nakai, T. [2, 3]-Wittig rearrangement using glucose as a chiral auxiliary: asymmetric transmission from the anomeric center. *Synlett*. **1995**, 321-322.
9. Lay, L.; Meldal, M.; Nicotra, F.; Panza, L.; Russo, G. Stereoselective synthesis of the C-analogue of β -D-glucopyranosyl serine. *Chem. Commun.* **1997**, 54, 1469-1470.
10. van Hooft, P. A. V.; Oualid, F. E.; Overkleeft, H. S.; van der Marel, G. A.; van Boom, J. H.; Leeuwenburgh, M. A. Synthesis and elaboration of functionalised carbohydrate-derived spiroketals. *Org. Biomol. Chem.* **2004**, 2, 1395-1403.
11. Li, X. L.; Ohtake, H.; Takahashi, H.; Ikegami, S. A facile synthesis of 1'-C-alkyl- α -disaccharides from 1-C-alkyl-hexopyranoses and methyl 1-C-methyl-hexopyranosides. *Tetrahedron* **2001**, 57, 4297-4309.
12. Yamanoi, T.; Nara, Y.; Matsuda, S.; Oda, Y.; Yoshida, A.; Katsuraya, K.; Watanabe, M. Synthetic approach to *exo*-glycals from 1-C-vinyl-D-glycopyranose derivatives via an S_N1' -substitution mechanism. *Synlett*. **2007**, 785-789.
13. Matsuda, S.; Matsumura, K.; Watanabe, M.; Yamanoi, T. Synthesis of a partially benzylated derivative of the anhydro-D-altro-heptulose found in *Coriaria japonica* A. *Tetrahedron Lett.* **2007**, 48, 5807-5810.
14. Yamanoi, T.; Inoue, R.; Matsuda, S.; Katsuraya, K.; Hamasaki, K. Synthesis of trehalose mimics by bismuth(III) triflate or bis(trifluoromethane)sulfonimide-catalyzed 1-C-methyl-D-hexopyranosylation. *Tetrahedron Asymmetry* **2006**, 17, 2914-2918.
15. Yamanoi, T.; Oda, Y.; Matsuda, S.; Yamazaki, I.; Matsumura, K.; Katsuraya, K.; Watanabe, M.; Inazu, T. Synthesis of 1-C-alkyl- α -D-glucopyranosides by Lewis acid- or Brønsted acid-catalyzed O-glycosidation. *Tetrahedron* **2006**, 62, 10383-10392.

Sample Availability: Samples of the compounds are available from the authors

© 2008 by the authors; licensee Molecular Diversity Preservation International, Basel, Switzerland. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).