

Article

## A Mild and Facile One-Pot Synthesis of *N*-Methyl-3-Acyl-Pyrroles

Hassan Valizadeh \* and Ashraf Fakhari

Department of Chemistry, Faculty of Sciences, Azarbaijan University of Tarbiat Moallem, Tabriz, Iran

\* Author to whom correspondence should be addressed; E-mail: h-valizadeh@azaruniv.edu;

Fax: +98-412-432-7541.

Received: 28 February 2010; in revised form: 23 March 2010 / Accepted: 18 April 2010 /

Published: 27 April 2010

---

**Abstract:** *N*-Methyl-3-acylpyrroles were synthesized via a multicomponent reaction of dimethylacetylene dicarboxylate (DMAD), *N*-methylhydroxylamine and acylchlorides in the presence of  $\text{KHCO}_3$ . For comparison both conventional and microwave protocols were examined in this procedure. The reaction is clean and gives the products in good to excellent yields under conventional heating conditions at 40 °C in anhydrous dichloromethane.

**Keywords:** 3-acylpyrrole; multicomponent reaction; dimethylacetylene dicarboxylate

---

### 1. Introduction

Pyrrole is one of the most important heterocyclic compounds, having become increasingly important in medicinal chemistry and organic synthesis [1-5]. Some of the recently isolated pyrrole-containing marine natural products have been found to exhibit considerable cytotoxicity and function as multi drug resistant (MDR) reversal agents. Many of these biologically active compounds have emerged as chemotherapeutic agents [6]. Heterocyclic compounds containing a 3-acylpyrrole fragment are of interest for making new pharmacological preparations. For example, the cannabinoid activity of 1-alkyl-3-(naphthoyl)pyrroles [7] and 1-alkyl-3-(naphthoyl)indoles [8] is known, as is the antibiotic activity of verrucarins (3-acetyl-4-hydroxymethylpyrrole) isolated from *Myrothecium verrucaria* [9]. 3-Acylpyrroles and other 3-substituted pyrroles obtained from them are precursors of liquid crystal materials [10] and polypyrroles [11], possessing high electrical conductivity in comparison with their


1-substituted analogs [12]. Such conjugated polymers may be used, for example, to construct gas sensors [13], and also sensors capable of distinguishing DNA molecules [14]. Many methods for the synthesis of diversely substituted pyrroles have been developed [15]. Conjugate addition reactions [16], transition metal-mediated reactions [17], reductive couplings [18], aza-Wittig reactions [19], and other multistep operations [20] have all been performed for the synthesis of pyrroles. Despite these huge developments, the Paal-Knorr [21] reaction is still considered to be the most attractive method for the synthesis of pyrroles. In the present work, we describe a simple method for the  $\text{KHCO}_3$ -catalyzed synthesis of 3-acylpyrroles in good yields starting from commercially available DMAD, *N*-methylhydroxylamine and various acylchlorides.

## 2. Results and Discussion

Our research group has reported several studies directed toward the synthesis of some heterocyclic compounds under solvent-free conditions [22-27]. Because of the biological activities of pyrrole derivatives, we became interested in the one-pot synthesis of *N*-methyl-3-acylpyrroles under mild and solvent-free conditions.

Electrophilic substitution in pyrrole occurs predominantly at the  $\text{C}_2$ -position [28-30]. Thus, the investigation of efficient methods for preparing  $\text{C}_3$ -substituted pyrroles is one of the important goals in pyrrole chemistry because of their frequent use for obtaining various biologically active compounds like porphyrins. For the substitution on  $\beta$ -( $\text{C}_3$ ) position, Friedel-Crafts acylation or alkylation on pyrrole bearing an electron withdrawing substituent at  $\text{C}_2$  [31-32] or  $\text{N}_1$  [33-34] has been widely investigated. Direct Friedel-Crafts acylation on 2,5-dimethylpyrrole also gives 3,4-symmetric acyl compounds, but normally the yields are very low (< 5%) when no electron withdrawing groups are on the substituted pyrroles [18]. Very recently we prepared pentasubstituted pyrroles via the reaction of DMAD with *N*-methylhydroxylamine under mild solvent-free conditions, so we studied the possibility of synthesizing *N*-methyl-3-acylpyrroles **4a-e** using a multicomponent reaction in anhydrous dichloromethane of DMAD (**1**), *N*-methylhydroxylamine (**2**) and various acylchlorides **3a-e** in the presence of  $\text{KHCO}_3$  (Scheme 1).

**Scheme 1.** One-pot synthesis of *N*-methyl-3-acylpyrroles.



We tried several different reaction conditions under microwave irradiation and conventional heating, and found that the reaction took place efficiently and in good yield at 40 °C in  $\text{CH}_2\text{Cl}_2$ . Firstly, benzoyl chloride (**3a**) was chosen as a model for the reaction with DMAD and *N*-

methylhydroxylamine to optimize reaction conditions. It was observed that a low yield of **4a** was obtained in the absence of  $\text{KHCO}_3$  when the reaction was stirred at room temperature for 10 h. The reaction in the presence of the  $\text{KHCO}_3$  at room temperature in  $\text{CH}_2\text{Cl}_2$ , afforded the corresponding product in 61% yield (Table 1, entry 1). The reaction was also examined at 40 °C and found to be faster and to give higher yield (81%) than at room temperature (Table 1, entry 2). The procedure was examined under microwave-assisted conditions and found that the yield of product was very low in comparison with conventional heating in anhydrous  $\text{CH}_2\text{Cl}_2$  (Table 1, entries 7-8). In another experiment, for the same reaction, several common solvents were examined in the presence of  $\text{KHCO}_3$  at 40 °C. The results showed that the reaction in  $\text{CH}_2\text{Cl}_2$  resulted in higher yields in comparison with the other tested solvents (Table 1, entries 2-5). There is no significant difference in the results using different bases such as  $\text{Na}_2\text{CO}_3$ ,  $\text{K}_2\text{CO}_3$ ,  $\text{KHCO}_3$ ,  $\text{NH}_4\text{OAc}$ ,  $\text{NaOH}$ , and  $\text{NH}_4\text{Cl}$  as the catalyst in this procedure. The reaction of enamine, derived from addition of an *N*-methylhydroxylamine to DMAD, and benzoylchloride proceeds by a smooth 1:1:1 addition reaction in anhydrous dichloromethane at 40°C to produce *N*-methyl-3-benzoylpyrrole (**4a**) in 81% yield.

**Table 1.** Study of the synthesis of 3-benzoylpyrrole **4a** under different conditions.

Entry	Solvent/Solid support	Mode of heating	Temperature (°C)	Time (h)	Yield <sup>a</sup> (%)
1	$\text{CH}_2\text{Cl}_2$	Conventional	25	3	61
2	$\text{CH}_2\text{Cl}_2$	Conventional	40	2.5	81
3	$\text{CHCl}_3$	Conventional	40	3	63
4	Dioxane	Conventional	40	2.5	60
5	THF	Conventional	40	2.5	63
6	Neat	Conventional	40	3	53
7	Neat	MW	Not detected	10 min	<30
8	Basic alumina	MW	Not detected	10 min	<30

<sup>a</sup> Isolated yields after column chromatography.

Having established the reaction conditions, various acyl chlorides were examined in this procedure to investigate the reaction scope, and several representative examples are summarized in Table 2. The reaction was clean and proceeded smoothly to give the corresponding 3-acylpyrroles in good to excellent yields. There were no remarkable differences in yields and reaction times between aromatic and aliphatic acyl halides in this procedure.

**Table 2.** One-pot synthesis of 3-acylpyrroles.


Entry	Product	Time (h)	Yield <sup>a</sup> (%)
1	<b>4a</b>	2.5	81
2	<b>4b</b>	3	83
3	<b>4c</b>	3	73
4	<b>4d</b>	3.5	78
5	<b>4e</b>	2.5	76

<sup>a</sup> Isolated yields after column chromatography.

The structures of compounds **4a–4e** were deduced from their elemental analyses and their IR,  $^1\text{H}$ -NMR, and  $^{13}\text{C}$ -NMR spectroscopic data and no side products were identified. For example, the  $^1\text{H}$ -NMR spectrum of **4a** exhibited one singlet identified as the *N*-methyl moiety ( $\delta$  3.50), phenyl proton multiplets ( $\delta$  7.30, 3H and  $\delta$  7.41, 2H), along with multiplets [ $\delta$  7.51, 1H,  $\delta$  7.60, 1H and  $\delta$  7.81, 1H), for the pyrrole protons. The  $^1\text{H}$ -decoupled  $^{13}\text{C}$ -NMR spectrum of **4a** showed 10 distinct resonances that confirm the proposed structure. The IR spectrum of **4a** displayed a characteristic carbonyl ( $1,732\text{ cm}^{-1}$ ) stretching vibration.

Although the mechanistic details of the reaction are not known, a plausible rationalization may be advanced to explain the product formation. Presumably, the enamine **5** formed from *N*-methylhydroxylamine and DMAD attacked to acyl chloride to furnish intermediate **6**, which is converted to intermediate **7**. Subsequent reaction of **7** with DMAD yields intermediate **8**, which undergoes an intramolecular cyclization reaction to generate **9**. Finally, products **4a–e** were formed from intermediate **9**.

**Scheme 2.** Plausible mechanism for the formation of the products **4a–e**.



### 3. Experimental

#### 3.1 General

All reagents were purchased from Merck and used without further purification.  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR spectra were recorded on a Bruker Avance AC-400 MHz instrument using  $\text{CDCl}_3$  as the deuterated solvent and TMS as internal standard. Elemental analyses were carried out on a Perkin-Elmer 240C

elemental analyzer and are reported in percent atomic abundance. All melting points are uncorrected and measured in open glass-capillaries using Stuart melting point apparatus. Microwave experiments were conducted in a Milestone MicroSYNTH apparatus.

### 3.2. General procedure for preparation of 3-acylpyrroles **4a-e**

Dimethylacethylene dicarboxylate (2 mmol) was added to a stirred solution of *N*-methylhydroylamine hydrochloride (2 mmol) and  $\text{KHCO}_3$  (3 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (10 mL). The mixture was stirred for 15 min, and then a solution of an acyl chloride (2 mmol) and dimethylacethylene dicarboxylate (2 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) was added slowly at 40 °C. After completion of the reaction (2.5–3.5 h) as indicated by TLC on silica gel F<sub>254</sub> (*n*-hexane/EtOAc 6:1), the solvent from the reaction mixture was evaporated under reduced pressure to leave a residue that was purified by column chromatography (*n*-hexane/EtOAc 6:1) to afford pure desired products **4a-e**.

*N*-Methyl-3-benzoyl pyrrole (**4a**): IR (KBr) ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ ): 1732 (C=O);  $^1\text{H-NMR}$   $\delta_{\text{H}}$  (ppm): 7.81 (m, 1H), 7.60 (m, 1H), 7.51 (m, 1H), 7.41 (m, Ph-H, 2H), 7.30 (m, Ph-H, 3H), 3.50 (s, 3H, N-Me);  $^{13}\text{C-NMR}$   $\delta_{\text{C}}$  (ppm): 163.32, 133.14, 132.24, 129.92, 128.77, 127.62, 127.10, 126.84, 125.75, 36.24; Anal. Calcd (%) for  $\text{C}_{12}\text{H}_{11}\text{NO}$ : C, 77.81; H, 5.99; N, 7.56. Found (%): C, 77.90; H, 5.91; N, 7.48.

*N*-Methyl-3-acetyl pyrrole (**4b**): IR (KBr) ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ ): 1727 (C=O);  $^1\text{H-NMR}$   $\delta_{\text{H}}$  (ppm): 7.75 (m, 1H), 7.48-7.55 (m, 2H), 3.53 (s, 3H, N-Me), 2.09 (s, 3H, COMe);  $^{13}\text{C-NMR}$   $\delta_{\text{C}}$  (ppm): 160.22, 131.33, 129.12, 126.12, 124.78, 33.45, 27.50; Anal. Calcd (%) for  $\text{C}_7\text{H}_9\text{NO}$ : C, 68.27; H, 7.37; N, 11.37. Found (%): C, 69.05; H, 7.29; N, 11.31.

*N*-Methyl-3-(4-chlorobenzoyl)-pyrrole (**4c**): IR (KBr) ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ ): 1720 (C=O);  $^1\text{H-NMR}$   $\delta_{\text{H}}$  (ppm): 7.77 (m, 1H), 7.51-7.58 (m, 2H), 7.38 (d,  $J=7.56\text{Hz}$ , 2H), 7.31 (d,  $J=7.56\text{Hz}$ , 2H), 3.47 (s, 3H, N-Me);  $^{13}\text{C-NMR}$   $\delta_{\text{C}}$  (ppm): 164.56, 135.66, 134.32, 131.11, 129.32, 128.09, 127.99, 126.25, 126.15, 33.36; Anal. Calcd (%) for  $\text{C}_{12}\text{H}_{10}\text{ClNO}$ : C, 65.61; H, 4.59; N, 6.38. Found (%): C, 65.65; H, 4.53; N, 6.34.

*N*-Methyl-3-(2-methoxybenzoyl)-pyrrole (**4d**): IR (KBr) ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ ): 1721 (C=O);  $^1\text{H-NMR}$   $\delta_{\text{H}}$  (ppm): 7.83 (m, 1H), 7.49-7.54 (m, 2H), 7.31-7.38 (m, Ph-H, 4H), 3.87 (s, 3H, OMe), 3.50 (s, 3H, N-Me);  $^{13}\text{C-NMR}$   $\delta_{\text{C}}$  (ppm): 162.21, 137.62, 136.54, 135.66, 134.10, 133.22, 132.20, 130.34, 129.79, 127.40, 126.15, 67.71, 34.16; Anal. Calcd (%) for  $\text{C}_{13}\text{H}_{13}\text{NO}_2$ : C, 72.54; H, 6.09; N, 6.51. Found (%): C, 72.58; H, 5.98; N, 6.48.

*N*-Methyl-3-(4-nitrobenzoyl)-pyrrole (**4e**): IR (KBr) ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ ): 1717 (C=O);  $^1\text{H-NMR}$   $\delta_{\text{H}}$  (ppm): 7.70 (m, 1H), 7.66 (d,  $J=7.87\text{Hz}$ , 2H), 7.45-7.53 (m, 2H), 7.37 (d,  $J=7.87\text{Hz}$ , 2H), 3.54 (s, 3H, N-Me);  $^{13}\text{C-NMR}$   $\delta_{\text{C}}$  (ppm): 160.06, 133.06, 133.02, 131.21, 129.70, 128.56, 127.65, 127.05, 126.55, 37.09; Anal. Calcd (%) for  $\text{C}_{12}\text{H}_{10}\text{N}_2\text{O}_3$ : C, 62.60; H, 4.38; N, 12.17. Found (%): C, 62.80; H, 4.35; N, 12.08.

#### 4. Conclusions

In summary, we have described a highly efficient one-pot procedure for the preparation of *N*-methyl-3-acylpyrroles by a three component reaction in CH<sub>2</sub>Cl<sub>2</sub> using KHCO<sub>3</sub> as catalyst. The reaction products were prepared in moderate to good yield, even with different substituted acyl halides. We believe that this procedure provides a valuable addition to current methodologies.

#### Acknowledgements

The partial financial assistance from the Research Vice Chancellor of Azarbaijan University of Tarbiat Moallem is gratefully acknowledged.

#### References and Notes

1. Lehedu, J.; Fauconneau, B.; Barrier, L.; Ourakow, M.; Piriou, A.; Vierfond, J.M. Synthesis and antioxidant activity of new tetraarylpyrroles. *Eur. J. Med. Chem.* **1999**, *34*, 991-996.
2. Raimondi, M.V.; Cascioferro, S.; Schillaci, D.; Petruso, S. Synthesis and antimicrobial activity of new bromine-rich pyrrole derivatives related to monodeoxyphytoluteorin. *Eur. J. Med. Chem.* **2006**, *41*, 1439-1445.
3. Cara, L.C.L.; Camacho, M.E.; Carrión, M.D.; Tapias, V.; Gallo, M.A.; Escames, G.; Acuña-Castroviejo, D.; Espinosa, A.; Entrena, A. Phenylpyrrole derivatives as neural and inducible nitric oxide synthase (nNOS and iNOS) inhibitors. *Eur. J. Med. Chem.* **2009**, *44*, 2655-2666.
4. Abu-Rabeah, K.; Marks, R.S. Impedance study of the hybrid molecule alginate-pyrrole: Demonstration as host matrix for the construction of a highly sensitive amperometric glucose biosensor. *Sens. Actuat. B. Chem.* **2009**, *136*, 516-522.
5. Liu, J.-H.; Yang, Q.-C.; Mak, T.C.W.; Wong, H.N.C. Highly regioselective synthesis of 2,3,4-trisubstituted 1*H*-pyrroles: A formal total synthesis of lukianol A. *J. Org. Chem.* **2000**, *65*, 3587-3595.
6. Tao, H.; Hwang, I.; Boger, D.L. Multidrug resistance reversal activity of permethyl ningalin B amide derivatives. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 5979-5981.
7. Lainton, A.H.; Huffman, J.W. 1-Alkyl-3-(1-naphthoyl)pyrroles: A new class of cannabinoid. *Tetrahedron Lett.* **1995**, *36*, 1401-1404.
8. Huffman, J.W.; Dai, D.; Martin, B.R.; Compton, D.R. Design, synthesis and pharmacology of cannabimimetic indoles. *Bioorg. Med. Chem. Lett.* **1994**, *4*, 563-566.
9. Pfäffli, P.; Tamm, C. Revidierte struktur von verrucarin E. Eine synthese des antibioticums und verwandter  $\beta$ -acetyl-pyrrol-derivate. Verrucarine und roridine. *Helv. Chim. Acta* **1969**, *52*, 1911-1920.
10. Langley, P.J.; Davis, F.J.; Mitchell, G.R. Approaches to 1,1-disubstituted cinnolin-3-ylidene oxides: synthesis and reactivity of a new class of heterocyclic betaines. *J. Chem. Soc. Perkin. Trans.* **1997**, *2*, 2229-2236.
11. Ruhe, J.; Ezquerra, T.A.; Wegner, G. New conducting polymers from 3-alkylpyrroles. *Synth Met.* **1989**, *28*, 177-181.

12. Delabouglise, D.; Roncali, J.; Lemaire, M.; Garnier, F. Control of the lipophilicity of polypyrrole by 3-alkyl substitution. *J. Chem. Soc. Chem. Commun.* **1989**, 475-477.
13. de Lacy Costello, B.P.J.; Evans, P.; Guernion, N.; Ratcliffe, N.M.; Sivanand, P.S.; Teare, G.C. The synthesis of a number of 3-alkyl and 3-carboxy substituted pyrroles; their chemical polymerisation onto poly(vinylidene fluoride) membranes, and their use as gas sensitive resistors. *Synth. Met.* **2000**, *114*, 181-188.
14. Garnier, F.; Korri-Youssoufi, H.; Srivastava, P.; Mandrand, B.; Delair, T. Toward intelligent polymers: DNA sensors based on oligonucleotide-functionalized polypyrroles. *Synth. Met.* **1999**, *100*, 89-94.
15. Gilchrist, T.L. Synthesis of aromatic heterocycles. *J. Chem. Soc. Perkin. Trans.* **1998**, *1*, 615-628.
16. Dieter, R.K.; Yu, H. A facile synthesis of polysubstituted pyrroles. *Org. Lett.* **2000**, *2*, 2283-2286.
17. Iwasawa, N.; Maeyama, K.; Saitou, M. Reactions of propargyl metallic species generated by the addition of alkynyllithiums to fischer-type carbene complexes. *J. Am. Chem. Soc.* **1997**, *119*, 1486-1487.
18. Furstner, A.; Weintritt, H.; Hupperts, A. A new, titanium-mediated approach to pyrroles: First synthesis of lukianol A and lamellarin O dimethyl ether. *J. Org. Chem.* **1995**, *60*, 6637-6641.
19. Katritzky, A.; Jiang, J.; Steel, P.J. 1-Aza-1,3-bis(triphenylphosphoranylidene)propane: A Novel :CHCH<sub>2</sub>N: Synthone. *J. Org. Chem.* **1994**, *59*, 4551-4555.
20. Periasamy, M.; Srinivas, G.; Bharati, P. Conversion of aryl methyl ketimines to 2,5-diarylpyrroles using TiCl<sub>4</sub>/Et<sub>3</sub>N. *J. Org. Chem.* **1999**, *64*, 4204-4205.
21. Chen, J.; Wu, H.; Zheng, Z.; Jin, C.; Zhang, X.; Su, W. An approach to the Paal–Knorr pyrroles synthesis catalyzed by Sc(OTf)<sub>3</sub> under solvent-free conditions. *Tetrahedron Lett.* **2006**, *47*, 5383-5387.
22. Valizadeh, H.; Shockravi, A. An efficient procedure for the synthesis of coumarin derivatives using TiCl<sub>4</sub> as catalyst under solvent-free conditions. *Tetrahedron Lett.* **2005**, *46*, 3501-3503.
23. Valizadeh, H.; Mamaghani, M.; Badrian, A. Effect of microwave irradiation on reaction of arylaldehyde derivatives with some active methylene compounds in aqueous media. *Synth. Commun.* **2005**, *35*, 785-790.
24. Valizadeh, H.; Vaghefi, H.; One-pot Wittig and Knoevenagel reactions in ionic liquid as convenient methods for the synthesis of coumarin derivatives. *Synth. Commun.* **2009**, *39*, 1666-1678.
25. Shockravi, A.; Shargi, H.; Valizadeh, H.; Heravi, M.M. Solvent free synthesis of coumarins. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2002**, *177*, 2555-2559.
26. Valizadeh, H.; Amiri, M.; Gholipur, H. Efficient and convenient method for the synthesis of isoxazoles in ionic liquid. *J. Heterocycl. Chem.* **2009**, *46*, 108-110.
27. Valizadeh, H.; Heravi, M.M.; Amiri, M. Unexpected synthesis of *N*-methylbenzo[d]isoxazolium hydroxides under microwave irradiation conditions. *Mol. Divers.* **2009**, doi:10.1007/s11030-009-9189-x.
28. Jones, R.A. *Advances in Heterocyclic Chemistry*; Katritzky, A.R., Boulton, A.J., Eds.; Academic Press: New York, NY, USA, 1970; Volume 11, p. 383.
29. Marino, G. *Advances in Heterocyclic Chemistry*; Katritzky, A.R., Boulton, A.J., Eds.; Academic Press: New York, NY, USA, 1970; Volume 13, p. 235.

30. Bird, C.W.; Cheesman, G.W.H. *Comprehensive Heterocyclic Chemistry*; Katritzky, A., Rens, C.W., Eds.; Pergamon Press: Oxford, UK, 1984; Volume 4, p. 39.
31. Cooper, G. Cyclopropyl 2-pyrrolyl ketone. *J. Org. Chem.* **1971**, *36*, 2897.
32. Tani, M.; Ariyasu, T.; Nishiyama, C.; Hagiwara, H.; Watanabe, T.; Yokoyama, Y.; Murakami, Y.  $\beta$ -Acylation of ethyl pyrrole-2-carboxylate by Friedel-Crafts acylation : Scope and limitations (synthetic studies on indoles and related compounds. XXXVIII). *Chem. Pharm. Bull.* **1996**, *44*, 48-54.
33. Kakushima, M.; Hamel, P.; Frenette, R.; Rokach, J. Regioselective synthesis of acylpyrroles. *J. Org. Chem.* **1983**, *48*, 3214-3219.
34. Anderson, H.J.; Loader, C.E.; Xu, R.X.; Le, N.; Gogan, N.J.; McDonald, R.; Edwards, L.G. Pyrrole chemistry. XXVIII. Substitution reactions of 1-(phenylsulfonyl)pyrrole and some derivatives. *Can. J. Chem.* **1985**, *63*, 896-902.

*Sample Availability:* Samples of compounds **4a-e** are available from the authors.

© 2010 by the authors; licensee MDPI, Basel, Switzerland. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).