

Article

One-Pot Tandem Synthesis of 2-Arylquinazolines by a Multicomponent Cyclization Reaction

Leping Ye *, Lin Yu, Lijun Zhu and Xiaodong Xia

The Second Affiliated Hospital and Yuying Children's Hospital of Wenzhou Medical University, Wenzhou 325000, Zhejiang, China

* Author to whom correspondence should be addressed; E-Mail: yeeping@163.com; Tel.: +86-577-8800-2124; Fax: +86-577-8883-2693.

Received: 8 October 2013; in revised form: 21 October 2013 / Accepted: 21 October 2013 /

Published: 8 November 2013

Abstract: A series of 2-arylquinazolines have been synthesized in moderate to excellent yields by one-pot tandem reaction of (2-aminophenyl)methanols, aldehydes and ceric ammonium nitrate (CAN). The utility of this transformation was demonstrated by its compatibility with a wide range of functional groups. Thus, the method represents a simple and practical procedure to access 2-arylquinazolines.

Keywords: quinazolines; multicomponent; cyclization reaction; one-pot tandem synthesis


1. Introduction

Quinazolines and derivatives thereof are ubiquitous structural motifs that frequently occur in a wide variety of bioactive natural products such as prazosin [1], lapatinib [2], icotinib [3], *etc.* They possess a wide range of biological and pharmacological activities, including antibacterial [4], anti-inflammatory [5], antiplasmodial [6], antitumor [7], antimicrobial and antioxidant properties [8]. In addition, they have also been used as photochemotherapeutic agents [9], DNA-gyrase, JAK2, PDE5, and EGFR tyrosine kinase inhibitors [10] as well as CB2 receptor agonists [11]. Recently, Yang reported that substituted quinazolines possess novel potent and selective FLT3 inhibitory and anti-acute myeloid leukemia activities [12]. Consequently, in the past few years a number of synthetic methods to prepare these compounds have been described. Fu and co-workers reported that copper-catalyzed cascade reactions provide attractive and valuable routes for the construction of quinazoline derivatives via sequential Ullmann-type coupling and intramolecular cyclization [13]. Walton reported a microwave-promoted


syntheses of quinazolines by the reaction of 2-(aminoaryl)alkanone *O*-phenyl oximes with aldehydes [14]. Tandem reaction of 2-aminobenzophenones with benzylic amines followed by C–H functionalization has also been reported [15–18]. Zhang reported a three-component synthesis of quinazoline derivatives using a low melting sugar–urea–salt mixture as a solvent [19]. In 2012, Beifuss and co-workers developed a copper-catalyzed coupling reaction of *o*-bromobenzylbromides and benzamidines for the synthesis of quinazolines in an aqueous medium [20]. The condensation reactions of 2-aminobenzylamines with aldehydes followed by subsequent oxidation with strong oxidants (e.g., 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) [21], MnO₂ [22], and NaClO [23]) provide a conventional but simple method to synthesize quinazolines (Scheme 1(a)). However, these methods are hampered by the need to use a stoichiometric amount of a nonrenewable oxidant and the yields have not always been satisfactory. Recently, three improved methods have been reported for constructing quinazolines: (i) copper-catalyzed aerobic oxidative cyclization reaction of 2-(aminomethyl)benzenamines with aldehydes [24]; (ii) aerobic oxidative cyclization reaction of 2-(aminomethyl)benzenamines with aldehydes by a cooperative catalytic system of platinum/iridium alloyed nanoclusters and a dimeric catechol derivative [25]; (iii) iridium-catalyzed hydrogen transfer reaction of 2-(aminomethyl)benzenamines with aldehydes using styrene as a hydrogen acceptor [26].

Scheme 1. Reaction of (2-aminophenyl)methanols with aldehydes under different conditions.

Previous work



This work



Herein, we wish to report a new method for the synthesis of quinazolines by using a one-pot tandem reaction of (2-aminophenyl)methanols, aldehydes and ceric ammonium nitrate (CAN) (Scheme 1(b)).

2. Results and Discussion

We began our study by examining the reaction between (2-aminophenyl)methanol (**1a**) and benzaldehyde (**2a**) to obtain 2-phenyl-4*H*-benzo[*d*][1,3]oxazine (**4a**) and screen the optimal reaction conditions. Surprisingly, a trace amount of unexpected 2-phenylquinazoline (**3a**) was detected by GC/MS (EI) analysis during the process of performing the above condensation reaction in the presence of ceric ammonium nitrate (CAN). To the best of our knowledge, the synthesis of quinazolines using a one-pot tandem reaction of (2-aminophenyl)methanols, aldehydes and ceric ammonium nitrate has never been investigated before. In the present study, this unexpected reaction was thoroughly investigated. To realize the selective generation of **3a**, we initially ran a series of trial experiments in

the presence of palladium catalysts by adjusting the reaction parameters. However, little-to-no product **3a** was detected. To our delight, after the catalyst was switched to a copper catalyst such as CuOTf, the desired product **3a** was isolated in 19% yield when the combination of KF and CH₃CN was employed (Table 1, entry 1). Encouraged by this promising result, we further adjusted reaction parameters including copper catalysts, bases, and solvents.

Table 1. Optimization of the reaction conditions ^a.

Entry	Cu sources	Base	Solvent	Yield (%) ^b
1	CuOTf	KF	CH ₃ CN	19
2	CuOTf	Li ₂ CO ₃	CH ₃ CN	32
3	CuOTf	CsOAc	CH ₃ CN	17
4	CuOTf	K ₂ CO ₃	CH ₃ CN	45
5	CuOTf	K ₃ PO ₄	CH ₃ CN	14
6	CuOTf	<i>t</i> -BuOK	CH ₃ CN	33
7	CuOTf	LiOH	CH ₃ CN	42
8	CuOTf	NaOH	CH ₃ CN	48
9	CuOTf	KOH	CH ₃ CN	64
10	CuOTf	CsOH	CH ₃ CN	68
11	CuBr	CsOH	CH ₃ CN	82
12	CuCl	CsOH	CH ₃ CN	91
13	Cu(OTf) ₂	CsOH	CH ₃ CN	67
14	Cu(OAc) ₂	CsOH	CH ₃ CN	56
15	CuBr ₂	CsOH	CH ₃ CN	51
16	CuCl ₂	CsOH	CH ₃ CN	57
17	CuO	CsOH	CH ₃ CN	34
18	none	CsOH	CH ₃ CN	0
19	CuCl	CsOH	Toluene	31
20	CuCl	CsOH	THF	65
21	CuCl	CsOH	PhCl	36
22	CuCl	CsOH	DMF	22

^a Reaction conditions: **1a** (0.2 mmol), **2a** (0.3 mmol), CAN (0.3 mmol), Cu source (10 mol%), 2,2'-bipyridine (10 mol%), TEMPO (10 mol%), base (0.5 mmol), and solvent (2 mL), O₂, 30 °C, 24 h then 60 °C for 24 h.


^b Isolated yield of **3a**.

Screening revealed that the use of CsOH or KOH as a base achieved the best results. Other bases, including KF, Li₂CO₃, CsOAc, K₂CO₃, K₃PO₄, *t*-BuOK, LiOH, and NaOH, were less efficient (Table 1, entries 2–10). Among the copper sources used (e.g., CuOTf, CuBr, CuCl, Cu(OTf)₂, Cu(OAc)₂, CuBr₂, CuCl₂, and CuO), CuCl exhibited the highest catalytic reactivity (91% yield, Table 1, entries 10–17). No reaction occurred when the procedure was carried out in the absence of copper catalyst (Table 1, entry 18). Finally, we studied the solvent effect and found that CH₃CN was superior

to toluene, THF, PhCl, and DMF (Table 1, entries 12, 19–22). In addition, the reaction failed to give the desired product when the procedure was carried out under a N₂ atmosphere.







With the optimized reaction conditions in hand, we next explored the substrate scope of (2-aminophenyl)methanols **1** and aldehydes **2** for the one-pot tandem multicomponent cyclization reaction (Table 2). First, the reaction of (2-aminophenyl)methanol (**1a**) with various aldehydes **2a–k** in the presence of CAN was investigated under the standard conditions. The results disclosed that a variety of aldehydes are suitable substrates, and the cyclization reactions provide the corresponding products in moderate to good yields (Table 2, entries 1–11).

Table 2. Substrate scope of (2-aminophenyl)methanols and aldehydes ^a.



Entry	(2-Aminophenyl)methanols (1)	Aldehydes (2)	Product (3)	Yield (%) ^b
1	(1a)	(2a)	3a	91
2	(1a)	(2b)	3b	84
3	(1a)	(2c)	3c	66
4	(1a)	(2d)	3d	82
5	(1a)	(2e)	3e	93
6	(1a)	(2f)	3f	86
7	(1a)	(2g)	3g	84
8	(1a)	(2h)	3h	71
9	(1a)	(2i)	3i	78
10	(1a)	(2j)	3j	87
11	(1a)	(2k)	3k	89
12	(1b)	(2a)	3l	87

Table 2. Cont.

Entry	(2-Aminophenyl)methanols (1)	Aldehydes (2)	Product (3)	Yield (%) ^b
13	 (1c)		3m	81
14	 (1d)		3n	80
15	 (1e)		3o	67

^a Reaction conditions: **1** (0.2 mmol), **2** (0.3 mmol), CAN (0.3 mmol), CuCl (10 mol%), 2,2'-bipyridine (10 mol%), TEMPO (10 mol%), CsOH (0.5 mmol), and CH₃CN (2 mL), O₂, 30 °C, 24 h then 60 °C for 24 h.


^b Isolated yield.

First, the mono-substituent positions at the aryl moiety of aldehydes were evaluated, and the results demonstrated that steric effects of substituents had some effects on the reaction. For example, when the cyclization reaction of **1a** with *para*-, and *ortho*-methylbenzaldehyde was examined, 84% of **3b** was isolated, while the yield of **3c** decreased to 66% (Table 2, entries 2–3). The electronic properties of the substituents on the phenyl ring of the aldehydes also affected the yields of the reaction to some extent. Generally, the aldehydes bearing an electron-withdrawing substituent (e.g., –F, –Cl and –Br, Table 2, entries 5–7) afforded a slightly higher yield of cyclization products than those analogues bearing an electron-donating substituent (e.g., –Me and –OMe, Table 2, entries 2–4). However, aldehydes bearing a strong electron-withdrawing substituent (e.g., –CF₃) at the *para* position afforded a slightly lower yield of product **3h** (Table 2, entry 8). A moderate yield of 2-(naphthalen-2-yl)quinazoline (**3i**) was observed using 1-naphthaldehyde (**2i**) as substrate (Table 2, entry 9). In addition, heterocyclic aldehydes, such as 2-furylaldehyde (**2j**) and 2-thienylaldehyde (**2k**) could also be used as the substrates, leading to the corresponding desired products 2-(furan-2-yl)quinazoline (**3j**) and 2-(thiophen-2-yl)quinazoline (**3k**) in 87% and 89% yields, respectively (Table 2, entries 10–11).

Subsequently, several (2-aminophenyl)methanols **1b–e**, which bear different substituents at the aryl moiety, were evaluated (Table 2, entries 12–15). The results showed that different functional groups, including methyl, fluoro, chloro, and nitro, were well tolerated under the standard conditions. In general, the (2-aminophenyl)methanols bearing an electron-donating substituent produced a slightly higher yield of cyclization products than those analogues bearing an electron-withdrawing substituent. For example, substrates **1b** and **1e**, bearing a methyl or nitro group, reacted with substrate **2a** to give the corresponding products **3l** and **3o** in 87% and 67% yields, respectively (Table 2, entries 12 and 15). It is noteworthy that halogen-containing substrates **1c** and **1d** produced the desired products **3m** and **3n** in 80% and 81% yields, respectively (Table 2, entries 13–14).

Two plausible reaction pathways for the formation of 2-arylquinazolines have been established and are shown in Scheme 2. First, aerobic alcohol oxidation reaction of (2-aminophenyl)methanols **1** generates 2-aminobenzaldehydes **4**. An imine intermediate III is formed from intermediate I or intermediate II by two plausible reaction pathways, respectively. Then intramolecular cyclization of intermediate III affords dihydroquinazolines IV. Aromatization of intermediate IV gives 2-arylquinazolines **3** as the desired products. However, the mechanism in detail on the formation of the 2-arylquinazolines remains unclear in current stage.

Scheme 2. Plausible reaction pathways.



3. Experimental

3.1. General

Melting points are uncorrected. $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra were measured on a 500 MHz or 300 MHz spectrometer using $\text{DMSO-}d_6$ or CDCl_3 as the solvent with tetramethylsilane (TMS) as an internal standard at room temperature. Chemical shifts are given in δ relative to TMS, and the coupling constants J are given in hertz. High-resolution mass spectra were recorded on an ESI-Q-TOF mass spectrometer. Other commercially obtained reagents were used without further purification. All reactions under nitrogen atmosphere were conducted using standard Schlenk techniques. Column chromatography was performed using EM silica gel 60 (300–400 mesh).

3.2. General Procedure for the Synthesis of 2-Arylquinazolins

To a Schlenk tube were added (2-aminophenyl)methanols **1** (0.2 mmol), aldehydes **2** (0.3 mmol), CAN (0.3 mmol), CuCl (0.02 mmol), 2,2'-bipyridine (0.02 mmol), TEMPO (0.02 mmol), CsOH (0.5 mmol), and CH_3CN (2 mL). Next the tube was charged with O_2 (1 atm), and was stirred constantly at 30 °C for 24 h, then at 60 °C for 24 h. After the completion of the reaction, the reaction mixture was cooled to room temperature, diluted with ethyl acetate, and washed with brine. After the aqueous layer was extracted with ethyl acetate, the combined organic layers were dried over anhydrous MgSO_4 and evaporated under reduced pressure. The residue was purified by flash column chromatography (hexane/ethyl acetate) to afford the desired products **3**.

2-Phenylquinazoline (3a) [24]. Pale yellow solid (91% yield), mp 98–99 °C (Lit. 97–98 °C); $^1\text{H-NMR}$ ($\text{DMSO-}d_6$, 300 MHz): δ 9.70 (s, 1H), 8.55–8.59 (m, 2H), 8.17 (d, $J = 8.1$ Hz, 1H), 8.00–8.08 (m, 2H), 7.71–7.76 (m, 1H), 7.55–7.59 (m, 3H); $^{13}\text{C-NMR}$ ($\text{DMSO-}d_6$, 75 MHz): δ 161.3, 159.8, 149.8, 137.4, 134.8, 130.8, 128.7, 128.1, 127.9, 127.8, 123.3.

2-p-Tolylquinazoline (3b) [24]. Yellow solid (84% yield), mp 109–111 °C (Lit. 107–109 °C); $^1\text{H-NMR}$ (CDCl_3 , 500 MHz): δ 9.44 (s, 1H), 8.52 (d, $J = 8.0$ Hz, 2H), 8.07 (d, $J = 8.0$ Hz, 1H), 7.87–7.90 (m, 2H), 7.56–7.60 (m, 1H), 7.35 (d, $J = 8.0$ Hz, 2H), 2.45 (s, 3H); $^{13}\text{C-NMR}$ (CDCl_3 , 125 MHz): δ 161.1, 160.4, 150.8, 140.8, 135.3, 134.0, 129.4, 128.52, 128.50, 127.1, 127.0, 123.5, 21.5.

2-o-Tolylquinazoline (3c) [25]. Colorless oil (66% yield); $^1\text{H-NMR}$ (CDCl_3 , 500 MHz): δ 9.49 (s, 1H), 8.11 (d, $J = 8.5$ Hz, 1H), 7.89–7.95 (m, 2H), 7.77 (d, $J = 7.5$ Hz, 1H), 7.63 (t, $J = 7.5$ Hz, 1H), 7.44

(t, $J = 7.5$ Hz, 1H), 7.05–7.11 (m, 2H), 3.87 (s, 3 H); ^{13}C -NMR (CDCl_3 , 125 MHz): δ 162.4, 160.0, 157.7, 150.6, 134.0, 131.7, 130.8, 129.0, 128.5, 127.5, 127.0, 123.1, 120.7, 111.9, 56.0.

2-(4-Methoxyphenyl)quinazoline (3d) [24]. White solid (82% yield), mp 93–94 °C (Lit. 91–93 °C); ^1H -NMR (CDCl_3 , 500 MHz): δ 9.40 (s, 1H), 8.57–8.59 (m, 2H), 8.03 (d, $J = 8.9$ Hz, 1H), 7.86 (t, $J = 9.7$ Hz, 2H), 7.55 (t, $J = 8.0$ Hz, 1H), 7.04 (d, $J = 9.0$ Hz, 2H), 3.89 (s, 3H); ^{13}C -NMR (CDCl_3 , 125 MHz): δ 161.9, 160.9, 160.4, 150.9, 134.0, 130.9, 130.2, 128.4, 127.1, 126.8, 123.3, 114.0, 55.4.

2-(4-Fluorophenyl)quinazoline (3e) [24]. White solid (93% yield), mp 138–139 °C (Lit. 135–137 °C); ^1H -NMR (CDCl_3 , 500 MHz): δ 9.42 (s, 1H), 8.61–8.64 (m, 2H), 8.06 (d, $J = 8.5$ Hz, 1H), 7.89 (t, $J = 8.0$ Hz, 2H), 7.60 (t, $J = 7.0$ Hz, 1H), 7.20 (t, $J = 8.5$ Hz, 2H); ^{13}C -NMR (CDCl_3 , 125 MHz): δ 165.6, 163.7, 160.5, 160.1, 150.7, 134.1, 130.7, 130.6, 128.5, 127.2, 127.1, 123.5, 115.6, 115.4.

2-(4-Chlorophenyl)quinazoline (3f) [24]. White solid (86% yield), mp 135–137 °C (Lit. 133–135 °C); ^1H -NMR ($\text{DMSO}-d_6$, 500 MHz): δ 9.72 (s, 1H), 8.57 (d, $J = 8.5$ Hz, 2H), 8.19 (d, $J = 8.0$ Hz, 1H), 8.04–8.08 (m, 2H), 7.75–7.78 (m, 1H), 7.64 (d, $J = 8.5$ Hz, 2H); ^{13}C -NMR ($\text{DMSO}-d_6$, 125 MHz): δ 161.5, 158.8, 149.8, 136.3, 135.8, 135.1, 129.9, 128.9, 127.93, 127.90, 123.4.

2-(4-Bromophenyl)quinazoline (3g) [23]. White solid (84% yield), mp 121–123 °C (Lit. 120–121 °C); ^1H -NMR (CDCl_3 , 500 MHz): δ 9.43 (s, 1H), 8.50 (d, $J = 8.5$ Hz, 2H), 8.06 (d, $J = 9.0$ Hz, 1H), 7.89–7.92 (m, 2H), 7.60–7.66 (m, 3H); ^{13}C -NMR (CDCl_3 , 125 MHz): δ 160.5, 160.1, 150.7, 137.0, 134.2, 131.8, 130.2, 128.6, 127.4, 127.1, 125.4, 123.6.

2-(4-(Trifluoromethyl)phenyl)quinazoline (3h) [23]. White solid (71% yield), mp 144–146 °C (Lit. not reported); ^1H -NMR ($\text{DMSO}-d_6$, 500 MHz): δ 9.76 (s, 1H), 8.75 (d, $J = 8.5$ Hz, 2H), 8.21 (d, $J = 8.0$ Hz, 1H), 8.06–8.12 (m, 2H), 7.93 (d, $J = 8.5$ Hz, 2H), 7.78–7.81 (m, 1H); ^{13}C -NMR ($\text{DMSO}-d_6$, 125 MHz): δ 161.6, 158.4, 149.7, 141.2, 135.1, 130.7, 150.5, 128.7, 128.4, 128.0, 127.9, 125.8, 125.7, 125.3, 123.6, 123.1.

2-(Naphthalene-1-yl)quinazoline (3i) [25]. white solid (78% yield), mp 121–123 °C (Lit. 120–121 °C); ^1H -NMR (CDCl_3 , 500 MHz): δ 9.60 (s, 1H), 8.70 (d, $J = 8.0$ Hz, 1H), 8.17–8.19 (m, 2H), 7.93–8.03 (m, 4H), 7.70 (t, $J = 7.5$ Hz, 1H), 7.64 (t, $J = 7.5$ Hz, 1H), 7.52–7.58 (m, 2H); ^{13}C -NMR (CDCl_3 , 125 MHz): δ 163.5, 160.4, 150.6, 136.3, 134.3, 134.2, 131.2, 130.4, 129.6, 128.7, 128.5, 127.7, 127.1, 126.8, 125.89, 125.87, 125.3, 123.1.

2-(Furan-2-yl)quinazoline (3j) [26]. Brown solid (87% yield), mp 132–134 °C (Lit. 131–132 °C); ^1H -NMR (CDCl_3 , 500 MHz): δ 9.36 (s, 1H), 8.08 (d, $J = 9.0$ Hz, 1H), 7.87–7.90 (m, 2H), 7.68 (s, 1H), 7.60 (t, $J = 6.0$ Hz, 1H), 7.44–7.45 (m, 1H), 6.60–6.61 (m, 1H); ^{13}C -NMR (CDCl_3 , 125 MHz): δ 160.7, 154.1, 152.5, 150.4, 145.3, 134.5, 128.4, 127.2, 123.4, 114.1, 112.3.

2-(Thiophene-2-yl)quinazoline (3k) [13]. White solid (89% yield), mp 132–133 °C (Lit. 132–134 °C); ^1H -NMR (CDCl_3 , 500 MHz): δ 9.34 (s, 1H), 8.14–8.15 (m, 1H), 8.00 (d, $J = 9.0$ Hz, 1H), 7.85–7.88 (m, 2H), 7.51–7.57 (m, 2H), 7.18–7.20 (m, 1H); ^{13}C -NMR (CDCl_3 , 125 MHz): δ 160.5, 157.9, 150.6, 143.8, 134.3, 129.9, 129.2, 128.3, 128.2, 127.2, 127.0, 123.4.

6-Methyl-2-phenylquinazoline (3l) [27]. Pale yellow solid (87% yield), mp 131–132 °C (Lit. 130–132 °C); $^1\text{H-NMR}$ (CDCl_3 , 500 MHz): δ 9.38 (s, 1H), 8.60 (d, $J = 8.5$ Hz, 2H), 7.98 (d, $J = 8.5$ Hz, 1H), 7.72–7.74 (m, 1H), 7.67 (s, 1H), 7.50–7.55 (m, 3H), 2.58 (s, 3H); $^{13}\text{C-NMR}$ (CDCl_3 , 125 MHz): δ 160.4, 159.7, 149.4, 138.2, 137.4, 136.4, 130.4, 128.6, 128.4, 128.3, 125.8, 123.6, 21.6.

6-Fluoro-2-phenylquinazoline (3m) [14]. White solid (81% yield), mp 121–122 °C (Lit. 120–121 °C); $^1\text{H-NMR}$ (CDCl_3 , 500 MHz): δ 9.43 (s, 1H), 8.59–8.61 (m, 2H), 8.09–8.11 (m, 1H), 7.65–7.70 (m, 1H), 7.51–7.55 (m, 4H); $^{13}\text{C-NMR}$ (CDCl_3 , 125 MHz): δ 161.4, 160.7, 159.8, 159.7, 159.4, 148.0, 128.2, 137.7, 131.4, 131.3, 130.7, 128.7, 128.5, 124.6, 124.4, 124.0, 123.9, 110.2, 110.0.

6-Chloro-2-phenylquinazoline (3n) [17]. Pale yellow solid (80% yield), mp 158–159 °C (Lit. 157–159 °C); $^1\text{H-NMR}$ (CDCl_3 , 500 MHz): δ 9.38 (s, 1H), 8.59–8.61 (m, 2H), 8.02 (d, $J = 9.0$ Hz, 1H), 7.89 (d, $J = 2.5$ Hz, 1H), 7.81–7.83 (m, 1H), 7.51–7.55 (m, 3H); $^{13}\text{C-NMR}$ (CDCl_3 , 125 MHz): δ 161.3, 159.5, 149.2, 137.6, 135.0, 132.8, 130.9, 130.4, 128.7, 128.6, 125.8, 124.0.

7-Nitro-2-phenylquinazoline (3o). White solid (67% yield), mp 142–144 °C; IR (KBr) 2925, 1687, 1567, 1526, 1421, 1326, 1291, 927, 823, 740, 706, 686, 668 cm^{-1} ; $^1\text{H-NMR}$ (CDCl_3 , 500 MHz): δ 9.62 (s, 1H), 8.97 (d, $J = 2$ Hz, 1H), 8.64–8.66 (m, 2H), 8.357–8.361 (m, 1H), 8.11 (d, $J = 8.5$ Hz, 1H), 7.56–7.58 (m, 3 H); $^{13}\text{C-NMR}$ (CDCl_3 , 125 MHz): δ 162.9, 160.8, 151.2, 150.5, 136.9, 131.6, 129.0, 128.9, 128.8, 125.7, 124.9, 120.7. HRMS (ESI) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_{14}\text{H}_{10}\text{N}_3\text{O}_2$ 251.0695; Found: 251.0681.

4. Conclusions

In summary, we have successfully developed a new protocol for the copper-catalyzed cascade reaction of (2-aminophenyl)methanols, aldehydes, and ceric ammonium nitrate to afford a series of 2-arylquinazolines in moderate to excellent yields. Further efforts to extend this catalytic system to the preparation of other useful heterocyclic compounds are currently underway in our laboratories.

Acknowledgments

We thank the National Natural Science Foundation of China (No. 81102150), Traditional Chinese Medicine Scientific Research Foundation of Zhejiang Province (No. 2010ZA091) and Medicine and Health Science and Technology Projects of Zhejiang Province (No. 2012KYA133) for financial support.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Antonello, A.; Hrelia, P.; Leonardi, A.; Marucci, G.; Rosini, M.; Tarozzi, A.; Tumiatti, V.; Melchiorre, C. Design, synthesis, and biological evaluation of prazosin-related derivatives as multipotent compounds. *J. Med. Chem.* **2005**, *48*, 28–31.

2. Oude Munnink, T.H.; de Vries, E.G.E.; Vedelaar, S.R.; Timmer-Bosscha, H.; Schröder, C.P.; Brouwers, A.H.; Lub-de Hooge, M.N. Lapatinib and 17AAG reduce ^{89}Zr -trastuzumab-F(ab')₂ uptake in SKBR3 tumor xenografts. *Mol. Pharm.* **2012**, *9*, 2995–3002.
3. Hu, S.-J.; Long, W.; Wang, F.; Li, Z.-Q. Methods of preparing icotinib and icotinib hydrochloride, and intermediates thereof. WO Patent 2013064128, 2013.
4. Tiwari, R.; Chhabra, G. Evaluation of antibacterial activity of novel quinazoline derivatives. *Asian J. Chem.* **2010**, *22*, 5981–5986.
5. Balakumar, C.; Lamba, P.; Kishore, D.P.; Narayana, B.L.; Rao, K.V.; Rajwinder, K.; Rao, A.R.; Shireesha, B.; Narsaiah, B. Synthesis, anti-inflammatory evaluation and docking studies of some new fluorinated fused quinazolines. *Eur. J. Med. Chem.* **2010**, *45*, 4904–4913.
6. Kabri, Y.; Azas, N.; Dumetre, A.; Hutter, S.; Laget, M.; Verhaeghe, P.; Gellis, A.; Vanelle, P. Original quinazoline derivatives displaying antiplasmodial properties. *Eur. J. Med. Chem.* **2010**, *45*, 616–622.
7. Noolvi, M.N.; Patel, H.M.; Bhardwaj, V.; Chauhan, A. Synthesis and *in vitro* antitumor activity of substituted quinazoline and quinoxaline derivatives: Search for anticancer agent. *Eur. J. Med. Chem.* **2011**, *46*, 2327–2346.
8. Kumar, A.; Sharma, P.; Kumari, P.; Lal Kalal, B. Exploration of antimicrobial and antioxidant potential of newly synthesized 2,3-disubstituted quinazoline-4(3H)-ones. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 4353–4357.
9. Barraja, P.; Caracausi, L.; Diana, P.; Montalbano, A.; Carbone, A.; Salvador, A.; Brun, P.; Castagliuolo, I.; Tisi, S.; Dall'Acqua, F.; *et al.* Pyrrolo[3,2-h]quinazolines as photochemotherapeutic agents. *ChemMedChem* **2011**, *6*, 1238–1248.
10. Yang, S.H.; Khadka, D.B.; Cho, S.H.; Ju, H.-K.; Lee, K.Y.; Han, H.J.; Lee, K.-T.; Cho, W.-J. Virtual screening and synthesis of quinazolines as novel JAK2 inhibitors. *Bioorg. Med. Chem.* **2011**, *19*, 968–977.
11. Saari, R.; Toermae, J.-C.; Nevalainen, T. Microwave-assisted synthesis of quinoline, isoquinoline, quinoxaline and quinazoline derivatives as CB2 receptor agonists. *Bioorg. Med. Chem.* **2011**, *19*, 939–950.
12. Li, W.-W.; Wang, X.-Y.; Zheng, R.-L.; Yan, H.-X.; Cao, Z.-X.; Zhong, L.; Wang, Z.-R.; Ji, P.; Yang, L.-L.; Wang, L.-J.; *et al.* Discovery of the novel potent and selective FLT3 inhibitor 1-{5-[7-(3-morpholinopropoxy)quinazolin-4-ylthio]-[1,3,4]thiadiazol-2-yl}-3-*p*-tolylurea and its anti-acute myeloid leukemia (AML) activities *in Vitro* and *in Vivo*. *J. Med. Chem.* **2012**, *55*, 3852–3866.
13. Wang, C.; Li, S.; Liu, H.; Jiang, Y.; Fu, H. Copper-catalyzed synthesis of quinazoline derivatives via ullmann-type coupling and aerobic oxidation. *J. Org. Chem.* **2010**, *75*, 7936–7938.
14. Portela-Cubillo, F.; Scott, J.S.; Walton, J.C. Microwave-promoted syntheses of quinazolines and dihydroquinazolines from 2-aminoarylalkanone *O*-phenyl oximes. *J. Org. Chem.* **2009**, *74*, 4934–4942.
15. Han, B.; Wang, C.; Han, R.F.; Yu, W.; Duan, X.Y.; Fang, R.; Yang, X.L. Efficient aerobic oxidative synthesis of 2-aryl quinazolines via benzyl C–H bond amination catalyzed by 4-hydroxy-TEMPO. *Chem. Commun.* **2011**, *47*, 7818–7820.

16. Karnakar, K.; Kumar, A.V.; Murthy, S.N.; Ramesh, K.; Nageswar, Y.V.D. Recyclable graphite oxide promoted efficient synthesis of 2-phenyl quinazoline derivatives in the presence of TBHP as an oxidant. *Tetrahedron Lett.* **2012**, *53*, 4613–4617.
17. Zhang, J.T.; Zhu, D.P.; Yu, C.M.; Wan, C.F.; Wang, Z.Y. A simple and efficient approach to the synthesis of 2-phenylquinazolines via sp^3 C–H functionalization. *Org. Lett.* **2010**, *12*, 2841–2843.
18. Zhang, J.T.; Yu, C.M.; Wang, S.J.; Wan, C.F.; Wang, Z.Y. A novel and efficient methodology for the construction of quinazolines based on supported copper oxide nanoparticles. *Chem. Commun.* **2010**, *46*, 5244–5246.
19. Zhang, Z.-H.; Zhang, X.-N.; Mo, L.-P.; Li, Y.-X.; Ma, F.-P. Catalyst-free synthesis of quinazoline derivatives using low melting sugar–urea–salt mixture as a solvent. *Green Chem.* **2012**, *14*, 1502–1506.
20. Malakar, C.C.; Baskakova, A.; Conrad, J.; Beifuss, U. Copper-catalyzed synthesis of quinazolines in water starting from *o*-bromobenzylbromides and benzamidines. *Chem. Eur. J.* **2012**, *18*, 8882–8885.
21. Vanden Eynde, J.J.; Godin, J.; Mayence, A.; Maquestiau, A.; Anders, E. A new and convenient method for the preparation of 2-substituted quinazolines. *Synthesis* **1993**, *9*, 867–869.
22. Peng, Y.; Zeng, Y.; Qiu, G. Cai, L.; Pike, V.W. A convenient one-pot procedure for the synthesis of 2-arylquinazolines using active MnO_2 as oxidant. *J. Heterocycl. Chem.* **2010**, *47*, 1240–1245.
23. Maheswari, C.U.; Kumar, G.S.; Venkateshwar, M.; Kumar, R.A.; Kantam, M.L.; Reddy, K.R. Highly efficient one-pot synthesis of 2-substituted quinazolines and 4*H*-benzo[*d*][1,3]oxazines via cross dehydrogenative coupling using sodium hypochlorite. *Adv. Synth. Catal.* **2010**, *352*, 341–346.
24. Han, B.; Yang, X.-L.; Wang, C.; Bai, Y.-W. Pan, T.-C.; Chen, X.; Yu, W. CuCl/DABCO/4-HO-TEMPO-catalyzed aerobic oxidative synthesis of 2-substituted quinazolines and 4*H*-3,1-benzoxazines. *J. Org. Chem.* **2012**, *77*, 1136–1142.
25. Yuan, H.; Yoo, W.-J.; Miyamura, H.; Kobayashi, S. A cooperative catalytic system of platinum/iridium alloyed nanoclusters and a dimeric catechol derivative: An efficient synthesis of quinazolines through a sequential aerobic oxidative process. *Adv. Synth. Catal.* **2012**, *354*, 2899–2904.
26. Fang, J.; Zhou, J.; Fang, Z. Synthesis of 2-substituted quinazolines via iridium catalysis. *RSC Adv.* **2013**, *3*, 334–336.
27. Rossi, E.; Stradi, R. A new synthesis of quinazoline ring. *Synthesis* **1989**, *1989*, 214–215.

Sample Availability: Contact the authors.

© 2013 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).