Supplementary Materials: Two New Oleanane-Type Saponins with Anti-Proliferative Activity from *Camellia oleifera* Abel. Seed Cake

Jian-Fa Zong, Yun-Ru Peng, Guan-Hu Bao, Ru-Yan Hou * and Xiao-Chun Wan *

Contents

Contents	S1
Figure S1. Positive ion mode HR-ESI-MS spectrum of compound 1	S2
Figure S2. Positive ion mode HR-ESI-MS spectrum of compound 2	S2
Figure S3. ¹ H-NMR spectrum of compound 1	S3
Figure S4. ¹³ C-NMR spectrum of compound 1	S3
Figure S5. ¹ H- ¹ H COSY spectrum of compound 1	S4
Figure S6. HSQC spectrum of compound 1	S4
Figure S7. HMBC spectrum of compound 1	S5
Figure S8. NOESY spectrum of compound 1	S5
Figure S9. ¹ H-NMR spectrum of compound 2	S6
Figure S10. ¹³ C-NMR spectrum of compound 2	S6
Figure S11. ¹ H- ¹ H COSY spectrum of compound 2	S7
Figure S12. HSQC spectrum of compound 2	S7
Figure S13. HMBC spectrum of compound 2	



Figure S1. Positive ion mode HR-ESI-MS spectrum of compound 1.



Figure S2. Positive ion mode HR-ESI-MS spectrum of compound 2.



Figure S3. ¹H-NMR spectrum of compound 1.



Figure S4. ¹³C-NMR spectrum of compound 1.





Figure S5. ¹H-¹H COSY spectrum of compound 1.



Figure S6. HSQC spectrum of compound 1.







Figure S8. NOESY spectrum of compound 1.

Molecules 2016, 21, 188; doi:10.3390/molecules21020188



Figure S9. ¹H-NMR spectrum of compound 2.



Figure S10. ¹³C-NMR spectrum of compound 2.



Figure S12. HSQC spectrum of compound 2.



Figure S13. HMBC spectrum of compound 2.