

Supplementary material: ^1H and ^{13}C NMR spectra of synthesized compounds

For manuscript entitled:



Deep eutectic solvents as convenient media for synthesis of novel coumarinyl Schiff bases and their QSAR studies

Maja Molnar^a, Mario Komar^a, Harshad Brahmbhatt^a, Jurislav Babić^a, Stela Jokić^a, Vesna Rastija^{b,*}



^aJosip Juraj Strossmayer University of Osijek, Faculty of Food Technology Osijek, Franje Kuhaca 20, 31000 Osijek, Croatia

^bJosip Juraj Strossmayer University of Osijek, Faculty of Agriculture in Osijek, Vladimira Preloga 1, 31000 Osijek, Croatia


*(E)-N'-(4-methoxybenzylidene)-2-((4-methyl-2-oxo-2H-chromen-7-yl)oxy)acetohydrazide (**10**) [2]*





(E)-N'-(4-(dimethylamino)benzylidene)-2-((4-methyl-2-oxo-2H-chromen-7-yl)oxy)acetohydrazide (**29**) [5]




(E)-N'-(3-(4-chlorophenyl)-1-(4-fluorophenyl)-1H-pyrazol-4-yl)methylene)-2-((4-methyl-2-oxo-2H-chromen-7-yl)oxy)acetohydrazide (30)





(E)-N'-(3-(4-bromophenyl)-1-(*p*-tolyl)-1*H*-pyrazol-4-yl)methylene)-2-((4-methyl-2-oxo-2*H*-chromen-7-yl)oxy)acetohydrazide (**31**)





*(E)-N'-(3-(4-iodophenyl)-1-(*p*-tolyl)-1*H*-pyrazol-4-yl)methylene)-2-((4-methyl-2-oxo-2*H*-chromen-7-yl)oxy)acetohydrazide (32)*





(E)-N'-(3-(4-methoxyphenyl)-1-(*p*-tolyl)-1*H*-pyrazol-4-yl)methylene)-2-((4-methyl-2-oxo-2*H*-chromen-7-yl)oxy)acetohydrazide (**33**)



*(E)-N'-(3-(4-chlorophenyl)-1-(*p*-tolyl)-1*H*-pyrazol-4-yl)methylene-2-((4-methyl-2-oxo-2*H*-chromen-7-yl)oxy)acetohydrazide (34)*



(E)-N'-(1-(4-chlorophenyl)-3-(4-nitrophenyl)-1H-pyrazol-4-yl)methylene)-2-((4-methyl-2-oxo-2H-chromen-7-yl)oxy)acetohydrazide (35)



(E)-N'-((4-chloro-2-oxo-2H-chromen-3-yl)methylene)-2-((4-methyl-2-oxo-2H-chromen-7-yl)oxy)acetohydrazide (36)

