

Review

Recent Synthesis Developments of Organoboron Compounds via Metal-Free Catalytic Borylation of Alkynes and Alkenes

Yanmei Wen, Chunmei Deng, Jianying Xie and Xinhuang Kang *

Faculty of Chemistry and Environmental Science, Guangdong Ocean University, Zhanjiang 524088, China; wenym@gdou.edu.cn (Y.W.); dcm2382405@163.com (C.D.); xiejy1961@126.com (J.X.)

* Correspondence: kang.xinhuang@aliyun.com; Tel.: +86-759-2383728

Academic Editor: Igor V. Alabugin

Received: 28 October 2018; Accepted: 19 December 2018; Published: 28 December 2018



Abstract: Diboron reagents have been traditionally regarded as “Lewis acids”, which can react with simple Lewis base to create a significant nucleophilic character in one of boryl moieties. In particular, bis(pinacolato)diboron (B_2pin_2) reacts with simple Lewis bases, such as *N*-heterocyclic carbenes (NHCs), phosphines and alkoxides. This review focuses on the application of trivalent nucleophilic boryl synthon in the selective preparation of organoboron compounds, mainly through metal-free catalytic diboration and the β -boration reactions of alkynes and alkenes.

Keywords: organoboron compound; metal-free catalytic boration; nucleophilic boron; alkyne; alkene


1. Introduction

Organoboron compounds play an important role in organic chemistry as they can be converted into a wide variety of functional groups [1–13]. Thus, the synthesis of different types of organoboron compounds continues to be an important research field [14–20]. Conventional methods for their preparation are based on either reactive organometallic reagents or transition-metal-mediated processes [21–37]. Recently, great effort has been made towards the development of transition-metal-free methods with high chemoselectivity or regioselectivity and environmental sustainability [38–48]. The transition-metal-free catalyzed borylation reaction of alkynes and alkenes offers an attractive method for generating valuable organoboron compounds. The interaction of a diboron reagent, in particular bis(pinacolato)diboron (B_2pin_2), with a simple Lewis base, such as *N*-heterocyclic carbenes (NHCs), phosphines and alkoxides, generates Lewis acid-base adducts, in which the formally intact sp^2 boryl moiety becomes the important source of nucleophilic boryl species, resulting in an appropriate approach towards transition-metal-free selective preparation of organoboron compounds. This review focuses on the application of trivalent nucleophilic boryl synthon in the selective preparation of organoboron compounds, mainly through diboration and β -boration reactions of alkynes and alkenes.


2. Organoboron Compounds via Transition-Metal-Free Catalytic Diboration of Unsaturated Hydrocarbons

In recent years, the preparation of 1,2-diborated or 1,1-diborated products has rapidly grown due to the addition of diboryl compounds to unsaturated organic compounds in the transition metal-free catalysis. The first diboration reaction of non-activated alkenes in the absence of transition metal complexes was reported by Fernández and coworkers (Scheme 1a) [49]. The diboration of non-activated olefins, such as terminal and internal alkenes, allenes and vinylarenes, can be produced via the formation of a 1,2-diborated product in a tetrahydrofuran (THF) solvent at 70 °C for ~6–16 h.

The interaction of B_2pin_2 (**1**) and a methoxide anion, derived from the combination of Cs_2CO_3 and MeOH, can generate in situ the adduct $[B_2pin_2OMe]^-$. This adduct reacts with the unsubstituted carbon atom (C^1) of the $C=C$ double bond via the transition state (TS1), which is the overlap between the strongly polarized B-B σ orbital of the activated diboron reagent and the $C-C$ π^* orbital of the olefin. When the B-B bond weakens, the negative charge density on the C^2 of the $C=C$ double bond increases. The negatively charged olefin-boryl olefin-B(pin) fragment attacks the boron atom with an electrophilic character to form the second transition state (TS2), which then protonates to form the diborated product (Scheme 2). The adduct $[B_2pin_2OMe]^-$ has been fully characterized by both ^{11}B -NMR spectroscopy and ESI-MS. A later report extended the methodology using unsymmetrical Bpin-Bdan (dan represents 1,8-diaminonaphthalene) (**2**) as the diboron reagents, where the methoxide selectively interacted with the stronger Lewis acid Bpin moiety. The Bdan moiety appears in the internal position (Scheme 1b) [50].




Scheme 1. (a) Transition-metal-free diboration reaction of non-activated olefins; (b) Mixed organocatalytic diboration of non-activated olefins.




Scheme 2. Suggested catalytic cycle for the diboration of non-activated olefins.

In 2015, Sawamura reported the formation of α,β -diboryl acrylates through trialkylphosphine catalyzed anti-selective diboration of the $C\equiv C$ triple bond (Scheme 3a) [51]. Through cross-coupling reaction of the differentiated heteroatom substituents of the α,β -diboryl acrylates in a stepwise manner, a diverse array of unsymmetrical tetrasubstituted alkenes can be accessed. Very recently, Santos employed an unsymmetrical Bpin–Bdan **2** as a diboron reagent for the direct diboration of alkynamides, which display broad functional group tolerance. The transition-metal-free catalyzed diboration of alkynamides with Bpin–Bdan, where the Bdan and Bpin moiety install α - and β -carbon atoms respectively. (Scheme 3b) [52].



Scheme 3. (a) Phosphine-Catalyzed 1,2-diboration of alkynoate; (b) 1,2-diboration of N-substituted phenylpropiolamides.

In 2014, Uchiyama and coworkers reported that the diboration of propargylic alcohols with diboron reagents in the presence of a stoichiometric amount of *n*-BuLi enables the trans-diboration of alkynes through aqueous workup. The present trans-diborylation gave oxaborole moieties which are key structural platforms and potent pharmacophores in material and pharmaceutical sciences (Scheme 4) [53]. Very recently, the direct diboration of alkynes using a more reactive unsymmetrical pinB-Bmes₂ (**3**) as the diboron reagent was reported by Yamashita's group. The mixture of two cis-isomers and one trans-isomer was obtained when the base-catalyzed diboration reaction of the terminal alkynes was conducted in the system consisting of *n*-BuLi and 1,2-dimethoxyethane in toluene (Scheme 5) [54].




Scheme 4. Trans-selective diborylation reaction of propargylic alcohols.


In 2015, Sawamura revealed a new method for the synthesis of 1,1-diborylalkenes through a Brønsted base catalyzed reaction between bis(pinacolato)diboron and various terminal alkynes containing propiolates, propiolamides and 2-ethynylazoles (Scheme 6a) [55]. In 2018, the Brønsted base catalyzed diboration of strong electron deficient alkynes, including various propiolates, ynones and propiolamides, was developed by Song's group (Scheme 6b) [56]. The terminal and internal alkynes were allowed to react under the "optimal" conditions, and good to excellent yields of the corresponding gem-diborylalkanes were obtained.

The asymmetric organocatalytic 1,2-diboration of alkenes has been accomplished utilizing economically accessible chiral alcohols as additives to form the Lewis acid-base $^*RO^- \rightarrow$ bis(pinacolato)diboron adduct that is added to cyclic and non-cyclic alkenes, providing moderate enantioselectivity. The authors reported moderate conversions and moderate levels of enantioselectivity for the synthesis of the 1,2-diborated product (Scheme 7a) [57]. Alternatively, the enantioselective diboration of alkenes can be achieved using inexpensive chiral carbohydrates, such


as the pseudoenantiomeric glycol 6-tertbutyldimethylsilyl-1,2-dihydroglucal (**4**) and dihydrorhamnal (**5**) (Scheme 7b) [58]. The alkoxide-catalyzed directed diboration of alkenyl alcohols through the intramolecular activation of the diboron reagent and the selective delivery of the nucleophilic boryl unit offers access to the diboration of cyclic and acyclic homoallylic and bishomoallylic alcohol substrates (Scheme 7c) [59].



Scheme 5. Reaction of pinB-BMes₂ with terminal alkynes.



Scheme 6. (a) Synthesis of 1,1-diborylalkenes, (b) Synthesis of functionalized geminal-diborylalkanes.




Scheme 7. Enantioselective diboration of olefins. (a) synthesis of the 1,2-diborated product; (b) synthesis of pseudoenantiomeric glycol 6-tertbutyldimethylsilyl-1,2-dihydroglucal (**4**) and dihydrorhamnal (**5**); (c) synthesis of diboration of cyclic and acyclic homoallylic and bishomoallylic alcohol substrates.


3. Organoboron Compounds via Transition-Metal-Free Catalytic β -boration of α,β -Unsaturated Compounds

3.1. *N*-Heterocyclic Carbene Catalysis


In 2009, Hoveyda and coworkers firstly reported that an imidazolium salt **6** in the absence of a transition metal was able to promote the boron conjugate addition of cyclic and acyclic α,β -unsaturated ketones and esters, providing β -boryl carbonyls (Scheme 8) [60]. It is noteworthy that in the presence of a base (NaOtBu) only, the borylation reaction did not take place. It was proposed that in this process a nucleophilic *N*-heterocyclic carbene (NHC) could associate with the diboron reagent in presence of a base and an in situ generated NHC·B₂(pin)₂ complex. The diboron reagent becomes nucleophilic and is able to transfer the “intact” sp² boryl group to the activated olefins in boron conjugate additions. A proton from alcohol additives or aqueous workup served as the electrophilic counterpart of the boron nucleophile (Scheme 9). Density functional theory (DFT) calculations and ¹¹B-NMR studies were carried out to determine the exact structure of the proposed sp²–sp³ hybridized NHC·B₂(pin)₂ adduct. The DFT calculations showed polarization of the B–B bond and diminished electrophilic character of the two boron atoms, where the electron density was strongly polarized towards the unbound boron center. However, due to the weak association of the NHC to the sp²–sp² diboron, causing dynamic exchange, the ¹¹B-NMR signal for B₂pin₂ disappeared within five minutes after treatment with the NHC. In 2012, the crystal structure of the neutral NHC adduct **7** was reported by Marder and coworkers (Scheme 10) [61].



Scheme 8. The β -borylation of α,β -unsaturated carbonyl compounds.




Scheme 9. Mechanism for B₂pin₂ activation and conjugate addition to an enone.



Scheme 10. Formation of the sp²–sp³ hybridized NHC·B₂(pin)₂ compound.

In a subsequent study, Hoveyda reported further work on enantioselective method for boron conjugate addition to α,β -unsaturated carbonyls with a chiral NHC **8** in combination with 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). This method can tolerate a broad range of substrates, including acyclic and cyclic α,β -unsaturated ketones, as well as acyclic esters, amides, and aldehydes.

The desired β -boryl carbonyls are obtained in up to a 94% isolated yield and an >98:2 enantiomer ratio (Scheme 11a) [62]. A later report extended this methodology to β -disubstituted enones, generating products containing boron-substituted quaternary carbon with enantioselectivity (Scheme 11b) [63]. The use of diboranes and *N*-heterocyclic carbenes in the absence of a transition metal complex has been reported by other groups [64,65].






Scheme 11. (a) NHC-catalyzed enantioselective boryl conjugate addition to unsaturated carbonyls, (b) NHC-catalyzed enantioselective boryl conjugate addition to enones.

3.2. Base Catalysis


In 2012, Fernández reported the utility of the base-catalyzed method for the β -boration of acyclic and cyclic activated olefins in cases involving α,β -unsaturated amides and phosphonates [66]. The methodology could be applied to different diboron reagents, such as bis(pinacolato)diboron (B_2pin_2), bis(catecholato)borane (B_2cat_2) and so on. Alternative inorganic and organic bases were explored, such as NaOMe, LiOMe, NaOtBu, K_2CO_3 , CsF and Verkade's base, the latter being the most efficient (Scheme 12). With an excess of MeOH, Verkade's base is completely protonated and the generated MeO^- forms a Lewis acid-base adduct with B_2pin_2 . Computational studies identified that there was an overlap between the strongly polarized B–B σ orbital and the C–C π^* orbital, and DFT calculations revealed that in adducts the sp^2 boron atom gains a strong nucleophilic character because the sp^3 boron atom loses negative charge density upon the charge transfer from the Lewis base. The intact sp^2 boron atom corresponds to a nucleophilic attack at the β -carbon atom of the activated olefins, forming a transition state (TS), which releases the "(pin)BOMe" and leads to the formation of an anionic intermediate, which is then protonated to provide the β -boration product (Scheme 13).

A later report extended the methodology to non-symmetrical diboron reagents, such as Bpin–Bdan **2**, where the RO^- selectively interacted with the stronger Lewis acid Bpin moiety (Scheme 14a) [67]. Experimental and theoretical investigation confirmed that the interaction of RO^- and the Bpin moiety preferred to form a $[RO \rightarrow Bpin-Bdan]^-$ intermediate, which then resulted in exclusively the β -carbon Bdan carbonyl compound with high yields. The adduct $[MeO \rightarrow Bpin-Bpin]^-$ which efficiently mediates the β -boration of α,β -unsaturated imines formed in situ was reported by Fernández's group (Scheme 14b) [68]. From a theoretical point of view, the mechanism of β -boration of α,β -unsaturated imines has been thought to involve quaternization of the diboron reagent with methoxide. The function of the phosphine has been regarded to the ion pair formation.


Scheme 12. Verkade's base mediates β -boration of ethyl crotonate.Scheme 13. Proposed reaction pathway for β -boration of methyl acrylate.Scheme 14. (a) The β -boration of α,β -unsaturated compounds, (b) The β -boration of in situ formed α,β -unsaturated imines.

3.3. Phosphine Catalysis

In 2010, Fernández reported the β -borylation of α,β -unsaturated compounds using a trivalent phosphorus nucleophile in combination with catalytic amounts of bases and stoichiometric amounts of methanol (Scheme 15) [69]. Further screening showed that a variety of phosphorus compounds including achiral, chiral mono and bidentate phosphines could also give reasonable conversions without a transition metal. The process was proposed that the phosphine interacted with the empty orbital of one of the boron atoms to result in the heterolytic cleavage of the B–B bond of B_2pin_2 . Upon such interaction, the other boron moiety could act as a nucleophile towards the α,β -unsaturated esters and ketones (Scheme 16) [69].




Scheme 15. Phosphine-mediated asymmetric β -boration of α,β -unsaturated compounds.




Scheme 16. Plausible mechanism for the phosphine-catalyzed β -boration of α,β -unsaturated carbonyl compound.

In a further publication, Fernández presented novel reaction conditions for the β -boration reaction of α,β -unsaturated carbonyl compounds without a transition metal. Eluding the use of Brønsted bases to promote the catalytic active species, the reaction can be carried out by the unique presence of catalytic amounts of phosphine in MeOH (Scheme 17) [70]. The experimental and theoretical studies demonstrated interactions between the phosphine and the substrate. Initially, the phosphine interacts with the substrate to form a strongly basic zwitterionic phosphonium enolate species, which would aid deprotonation of the MeOH to eventually form the ion pair $[\alpha\text{-(H),}\beta\text{-(PR}_3\text{)-ketone}]^+ [B_2pin_2 \cdot MeO]^-$ (Scheme 18).




Scheme 17. Phosphine assisted β -boration reaction of α,β -unsaturated carbonyl compounds in MeOH.



Scheme 18. Ion pair formation.


Córdova developed the first metal-free one-pot three-component catalytic selective reaction between a diboron reagent, α,β -unsaturated aldehydes, and 2-(triphenylphosphoranylidene) acetate esters for the synthesis of homoallylboronates. The multicomponent reaction undergoes a catalytic β -boration of α,β -unsaturated aldehydes to give the β -borylaldehyde. A subsequent Wittig reaction between the β -borylaldehyde and the 2-(triphenylphosphoranylidene)acetate esters yields the corresponding homoallylboronates with high chemoselectivity and regioselectivity (Scheme 19) [71].




Scheme 19. One-pot three-component synthesis of homoallylboronates.

4. Organoboron Compounds via Transition-Metal-Free Catalytic Borylation of Allylic and Propargylic Alcohols

The base-catalyzed borylation of tertiary allylic alcohols can be efficiently performed to prepare 1,1-disubstituted allyl boronates with a high yields (Scheme 20a) [72]. The reactions are performed in the presence of a catalytic amount of Cs_2CO_3 and MeOH to promote the formation of the Lewis acid-base adduct, $[\text{Hbase}]^+[\text{MeOB}_2\text{pin}_2]^-$, which may also be responsible for the tandem allylic borylation/diboration of allylic alcohols that generate 1,2,3-polyborated products (Scheme 20a). To complement the study, the scope of the metal-free allylic borylation was further expanded to 1-propargylic cyclohexanol, to form the corresponding product with a working hydroborated triple bond at 50 °C. Interestingly, by increasing the temperature to 90 °C, the alkenyl borane was selectively formed (Scheme 20b). The author suggested a possible mechanism to account for this process. First, the adduct $[\text{B}_2\text{pin}_2\cdot\text{MeO}]^-$ was formed through a base-mediated nucleophilic attack at one of the Bpin moieties of B_2pin_2 . Second, the other Bpin unit with nucleophilic activity attacks the terminal carbon atom of the allylic alcohol, which leads cleavage of the C–O bond. Last, the base catalyst is regenerated via the reaction of the conjugated acid of the base (BaseH^+) and the OH group (Scheme 21).




Scheme 20. (a) Borylation of tertiary allylic alcohols, (b) Borylation of propargylic alcohols.




Scheme 21. Suggested mechanism for the metal-free allylic borylation.

Another facile synthetic process for converting tertiary allylic alcohols to corresponding allylic boronates in moderate to good yields was reported. The mechanistic approach suggested that allylic boronate formation was achieved via the sequential processes of Fernández-type alkoxide-mediated diborylation of an alkene and the boron-Wittig reaction (Scheme 22) [73].



Scheme 22. Direct conversion of allylic alcohols to allylic boronates.

On the basis of “pseudo-intramolecular activation” [53] of diborons enabling the trans-selective diboration of alkynes, Uchiyama and coworkers found that the carboboration reaction was enabled by pseudo-intramolecular activation of alkynylboronates using propargylic alcohols to produce oxaborole products (Scheme 23) [74]. The synthetic utility of this product was investigated using medicinal chemistry and strong blue fluorescence emission.




Scheme 23. Trans-selective alkynylboration reaction of alkynes.

5. Organoboron Compounds via Transition-Metal-Free Catalytic Hydroboration of Unsaturated Hydrocarbons


In 2016, Song reported that the transition-metal-free catalytic borylation of arylacetylenes or vinyl arenes can be efficiently performed to produce alkylboronates through tandem borylation

and protodeboronation (Scheme 24a) [75]. This method has high regioselectivity and chemoselectivity, good to excellent yields and can tolerate a broad range functional groups. The transition-metal-free method for borylation of alkynes and alkenes with bis(pinacolato)diboron can also easily afford vinylboronates (Scheme 24b) [76] or diboryl alkanes (Scheme 24c) [77]. In 2018, Fernández and coworkers reported that the 1,4-hydroboration of 1,3-dienes was followed by an in situ diboration reaction of the internal double bond to produce a 1,2,3-triboration product (Scheme 24d) [78].



Scheme 24. (a) Synthesis of alkylboronates from arylacetylenes and vinyl arenes, (b) Synthesis of β -vinylboronates from terminal alkynes, (c) Synthesis of 1,2-diborylalkanes from non-activated olefins, (d) Synthesis of 1,2,3-triborated compounds from 1,3-dienes.

A new approach to the transition-metal-free synthesis of alkenyl boronates was based on the hydroboration of the distal double bond of allenamides (Scheme 25) [79]. In order to obtain exclusively the complete stereoselective product, the acyl groups on the amine moiety were crucial, which formed a stable allylic anion intermediate that was further regioselectively protonated to obtain a *Z*-isomer.



Scheme 25. Transition-metal-free hydroboration of allenamides.

6. Conclusions

In this review, we have presented numerous very useful processes for the synthesis of valuable organoboron compounds, which involve transition-metal-free catalyzed diboration, β -boration and boryl substitution reactions. The key to the success of this reaction is to use a simple Lewis base capable of activating the diboron reagent to create a significant nucleophilic boryl synthon. The reactions tolerate a wide variety of functional groups and progress under metal-free reaction conditions, thus avoiding contaminating the final boron products with heavy metals, a problem which is frequently encountered in the manufacturing process. Although the metal-free catalyzed borylation of alkynes

and alkenes is relatively unexplored at this time, there is no doubt that the future is promising for this reaction and there will be many exciting discoveries in this field.

Author Contributions: Conceptualization: Y.W., C.D., J.X., X.K.; Writing-Original Draft Preparation, Y.W., C.D., J.X., X.K.; Writing-Review & Editing, Y.W., C.D., J.X., X.K.; Funding Acquisition, X.K.

Acknowledgments: The authors of this paper appreciate the financial support from the Guangdong Natural Science Foundation (No. 2016A030313750), Project of Enhancing School with Innovation of Guangdong Ocean University (No. GDOU2017052502), Zhanjiang Natural Science Foundation (No. A15445, No. A14457).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Paton, R.S.; Goodman, J.M.; Pellegrinet, S.C. Theoretical Study of the Asymmetric Conjugate Alkenylation of Enones Catalyzed by Binaphthols. *J. Org. Chem.* **2008**, *73*, 5078–5089. [[CrossRef](#)] [[PubMed](#)]
2. Hatakeyama, T.; Nakamura, M.; Nakamura, E. Diastereoselective Addition of Zincated Hydrazones to Alkenylboronates and Stereospecific Trapping of Boron/Zinc Bimetallic Intermediates by Carbon Electrophiles. *J. Am. Chem. Soc.* **2008**, *130*, 15688–15701. [[CrossRef](#)] [[PubMed](#)]
3. Carson, M.W.; Giese, M.W.; Coghlan, M.J. An Intra/Intermolecular Suzuki Sequence to Benzopyridyloxepines Containing Geometrically Pure Exocyclic Tetrasubstituted Alkenes. *Org. Lett.* **2008**, *10*, 2701–2704. [[CrossRef](#)] [[PubMed](#)]
4. Yang, M.; Yokokawa, N.; Ohmiya, H.; Sawamura, M. Synthesis of Conjugated Allenes through Copper-Catalyzed γ -Selective and Stereospecific Coupling between Propargylic Phosphates and Aryl-or Alkenylboronates. *Org. Lett.* **2012**, *14*, 816–819. [[CrossRef](#)] [[PubMed](#)]
5. He, Z.; Kirchberg, S.; Fröhlich, R.; Studer, A. Oxidative Heck Arylation for the Stereoselective Synthesis of Tetrasubstituted Olefins Using Nitroxides as Oxidants. *Angew. Chem. Int. Ed.* **2012**, *51*, 3699–3702. [[CrossRef](#)] [[PubMed](#)]
6. Pospiech, S.; Bolte, M.; Lerner, H.-W.; Wagner, M. Insertion Reactions into the Boron–Boron Bonds of Barrelene-Type 1,2-Diaminodiboranes (4). *Organometallics* **2014**, *33*, 6967–6974. [[CrossRef](#)]
7. Hyodo, K.; Suetsugu, M.; Nishihara, Y. Diborylation of Alkynyl MIDA Boronates and Sequential Chemoselective Suzuki–Miyaura Couplings: A Formal Carboborylation of Alkynes. *Org. Lett.* **2014**, *16*, 440–443. [[CrossRef](#)]
8. Mora-Radó, H.; Bialy, L.; Czechtizky, W.; Méndez, M.; Harrity, J.P.A. An Alkyne Diboration/ 6π -Electrocyclization Strategy for the Synthesis of Pyridine Boronic Acid Derivatives. *Angew. Chem. Int. Ed.* **2016**, *55*, 5834–5836. [[CrossRef](#)]
9. Kuang, Z.J.; Li, B.N.; Song, Q.L. Cu/Pd cooperatively catalyzed tandem intramolecular anti-Markovnikov hydroarylation of unsaturated amides: Facile construction of 3,4-dihydroquinolinones via borylation/intramolecular C(sp³)–C(sp²) cross coupling. *Chem. Commun.* **2018**, *54*, 34–37. [[CrossRef](#)]
10. Stephens, T.C.; Pattison, G. Transition-Metal-Free Homologative Cross-Coupling of Aldehydes and Ketones with Geminal Bis(boron) Compounds. *Org. Lett.* **2017**, *19*, 3498–3501. [[CrossRef](#)]
11. Kuang, Z.J.; Chen, H.H.; Yan, J.X.; Yang, K.; Lan, Y.; Song, Q.L. Base-Catalyzed Borylation/B–O Elimination of Propynols and B₂pin₂ Delivering Tetrasubstituted Alkenylboronates. *Org. Lett.* **2018**, *20*, 5153–5157. [[CrossRef](#)] [[PubMed](#)]
12. Okura, K.; Teranishi, T.; Yoshida, Y.; Shirakawa, E. Electron-Catalyzed Cross-Coupling of Arylboron Compounds with Aryl Iodides. *Angew. Chem. Int. Ed.* **2018**, *57*, 7186–7190. [[CrossRef](#)] [[PubMed](#)]
13. He, Z.Q.; Song, F.F.; Sun, H.; Huang, Y. Transition-Metal-Free Suzuki-Type Cross-Coupling Reaction of Benzyl Halides and Boronic Acids via 1,2-Metalate Shift. *J. Am. Chem. Soc.* **2018**, *140*, 2693–2699. [[CrossRef](#)] [[PubMed](#)]
14. Iafe, R.G.; Kuo, J.L.; Hochstatter, D.G.; Saga, T.; Turner, J.W.; Merlic, C.A. Increasing the Efficiency of the Transannular Diels–Alder Strategy via Palladium(II)-Catalyzed Macrocyclizations. *Org. Lett.* **2013**, *15*, 582–585. [[CrossRef](#)] [[PubMed](#)]
15. Shi, X.L.; Kiesman, W.F.; Levina, A.; Xin, Z.L. Catalytic Asymmetric Petasis Reactions of Vinylboronates. *J. Org. Chem.* **2013**, *78*, 9415–9423. [[CrossRef](#)] [[PubMed](#)]

16. Yoshida, H.; Kageyuki, I.; Takaki, K. Silver-Catalyzed Highly Regioselective Formal Hydroboration of Alkynes. *Org. Lett.* **2014**, *16*, 3512–3515. [[CrossRef](#)] [[PubMed](#)]
17. Obligacion, J.V.; Neely, J.M.; Yazdani, A.N.; Pappas, I.; Chirik, P.J. Cobalt Catalyzed Z-Selective Hydroboration of Terminal Alkynes and Elucidation of the Origin of Selectivity. *J. Am. Chem. Soc.* **2015**, *137*, 5855–5858. [[CrossRef](#)]
18. Huang, X.; Hu, J.J.; Wu, M.Y.; Wang, J.Y.; Peng, Y.Q.; Song, G.H. Catalyst-free chemoselective conjugate addition and reduction of α,β -unsaturated carbonyl compounds via a controllable boration/protodeboration cascade pathway. *Green Chem.* **2018**, *20*, 255–260. [[CrossRef](#)]
19. Lavergne, J.L.; Jayaraman, A.; Castro, L.C.M.; Rochette, É.; Fontaine, F.-G. Metal-Free Borylation of Heteroarenes Using Ambiphilic Aminoboranes: On the Importance of Sterics in Frustrated Lewis Pair C–H Bond Activation. *J. Am. Chem. Soc.* **2017**, *139*, 14714–14723. [[CrossRef](#)]
20. Farrell, J.M.; Schmidt, D.; Grande, V.; Wurthner, F. Synthesis of a Doubly Boron-Doped Perylene through NHC-Borenium Hydroboration/C–H Borylation/Dehydrogenation. *Angew. Chem. Int. Ed.* **2017**, *56*, 11846–11850. [[CrossRef](#)]
21. Pintaric, C.; Olivero, S.; Gimbert, Y.; Chavant, P.Y.; Duñach, E. An Opportunity for Mg-Catalyzed Grignard-Type Reactions: Direct Coupling of Benzylic Halides with Pinacolborane with 10 mol% of Magnesium. *J. Am. Chem. Soc.* **2010**, *132*, 11825–11827. [[CrossRef](#)] [[PubMed](#)]
22. Yang, C.-T.; Zhang, Z.-Q.; Tajuddin, H.; Wu, C.-C.; Liang, J.; Liu, J.-H.; Fu, Y.; Czyzewska, M.; Steel, P.G.; Marder, T.B.; et al. Alkylboronic Esters from Copper-Catalyzed Borylation of Primary and Secondary Alkyl Halides and Pseudohalides. *Angew. Chem. Int. Ed.* **2012**, *124*, 543–547. [[CrossRef](#)]
23. Kubota, K.; Iwamoto, H.; Ito, H. Formal nucleophilic borylation and borylative cyclization of organic halides. *Org. Biomol. Chem.* **2017**, *15*, 285–300. [[CrossRef](#)] [[PubMed](#)]
24. Verma, P.K.; Mandal, S.; Geetharani, K. Efficient Synthesis of Aryl Boronates via Cobalt-Catalyzed Borylation of Aryl Chlorides and Bromides. *ACS Catal.* **2018**, *8*, 4049–4054. [[CrossRef](#)]
25. Yoshida, T.; Iliés, L.; Nakamura, E. Iron-Catalyzed Borylation of Aryl Chlorides in the Presence of Potassium *t*-Butoxide. *ACS Catal.* **2017**, *7*, 3199–3203. [[CrossRef](#)]
26. Mkhaliid, I.A.I.; Barnard, J.H.; Marder, T.B.; Murphy, J.M.; Hartwig, J.F. C–H Activation for the Construction of C–B Bonds. *Chem. Rev.* **2010**, *110*, 890–931. [[CrossRef](#)] [[PubMed](#)]
27. Wei, C.S.; Jiménez-Hoyos, C.A.; Videa, M.F.; Hartwig, J.F.; Hall, M.B. Origins of the Selectivity for Borylation of Primary over Secondary C–H Bonds Catalyzed by Cp*-Rhodium Complexes. *J. Am. Chem. Soc.* **2010**, *132*, 3078–3091. [[CrossRef](#)]
28. Xu, L.; Wang, G.H.; Zhang, S.; Wang, H.; Wang, L.H.; Liu, L.; Jiao, J.; Li, P.F. Recent advances in catalytic C–H borylation reactions. *Tetrahedron* **2017**, *73*, 7123–7157. [[CrossRef](#)]
29. Jayaraman, A.; Castro, L.C.M.; Desrosiers, V.; Fontaine, F.G. Metal-free borylative dearomatization of indoles: Exploring the divergent reactivity of aminoborane C–H borylation catalysts. *Chem. Sci.* **2018**, *9*, 5057–5063. [[CrossRef](#)]
30. Lee, Y.; Jang, H.; Hoveyda, A.H. Vicinal Diboronates in High Enantiomeric Purity through Tandem Site-Selective NHC–Cu-Catalyzed Boron–Copper Additions to Terminal Alkynes. *J. Am. Chem. Soc.* **2009**, *131*, 18234–18235. [[CrossRef](#)]
31. Chen, Q.; Zhao, J.; Ishikawa, Y.; Asao, N.; Yamamoto, Y.; Jin, T. Remarkable Catalytic Property of Nanoporous Gold on Activation of Diborons for Direct Diboration of Alkynes. *Org. Lett.* **2013**, *15*, 5766–5769. [[CrossRef](#)] [[PubMed](#)]
32. Jung, H.-Y.; Yun, J. Copper-Catalyzed Double Borylation of Silylacetylenes: Highly Regio- and Stereoselective Synthesis of Syn-Vicinal Diboronates. *Org. Lett.* **2012**, *14*, 2606–2609. [[CrossRef](#)] [[PubMed](#)]
33. Bidal, Y.D.; Lazreg, F.; Cazin, C.S.J. Copper-Catalyzed Regioselective Formation of Tri- and Tetrasubstituted Vinylboronates in Air. *ACS Catal.* **2014**, *4*, 1564–1569. [[CrossRef](#)]
34. Yuan, W.M.; Ma, S.M. CuCl–K₂CO₃-catalyzed highly selective borylcupration of internal alkynes—Ligand effect. *Org. Biomol. Chem.* **2012**, *10*, 7266–7268. [[CrossRef](#)] [[PubMed](#)]
35. Jung, H.-Y.; Feng, X.H.; Kim, H.; Yun, J. Copper-catalyzed boration of activated alkynes. Chiral boranes via a one-pot copper-catalyzed boration and reduction protocol. *Tetrahedron* **2012**, *68*, 3444–3449. [[CrossRef](#)]

36. Zhao, F.; Jia, X.W.; Li, P.Y.; Zhao, J.W.; Zhou, Y.; Wang, J.; Liu, H. Catalytic and catalyst-free diboration of alkynes. *Org. Chem. Front.* **2017**, *4*, 2235–2255. [[CrossRef](#)]
37. Wen, Y.M.; Xie, J.Y.; Deng, C.M.; Li, C.D. Selective Synthesis of Alkylboronates by Copper(I)-Catalyzed Borylation of Allyl or Vinyl Arenes. *J. Org. Chem.* **2015**, *80*, 4142–4147. [[CrossRef](#)] [[PubMed](#)]
38. Zhao, H.; Tong, M.; Wang, H.; Xu, S. Transition-metal-free synthesis of 1,1-diboronate esters with a fully substituted benzylic center via diborylation of lithiated carbamates. *Org. Biomol. Chem.* **2017**, *15*, 3418–3422. [[CrossRef](#)]
39. Yamamoto, E.; Izumi, K.; Horita, Y.; Ito, H. Anomalous Reactivity of Silylborane: Transition-Metal-Free Boryl Substitution of Aryl, Alkenyl, and Alkyl Halides with Silylborane/Alkoxy Base Systems. *J. Am. Chem. Soc.* **2012**, *134*, 19997–20000. [[CrossRef](#)]
40. Zhang, J.M.; Wu, H.-H.; Zhang, J.L. Cesium Carbonate Mediated Borylation of Aryl Iodides with Diboron in Methanol. *Eur. J. Org. Chem.* **2013**, 6263–6266. [[CrossRef](#)]
41. Lee, Y.; Baek, S.; Park, J.; Kim, S.T.; Tussupbayev, S.; Kim, J.; Baik, M.H.; Cho, S.H. Chemoselective Coupling of 1,1-Bis[(pinacolato)boryl]alkanes for the Transition-Metal-Free Borylation of Aryl and Vinyl Halides: A Combined Experimental and Theoretical Investigation. *J. Am. Chem. Soc.* **2017**, *139*, 976–984. [[CrossRef](#)] [[PubMed](#)]
42. Miralles, N.; Romero, R.M.; Fernández, E.; Munñiz, K. A mild carbon-boron bond formation from diaryliodonium salts. *Chem. Commun.* **2015**, *51*, 14068–14071. [[CrossRef](#)] [[PubMed](#)]
43. Qi, X.X.; Li, H.-P.; Peng, J.-B.; Wu, X.-F. Borylation of aryldiazonium salts at room temperature in an aqueous solution under catalyst-free conditions. *Tetrahedron Lett.* **2017**, *58*, 3851–3853. [[CrossRef](#)]
44. Zhu, C.; Yamane, M. Transition-Metal-Free Borylation of Aryltriazene Mediated by $\text{BF}_3 \cdot \text{OEt}_2$. *Org. Lett.* **2012**, *14*, 4560–4563. [[CrossRef](#)] [[PubMed](#)]
45. Li, H.; Wang, L.; Zhang, Y.; Wang, J.B. Transition-Metal-Free Synthesis of Pinacol Alkylboronates from Tosylhydrazones. *Angew. Chem. Int. Ed.* **2012**, *51*, 2943–2946. [[CrossRef](#)] [[PubMed](#)]
46. Chena, K.; Wang, L.H.; Meng, G.; Li, P.F. Recent Advances in Transition-Metal-Free Aryl C–B Bond Formation. *Synthesis* **2017**, *49*, 4719–4730.
47. Liu, Y.L.; Kehr, G.; Daniliuc, C.G.; Erker, G. Metal-Free Arene and Heteroarene Borylation Catalyzed by Strongly Electrophilic Bis-boranes. *Chem. Eur. J.* **2017**, *23*, 12141–12144. [[CrossRef](#)]
48. McGough, J.S.; Cid, J.; Ingleson, M.J. Catalytic Electrophilic C–H Borylation Using NHC center dot Boranes and Iodine Forms C_2- , not C_3- , Borylated Indoles. *Chem. Eur. J.* **2017**, *23*, 8180–8184. [[CrossRef](#)]
49. Bonet, A.; Pubill-Ulldemolins, C.; Bo, C.; Gulyás, H.; Fernández, E. Transition-Metal-Free Diboration Reaction by Activation of Diboron Compounds with Simple Lewis Bases. *Angew. Chem. Int. Ed.* **2011**, *50*, 7158–7161. [[CrossRef](#)]
50. Miralles, N.; Cid, J.; Cuenca, A.B.; Carbó, J.J.; Fernández, E. Mixed diboration of alkenes in a metal-free context. *Chem. Commun.* **2015**, *51*, 1693–1696. [[CrossRef](#)]
51. Nagao, K.; Ohmiya, H.; Sawamura, M. Anti-Selective Vicinal Silaboration and Diboration of Alkynoates through Phosphine Organocatalysis. *Org. Lett.* **2015**, *17*, 1304–1307. [[CrossRef](#)] [[PubMed](#)]
52. Verma, A.; Snead, R.F.; Dai, Y.; Slebodnick, C.; Yang, Y.; Yu, H.; Yao, F.; Santos, W.L. Substrate-Assisted, Transition-Metal-Free Diboration of Alkynamides with Mixed Diboron: Regio- and Stereoselective Access to trans-1,2-Vinyldiboronates. *Angew. Chem. Int. Ed.* **2017**, *56*, 5111–5115. [[CrossRef](#)] [[PubMed](#)]
53. Nagashima, Y.; Hirano, K.; Takita, R.; Uchiyama, M. Trans-Diborylation of Alkynes: Pseudo-Intramolecular Strategy Utilizing a Propargylic Alcohol Unit. *J. Am. Chem. Soc.* **2014**, *136*, 8532–8535. [[CrossRef](#)] [[PubMed](#)]
54. Kojima, C.; Lee, K.-H.; Lin, Z.; Yamashita, M. Direct and Base-Catalyzed Diboration of Alkynes using The Unsymmetrical Diborane(4), pinB-BMes₂. *J. Am. Chem. Soc.* **2016**, *138*, 6662–6669. [[CrossRef](#)] [[PubMed](#)]
55. Morinaga, A.; Nagao, K.; Ohmiya, H.; Sawamura, M. Synthesis of 1,1-Diborylalkenes through a Brønsted Base Catalyzed Reaction between Terminal Alkynes and Bis(pinacolato)diboron. *Angew. Chem. Int. Ed.* **2015**, *54*, 15859–15862. [[CrossRef](#)] [[PubMed](#)]
56. Gao, G.L.; Kuang, Z.J.; Song, Q.L. Functionalized Geminal-diborylalkanes from Various Electron-Deficient Alkynes and B_2pin_2 . *Org. Chem. Front.* **2018**, *5*, 2249–2253. [[CrossRef](#)]

57. Bonet, A.; Sole, C.; Gulyás, H.; Fernández, E. Asymmetric organocatalytic diboration of alkenes. *Org. Biomol. Chem.* **2012**, *10*, 6621–6623. [[CrossRef](#)] [[PubMed](#)]
58. Fang, L.; Yan, L.; Haeffner, F.; Morken, J.P. Carbohydrate-Catalyzed Enantioselective Alkene Diboration: Enhanced Reactivity of 1,2-Bonded Diboron Complexes. *J. Am. Chem. Soc.* **2016**, *138*, 2508–2511. [[CrossRef](#)] [[PubMed](#)]
59. Blaisdell, T.P.; Caya, T.C.; Zhang, L.; Sanz-Marco, A.; Morken, J.P. Hydroxyl-Directed Stereoselective Diboration of Alkenes. *J. Am. Chem. Soc.* **2014**, *136*, 9264–9267. [[CrossRef](#)] [[PubMed](#)]
60. Lee, K.-S.; Zhugralin, A.R.; Hoveyda, A.H. Efficient C-B Bond Formation Promoted by N-Heterocyclic Carbenes: Synthesis of Tertiary and Quaternary B-Substituted Carbons through Metal-Free Catalytic Boron Conjugate Additions to Cyclic and Acyclic α,β -Unsaturated Carbonyls. *J. Am. Chem. Soc.* **2009**, *131*, 7253–7255. [[CrossRef](#)] [[PubMed](#)]
61. Kleeberg, C.; Crawford, A.G.; Batsanov, A.S.; Hodgkinson, P.; Apperley, D.C.; Cheung, M.S.; Lin, Z.; Marder, T.B. Spectroscopic and Structural Characterization of the CyNHC Adduct of B₂pin₂ in Solution and in the Solid State. *J. Org. Chem.* **2012**, *77*, 785–789. [[CrossRef](#)] [[PubMed](#)]
62. Wu, H.; Radomkit, S.; O'Brien, J.M.; Hoveyda, A.H. Metal-Free Catalytic Enantioselective C–B Bond Formation: (Pinacolato)boron Conjugate Additions to α,β -Unsaturated Ketones, Esters, Weinreb Amides, and Aldehydes Promoted by Chiral N-Heterocyclic Carbenes. *J. Am. Chem. Soc.* **2012**, *134*, 8277–8285. [[CrossRef](#)] [[PubMed](#)]
63. Radomkit, S.; Hoveyda, A.H. Enantioselective Synthesis of Boron-Substituted Quaternary Carbon Stereogenic Centers through NHC-Catalyzed Conjugate Additions of (Pinacolato)boron Units to Enones. *Angew. Chem. Int. Ed.* **2014**, *53*, 3387–3391. [[CrossRef](#)] [[PubMed](#)]
64. Wang, L.; Chen, Z.; Ma, M.; Duan, W.; Song, C.; Ma, Y. Synthesis and application of a dual chiral [2.2]paracyclophane-based N-heterocyclic carbene in enantioselective β -boration of acyclic enones. *Org. Biomol. Chem.* **2015**, *13*, 10691–10698. [[CrossRef](#)] [[PubMed](#)]
65. Wen, K.; Chen, J.; Gao, F.; Bhadury, P.S.; Fanb, E.; Sun, Z. Metal free catalytic hydroboration of multiple bonds in methanol using N-heterocyclic carbenes under open atmosphere. *Org. Biomol. Chem.* **2013**, *11*, 6350–6356. [[CrossRef](#)] [[PubMed](#)]
66. Pubill-Ulldemolins, C.; Bonet, A.; Bo, C.; Gulyás, H.; Fernández, E. Activation of Diboron Reagents with Brønsted Bases and Alcohols: An Experimental and Theoretical Perspective of the Organocatalytic Boron Conjugate Addition Reaction. *Chem. Eur. J.* **2012**, *18*, 1121–1126. [[CrossRef](#)]
67. Cid, J.; Carbó, J.J.; Fernández, E. A Clear-Cut Example of Selective Bpin-Bdan Activation and Precise Bdan Transfer on Boron Conjugate Addition. *Chem. Eur. J.* **2014**, *20*, 3616–3620. [[CrossRef](#)]
68. Cascia, E.L.; Sanz, X.; Bo, C.; Whiting, A.; Fernández, E. Asymmetric metal free β -boration of α,β -unsaturated imines assisted by (S)-MeBoPhoz. *Org. Biomol. Chem.* **2015**, *13*, 1328–1332. [[CrossRef](#)]
69. Bonet, A.; Gulyás, H.; Fernández, E. Metal-Free Catalytic Boration at the β -Position of α,β -Unsaturated Compounds: A Challenging Asymmetric Induction. *Angew. Chem. Int. Ed.* **2010**, *49*, 5130–5134. [[CrossRef](#)]
70. Pubill-Ulldemolins, C.; Bonet, A.; Gulyás, H.; Bo, C.; Fernández, E. Essential role of phosphines in organocatalytic β -boration reaction. *Org. Biomol. Chem.* **2012**, *10*, 9677–9682. [[CrossRef](#)]
71. Ibrahim, I.; Breistein, P.; Córdova, A. One-Pot Three-Component Highly Selective Synthesis of Homoallylboronates by Using Metal-Free Catalysis. *Chem. Eur. J.* **2012**, *18*, 5175–5179. [[CrossRef](#)] [[PubMed](#)]
72. Miralles, N.; Alam, R.; Szabó, K.J.; Fernández, E. Transition-Metal-Free Borylation of Allylic and Propargylic Alcohols. *Angew. Chem. Int. Ed.* **2016**, *55*, 4303–4307. [[CrossRef](#)] [[PubMed](#)]
73. Harada, K.; Nogami, M.; Hirano, K.; Kurauchi, D.; Kato, H.; Miyamoto, K.; Saito, T.; Uchiyama, M. Allylic borylation of tertiary allylic alcohols: A divergent and straightforward access to allylic boronates. *Org. Chem. Front.* **2016**, *3*, 565–569. [[CrossRef](#)]
74. Nogami, M.; Hirano, K.; Kanai, M.; Wang, C.; Saito, T.; Miyamoto, K.; Muranaka, A.; Uchiyama, M. Transition Metal-Free trans-Selective Alkynylboration of Alkynes. *J. Am. Chem. Soc.* **2017**, *139*, 12358–12361. [[CrossRef](#)] [[PubMed](#)]
75. Yanga, K.; Song, Q. Transition-metal-free regioselective synthesis of alkylboronates from arylacetylenes and vinyl arenes. *Green Chem.* **2016**, *18*, 932–936. [[CrossRef](#)]
76. Hong, S.B.; Zhang, W.; Liu, M.Y.; Yao, Z.J.; Deng, W. Transition-metal-free hydroboration of terminal alkynes activated by base. *Tetrahedron Lett.* **2016**, *57*, 1–4. [[CrossRef](#)]

77. Deng, C.M.; Ma, Y.F.; Wen, Y.M. Transition-Metal-Free Borylation of Alkynes and Alkenes. *Chem. Sel.* **2018**, *3*, 1202–1204. [[CrossRef](#)]
78. Davenport, E.; Fernández, E. Transition-metal-free synthesis of vicinal triborated compounds and selective functionalisation of the internal C–B bond. *Chem. Commun.* **2018**, *54*, 10104–10107. [[CrossRef](#)]
79. García, L.; Sendra, J.; Miralles, N.; Reyes, E.; Carbó, J.J.; Vicario, J.L.; Fernández, E. Transition-metal-free stereoselective borylation of allenamides. *Chem. Eur. J.* **2018**, *24*, 14059–14063. [[CrossRef](#)]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).