

Supplementary Information

Anti-Hepatitis B Virus Activity of Esculetin from *Microsorium fortunei* In Vitro and In Vivo

Si-Xin Huang^{1,†}, Jun-Fei Mou^{2,†}, Qin Luo³, Qing-Hu Mo¹, Xian-Li Zhou^{2,*}, Xiao Huang¹, Qing Xu⁴, Xiang-Duan Tan¹, Xu Chen¹, Cheng-Qin Liang^{1,**}

¹ College of Pharmacy, Guilin Medical University, Guilin 541004, Guangxi, China; huangsixin@stu.glmc.edu.cn (S.-X. H.); moxqxom@163.com (Q.-H. M.); gyhuangxiao@163.com (X. H.); tandu@glmc.edu.cn (X.-D. T.); chenxu@glmc.edu.cn (X.C.); cqliang@glmc.edu.cn (C.-Q. L.)

² Biotechnology Institute, Guilin Medical University, Guilin 541004, Guangxi, China; qiaoyifei@stu.glmc.edu.cn (J.-F.M.); xlzhou2019@glmc.edu.cn (X.-L. Z.)

³ Science Experiment Center, Guilin Medical University, Guilin 541004, Guangxi, China; luoqin5020@glmc.edu.cn (Q. L.)

⁴ Guangxi Key Laboratory of Molecular Medicine in Liver Injury and Repair, Guilin Medical University, Guilin 541004, Guangxi, China; xq5895801@163.com (Q. X.)

* Correspondence: cqliang@glmc.edu.cn (C.-Q.L.); xlzhou2019@glmc.edu.cn (X.-L. Z.); Tel:+0773-5891498 (C.-Q.L.); +0773-3680722 (X.-L. Z.); Fax:+0773-5894158 (C.-Q.L.); +0773-3680722 (X.-L. Z.)

Figure S1. High performance liquid chromatogram of esculetin.

Figure S2. ^1H -NMR of esculetin ($\text{DMSO-}d_6$, 500 MHz).

Figure S3. ^{13}C -NMR of esculetin ($\text{DMSO-}d_6$, 125 MHz).




Figure S1. HPLC of esculetin: Wave length = 349 nm; Flow rate = 1 mL/min; Column temperature = 36 °C; Mobile phase: Acetonitrile :0.5% acid water = 7:93; t_R = 15.479min.




Figure S2. ^1H -NMR of esculetin ($\text{DMSO-}d_6$, 500 MHz).




Figure S3. ^{13}C -NMR of esculetin ($\text{DMSO-}d_6$, 125 MHz).