HPLC profile of longan pericarp-sourced phenolics and their antioxidant and cytotoxic effects

Xuelian Bai, Rui Pan, Mingzhu Li, Xiuting Li *, Huawei Zhang *

Figure S1	¹ H NMR spectrum of A (600 MHz, in MeOH-d ₄)	S2
Figure S2	HR-ESI-MS spectrum of A	S3
Figure S3	¹ H NMR spectrum of B (600 MHz, in MeOH-d ₄)	S4
Figure S4	HR-ESI-MS spectrum of B	S5
Figure S5	¹ H NMR spectrum of C (600 MHz, in MeOH-d ₄)	\$6
Figure S6	HR-ESI-MS spectrum of C	S7
Figure S7	¹ H NMR spectrum of D (600 MHz, in DMSO-d ₆)	S8
Figure S8	HR-ESI-MS spectrum of D	S9
Figure S9	¹ H NMR spectrum of E (600 MHz, in MeOH-d ₄)	S10
Figure S10	HR-ESI-MS spectrum of E	\$11
Figure S11	¹ H NMR spectrum of F (600 MHz, in MeOH-d ₄)	S12
Figure S12	HR-ESI-MS spectrum of F	S13

Table of Contents



Figure S1 ¹H NMR spectrum of A (600 MHz, in MeOH-d₄)



Figure S2 HR-ESI-MS spectrum of A



Figure S3 ¹H NMR spectrum of B (600 MHz, in MeOH-d₄)



Figure S4 HR-ESI-MS spectrum of B



Figure S5 ¹H NMR spectrum of C (600 MHz, in MeOH-d₄)



Figure S6 HR-ESI-MS spectrum of C



Figure S7 ¹H NMR spectrum of D (600 MHz, in DMSO-d₆)

Analysis Info					
Analysis Inio	D:\Data\2018\1222\G_d			Acquisition Date	12/22/2018 3:00:38 PM
Method	tune_pos_low2.m			Operator	BDAL@DE
Sample Name Comment	G			Instrument / Ser#	micrOTOF-Q II 10366
Acquisition Par	ameter				
Source Type	ESI	Ion Polarity	Positive	Set Nebulizer	0.3 Bar
Scan Begin	100 m/z	Set End Plate Offset	-550 V	Set Dry Gas	4.0 Vmin
Scan End	1000 m/z	Set Collision Cell RF	300.0 Vpp	Set Divert Val	ve Waste
Intens.					+MS, 0.0min #
2.5					
2.0					
1.5					324.9962
1.0					
0.5 301 1400) 307.0608	312.7403			
				320.0397 322.	9758
300	305	310	315	320	325 m/

Figure S8 HR-ESI-MS spectrum of D



Figure S9 ¹H NMR spectrum of E (600 MHz, in MeOH-d₄)



Figure S10 HR-ESI-MS spectrum of E



Figure S11 ¹H NMR spectrum of F (600 MHz, in MeOH-d₄)



Figure S12 HR-ESI-MS spectrum of F