Supplementary information

RA-XII Suppresses the Development and Growth of Liver Cancer by Inhibition of Lipogenesis via SCAP-dependent SREBP Supression

Di Guo ^{1,†}, Yurong Wang ^{1,†}, Jing Wang ¹, Lihua Song ¹, Zhe Wang ^{1,2}, Bingyu Mao ³ and Ninghua Tan ^{1,2,*}

- ¹ State Key Laboratory of Natural Medicines, Department of TCMs Pharmaceuticals, School of Traditional Chinese Pharmacy, China Pharmaceutical University, 211198 Nanjing, China; guodi33@163.com (D.G.); yurong1987213@163.com (Y.W.); 18851107621@163.com (J.W.); songlihua4835@163.com (L.S.); wangzhe153807105@163.com (Z.W.)
- ² State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, 650201 Kunming, China
- ³ State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, 650223 Kunming, China; mao@mail.kiz.ac.cn
- * Correspondence: nhtan@cpu.edu.cn; Tel.: +86-25-8618-5772
- ⁺ These authors contributed equally to this work.



Figure S1. SRB measurement of cell vitality. Cells were treated with different concentrations of RA-XII for 48 h. Results are means ± SEM of three independent experiments.



Figure S2. Hematoxylin and eosin staining of kidney, liver and lung sections of the xenograft mouse model (Scale bar: 75 μ m).



Reported by User: Public	Project Name: HYY
Report Method: Overlay Report	Date Printed:
Report Method ID: 6288	4/26/2019
Page: 1 of 1	4:19:21 PM PRC

Figure S3. HPLC-DAD chromatogram of RA-XII, which indicates that RA-XII used in this study is pure.