

Supplementary Materials for

Study on the Microwave-Assisted Batch and Continuous Flow Synthesis of N-Alkyl-Isoindolin-1-One-3-Phosphonates by a Special Kabachnik-Fields Condensation

Adám Tajti ¹, Nóra Tóth ¹, Bettina Rávai ¹, István Csontos ¹, Pál Tamás Szabó ² and Erika Bálint ^{1,*}

¹ Department of Organic Chemistry and Technology, Budapest University of Technology and Economics, 1521 Budapest, Hungary; tajti.adam@mail.bme.hu (Á.T.); toth.nora@mail.bme.hu (N.T.); betti.ravai08@gmail.com (B.R.); icsontos@mail.bme.hu (I.C.)

² MS Metabolomics Laboratory, Instrumentation Center, Research Centre for Natural Sciences, Hungarian Academy of Sciences, Magyar tudósok krt. 2., H-1117 Budapest, Hungary; szabo.pal@ttk.mta.hu

* Correspondence: ebalint@mail.bme.hu; Tel.: +36-1-463-3653

Table of contents

Figures of the continuous flow system Figure S2

³¹P NMR, ¹H NMR and ¹³C NMR spectra




Figure S1. Design parameters of the continuous flow system.




Figure S2. Sketch of the continuous flow cell.

^{31}P NMR, ^1H NMR and ^{13}C NMR spectra

