

Supplemental Material

Simple synthesis of 17- β -O-hemisuccinate of stanozolol for immunoanalytical methods

Silvana Casati^a, Roberta Ottria^a, Pierangela Ciuffreda^{a, #}

^a *Dipartimento di Scienze Biomediche e Cliniche "Luigi Sacco", Università degli Studi di Milano, Via G.B. Grassi 74,
20157 Milano, Italy
silvana.casati@unimi.it, roberta.ottria@unimi.it; pierangela.ciuffreda@unimi.it*

Table of Contents

Figures

Figure S1a/S1b	Pag. 2
¹H and ¹³C-NMR data	
¹ H spectrum of mixture 2a/2b	Pag. 3
¹ H spectrum of mixture 3a/3b	Pag. 4
¹ H spectrum of mixture 4	Pag. 5
¹ H and ¹³ C NMR spectra of stanozolol 17 β -O-hemisuccinate (1)	Pag. 6




Figure S1a. ^1H -NMR spectrum from 8.40 ppm to 6.00 ppm of Stanozolol (top) and mixture **2a/2b** (bottom).








Figure S1b. ^1H -NMR spectrum from 4.20 ppm to 2.00 ppm of Stanozolol (top) and mixture **2a/2b** (bottom).



^1H spectrum of mixture **2a/2b**



^1H spectrum of mixture **3a/3b**



^1H spectrum of compound **4**



^1H and ^{13}C NMR spectra of stanazolol 17 β -O-Hemisuccinate (**1**)

