

Silver Nanoparticles Using Eucalyptus or Willow Extracts (AgNPs) as Contact Lens Hydrogel Components to Reduce the Risk of Microbial Infection

Andreas K. Rossos ¹, Christina N. Banti ^{1,*}, Panagiotis K. Raptis ¹, Christina Papachristodoulou ², Ioannis Sainis ³, Panagiotis Zoumpoulakis ⁴, Thomas Mavromoustakos ⁵ and Sotiris K. Hadjikakou ^{1,6,*}

¹ Section of Inorganic and analytical Chemistry, Department of Chemistry, University of Ioannina, 45110 Ioannina, Greece; arossos@uoi.gr (A.K.R.); panagiwtisraptis93@yahoo.gr (P.K.R.)

² Department of Physics, University of Ioannina, 45110 Ioannina, Greece; xpapaxri@uoi.gr

³ Cancer Biobank Center, University of Ioannina, 45110 Ioannina, Greece; isainis@uoi.gr

⁴ Laboratory of Chemistry, Analysis and Design of Food Processes, Department of Food Science and Technology, University of West Attica, Egaleo, 12243 Attica, Greece; pzoump@eie.gr

⁵ Organic Chemistry Laboratory, Department of Chemistry, University of Athens Greece, 15571 Athens, Greece; tmavrom@chem.uoa.gr

⁶ Institute of Materials Science and Computing, University Research Center of Ioannina (URCI), 45110 Ioannina, Greece

* Correspondence: cbanti@uoi.gr (C.N.B.), shadjika@uoi.gr (S.K.H.); Tel.: +302651008362 (C.N.B.), +302651008374 (S.K.H.)




Figure S1. TGA diagrams of AgNPs(ELE) (A), AgNPs(WBE) (B), pHEMA@AgNPs(ELE)_2 (C) and pHEMA@AgNPs(WBE)_2 (D).




Figure S2. Bacteria viability of pHEMA, pHEMA@ELE_2, pHEMA@AgNPs(ELE)_2, pHEMA@WBE_2 and pHEMA@AgNPs(WBE)_2 against *P. aeruginosa* (A), *S. epidermidis* (B), *S. aureus* (C).