

Supplementary Materials

The Inhibitory Effects of Plant Derivate Polyphenols on the Main Protease of SARS Coronavirus 2 and Their Structure–Activity Relationship

Thi Thanh Hanh Nguyen ¹, Jong-Hyun Jung ², Min-Kyu Kim ², Sangyong Lim ²,
Jae-Myoung Choi ³, Byoungsang Chung ³, Do-Won Kim ⁴ and Doman Kim ^{1,5,*}

¹ Institutes of Green Bioscience and Technology, Seoul National University, Pyeongchang-gun, Gangwon-do 25354, Korea; hara2910@snu.ac.kr

² Radiation Research Division, Korea Atomic Engery Research Institute, Jeongeup 56212, Korea; Jungjh83@kaeri.re.kr (J.-H.J.); mkkim@kaeri.re.kr (M.-K.K.); saylim@kaeri.re.kr (S.L.)

³ Ottogi Sesame Mills Co., Ltd., Chungcheongbuk-do 27623, Korea; mjchoi@ottogism.co.kr (J.-M.C.); bschung@ottogism.co.kr (B.C.)

⁴ Department of Physics, Gangneung-Wonju National University, Gangneung 25457, Korea; Do.Won.Kim@cern.ch

⁵ Graduate School of International Agricultural Technology, Seoul National University, Pyongchang-gun, Gangwon-do 25354, Korea

* Correspondence: kimdm@snu.ac.kr; Tel.: +82-33-339-5720; Fax: +82-33-339-5716




Figure S1 Michaelis-Menten plot for determination of K_m value of purified SARS-CoV-2 M^{pro}.