

Supporting Information

Mononuclear Tricoordinate Copper(I) and Silver(I) Halide Complexes of a Sterically Bulky Thiourea Ligand and a Computational Insight of their Interaction with Human Insulin


Awal Noor ^{1,*}, Sadaf Qayyum ¹, Farukh Jabeen ² and Ashfaq Ur Rehman ³


¹ Department of Basic Sciences, Preparatory Year Deanship, King Faisal University, Al-Hassa 31982, Saudi Arabia; sqayyum@kfu.edu.sa

² Department of Chemistry and Biochemistry, Laurentian University, 935 Ramsey Lake Road, Sudbury, ON P3E 2C6, Canada; fjabeen@laurentian.ca

³ Department of Molecular Biology and Biochemistry, University of California, Irvine, CA 92697-3900, USA; aurehman@uci.edu

* Correspondence: anoor@kfu.edu.sa








Figure S4. ¹³C NMR spectrum of **1b**.




Figure S5. ¹H NMR spectrum of **1c**.




Figure S6. ¹³C NMR spectrum of 1c.




Figure S7. ¹H NMR spectrum of 2a.




Figure S8. ¹³C NMR spectrum of 2a.




Figure S9. ¹H NMR spectrum of 2b.




Figure S10. ¹³C NMR spectrum of **2b**.




Figure S11. ¹H NMR spectrum of **2c**.




Figure S12. ¹³C NMR spectrum of **2c**.