

Phytochemicals from *Vanda bensonii* and Their Bioactivities to Inhibit Growth and Metastasis of Non-Small Cell Lung Cancer Cells

Tajudeen O. Jimoh ^{1,2}, Narawat Nuamnaichati ³, Rungroch Sungthong ³, Chaisak Chansriniyom ^{4,5}, Pithi Chanvorachote ^{6,7}, Kittisak Likhitwitayawuid ⁴, Chatchai Chaotham ^{3,7,*} and Boonchoo Sritularak ^{4,8,*}

¹ Pharmaceutical Sciences and Technology Program, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand

² Department of Biochemistry, Faculty of Health Sciences, Islamic University in Uganda, Kampala P.O. Box 7689, Uganda

³ Department of Biochemistry and Microbiology, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand

⁴ Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand

⁵ Natural Products and Nanoparticles Research Unit, Chulalongkorn University, Bangkok 10330, Thailand

⁶ Department of Pharmacology and Physiology, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand

⁷ Center of Excellence in Cancer Cell and Molecular Biology, Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand

⁸ Center of Excellence in Natural Products for Ageing and Chronic Diseases, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand;

* Correspondence: cchoatham@gmail.com (C.Chaotham); Tel.: (+66 (0) 2218 8372), boonchoo.sr@chula.ac.th (B.S.); Tel. (+66 (0) 2218 8356)

Figure S1. Sigmoidal dose-response plots of NCI-H460 cells treated with varying concentrations of *Vanda bensonii* metabolites. Percentages of cell viability after treating cells with varying concentrations of *V. bensonii* metabolites for 24 h, assessed by a colorimetric method. All experiments were done at least in triplicates, and 0.5% (v/v) dimethyl sulfoxide served as the vehicle control. The graphical results represent means \pm standard deviations.

