Molecules 1999, 4, M120

4-{[3-cyano-4-(methoxymethyl)-6-methyl-2-pyridinyl]oxy}benzoic acid

Yelena Kaigorodova, Vladimir Vasilin and Gennady Krapivin

Department of organic chemistry, Kuban State Technological University, Moskovskaya, 2, Krasnodar, 350072, Russian Federation. Phone +7 8612 559556, E-mail: organics@kubstu.ru

Received: 13 October 1999 / Accepted: 14 October 1999 / Published: 23 November 1999

4-{[3-cyano-4-(methoxymethyl)-6-methyl-2-pyridinyl]oxy} benzoic acid **3** was prepared by the reaction of the 2-chloro-3-cyanopyridine **1** with 4-hydroxybenzoic acid **2** using a literature procedure [1].

A mixture of 1.84g (10 mmol) of chlorocyanopyridine in 30 ml of ethanol, 1.38g (10 mmol) of 4-hydroxybenzoic acid and 1.12g (20 mmol) of potassium hydroxide in 8 ml water was boiled for 40 hours (water bath temperature). The reaction mixture was diluted with 5 ml of water and acidified with 10% aqueous hydrochloric acid until a precipitate formed. The precipitate was filtered and recrystallized from ethanol. Compound **3** - colourless solid. Yield 1.83g (61%).

M.p. 177°C (ethanol).

¹H NMR (DMSO-d₆, 60MHz): 2.38 (s, 3H, CH₃); 2.53 (s, 3H, CH₂OCH₃); 4.64 (s, 2H, CH₂OCH₃); 7.23 (s, 1H, H_{Het}), 7.32 (d, J=7Hz, 2H, H_{Ar}(1)), 8.03 (d, J=7Hz, 2H, H_{Ar}(2)), 11.72 (broad s, 1H, OH).

IR (vaseline oil, cm⁻¹): 3270-3150, 3150-3050, 2230, 1725, 1630, 1625, 1575, 1520, 1480, 1415, 1400, 1360, 1240, 1195, 1145, 1050, 1020, 980, 920, 900, 870, 850, 815, 790, 725.

Anal. calc. for C₁₆ H₁₄ N₂ O₄ (298.29): C 64.42, H 4.73, N 9.39; found: C 64.31, H 4.45, N 9.42.

References

1. Kulnevich, V.G.; Kaigorodova, Ye.A.; Arustamova, I.S.; Korobchenko, L.V; Vladiko, G.V; Boreko, E.I. *Khim.-Pharm. Zhurnal* **1990**, 25, 132.

Sample availability: available from the authors and MDPI.

©1999 MDPI. All rights reserved. *Molecules* website <u>www.mdpi.org/molecules</u>/

1 von 1 05.05.2009 15:30