

*Molecules* **2001**, *6*, M238

### 3-Amino-9-methyl-2*H*,6*H*--[1,2,4]triazino[5,4-*b*][1,3,4]thiadiazin-6-one

Majid M. Heravi<sup>1,3</sup>, Ghadir Rajabzadeh<sup>1</sup>, Mohammad Rahimizadeh<sup>1</sup>, Mehdi Bakavoli<sup>1</sup> and Mitra Ghassemzadeh<sup>2</sup>


1. Department of Chemistry, School of Sciences, Ferdowsi University of Mashhad, Iran

2. Chemistry & Chemical Engineering Research Center of Iran, P.O. Box 14335-186, Tehran, Iran.

E-mail: [mghassemzadeh@gmx.de](mailto:mghassemzadeh@gmx.de)

3. Department of Chemistry, School of Sciences, Azzahra University, Vanak, Tehran, Iran. E-mail: mmheravi@azzahra.ac.ir

Received: 17 December 2001 / Accepted: 15 May 2001 / Published: 25 May 2001



Condensation of 4-amino-6-methyl-5-thio-1,2,4-triazin-3-one **1** with chloro acetonitrile in the presence of triethyl amine afforded the corresponding 5-thiocyanomethyl derivative **2**. The latter was refluxed in a mixture of acetonitrile and triethylamine to give the title compound **3**. Compound **1** (0.316 g, 2 mmol) was dissolved in a solution of acetonitrile (8 mL) and triethylamine (0.6 mL). To this solution chloro acetonitrile (0.2 mL, excess) was added. The reaction mixture was refluxed for 12 hrs at room temperature. The solvent was evaporated and the residue was washed with hot EtOH to afford **2**. The latter (0.2 g, 1 mmol) was refluxed in a mixture of acetonitrile (5 mL) and triethylamine (0.3 mL) for 4 hrs. The solvent was evaporated and the residue was crystallized from CHCl<sub>3</sub> to afford **3**.

Selected Data for **2**. Yield: 72%, mp.: 188-9° C, <sup>1</sup>HNMR (CDCl<sub>3</sub>)d, 2.3(s, 3H, Me), 4.1(s, 2H, CH<sub>2</sub>), 5.9(s, 2H, NH<sub>2</sub>, exchangeable with D<sub>2</sub>O). IR,  $\tilde{\nu}$  (KBr disc): 3400, 3300, 2400, 1690, 1210, 1100 cm<sup>-1</sup>, M.S., m/z, M<sup>+</sup>, 196(3), 195(44), 194(12), 193(100), 177(47), 56(72).

Selected data for **3**. Yield: 75%, mp.: 159-60° C, <sup>1</sup>HNMR d (CDCl<sub>3</sub>), 2.47(s, 3H, Me), 4.88(s, 2H, CH<sub>2</sub>), 6.55(s, 2H, NH<sub>2</sub>, exchangeable with D<sub>2</sub>O). IR,  $\tilde{\nu}$  (KBr disc): 3450, 1690, 1400, 1110 cm<sup>-1</sup>, M.S., m/z, M<sup>+</sup>, 196(3), 195(11), 194(18), 73(100), 42(23).

#### Reference

1. Heravi, M.M.; Rajabzadeh, Gh.; Rahimizadeh, M.; Bakavoli, M.; Ghassemzadeh, M. Work on *P*, *S*, *Si* and related elements. Submitted in **2000**.

*Sample Availability*: Available from the authors and from MDPI.

© 2001 MDPI. All rights reserved. *Molecules* website <http://www.mdpi.org/molecules/>