

Article

Two Distinct Channels Mediated by m2mAChR and $\alpha 9$ nAChR Co-Exist in Type II Vestibular Hair Cells of Guinea Pig

Tao Zhou ^{1,†}, Yi Wang ^{1,†}, Chang-Kai Guo ^{1,†}, Wen-Juan Zhang ¹, Hong Yu ¹, Kun Zhang ¹
and Wei-Jia Kong ^{1,2,3,*}

¹ Department of Otorhinolaryngology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, 1277 Jiefang Avenue, Wuhan 430022, China; E-Mails: zhoutao2012ent@yahoo.cn (T.Z.); wangyi821@yahoo.cn (Y.W.); ckguo2255@sina.com (C.-K.G.); juan_364@163.com (W.-J.Z.); yuhong_0706@163.com (H.Y.); kkwd_010@163.com (K.Z.)

² Institute of Otorhinolaryngology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, 1277 Jiefang Avenue, Wuhan 430022, China

³ Key Laboratory of Neurological Disorders of Education Ministry, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, China

[†] These authors contributed equally to this work.

* Author to whom correspondence should be addressed; E-Mail: entwjkong@yahoo.com.cn; Tel.: +86-27-8572-6900; Fax: +86-27-8577-6343.

Received: 25 February 2013; in revised form: 29 March 2013 / Accepted: 17 April 2013 /

Published: 24 April 2013

Abstract: Acetylcholine (ACh) is the principal vestibular efferent neurotransmitter among mammals. Pharmacologic studies prove that ACh activates a small conductance Ca^{2+} -activated K^{+} channels (KCa) current (SK2), mediated by $\alpha 9$ -containing nicotinic ACh receptor ($\alpha 9$ nAChR) in mammalian type II vestibular hair cells (VHCs II). However, our studies demonstrate that the m2 muscarinic ACh receptor (m2mAChR) mediates a big conductance KCa current (BK) in VHCs II. To better elucidate the correlation between these two distinct channels in VHCs II of guinea pig, this study was designed to verify whether these two channels and their corresponding AChR subtypes co-exist in the same VHCs II by whole-cell patch clamp recordings. We found that m2mAChR sensitive BK currents were activated in VHCs II isolated by collagenase IA, while $\alpha 9$ nAChR sensitive SK2 currents were activated in VHCs II isolated by trypsin. Interestingly, after exposing the patched cells isolated by trypsin to collagenase IA for 3 min, the $\alpha 9$ nAChR sensitive

SK2 current was abolished, while m2mAChR-sensitive BK current was activated. Therefore, our findings provide evidence that the two distinct channels and their corresponding AChR subtypes may co-exist in the same VHCs II, and the alternative presence of these two ACh receptors-sensitive currents depended on isolating preparation with different enzymes.

Keywords: calcium-dependent potassium channel; muscarinic ACh receptor; nicotinic ACh receptor; vestibular hair cell

1. Introduction

Hair cells in a higher vertebrate's vestibule are generally divided into two categories: the flask shaped type I vestibular hair cells (VHCs I) and the cylindrical shaped type II vestibular hair cells (VHCs II) [1]. VHCs II are innervated by a number of button-shaped efferent nerve endings, which are cholinergic efferent fibers demonstrated by investigation of ACh-esterase activities in rat [2] and studies of choline acetyltransferase (ChAT) activities in rat [2,3], human [4] and mouse [5]. All these results support that acetylcholine (ACh) is the principal inhibitory neurotransmitter mediating the efferent activities to the vestibule [6].

Cholinergic neurotransmission is mediated by two classes of ACh receptors (AChRs): ionotropic nicotinic AChRs (nAChRs), which are nonselective cation channels, and G-protein-coupled muscarinic AChRs (mAChRs), which act through second-messenger systems [7,8]. The differential expressions of $\alpha 2$ -7, $\alpha 9$ $\alpha 10$, $\beta 2$ -4nAChR subunits [9] and m1-m5mAChR subtypes [10] have been observed in the vestibular end organs of rat. Besides, m1, m2, m5mAChR subtypes have also been observed in human vestibular periphery by reverse transcription polymerase chain reaction (RT-PCR) studies [10,11]. Recently, immunohistochemical studies have further supported that m1-m5 mAChR subtypes were located in isolated VHCs II of pigeons [12], and RT-PCR has also provided strong evidence that m1-m5 mAChR subtypes were detected in isolated VHCs II of guinea pig [13].

Previous pharmacologic studies have proved that ACh activated a small conductance Ca^{2+} -activated K^{+} (KCa) current (SK2) in mammalian VHCs II and outer hair cells (OHCs), which was mediated by the well-known $\alpha 9$ nAChR and sensitive to apamin and strychnine [14–16]. However, our recent studies showed that ACh activated a big conductance KCa (BK) current in VHCs II of guinea pig, which was mediated by m2mAChR and sensitive to iberitoxin (IBTX) and methoctramine [17–19], yet insensitive to strychnine and apamin. These two discrepant results implied that m2mAChR mediated BK currents and $\alpha 9$ nAChR mediated SK2 currents did not appear at the same time, which also indicated that the two distinct channels mediated by m2mAChR and $\alpha 9$ nAChR may exist in different VHCs II.

We found that collagenase IA was conventionally used to dissociate VHCs II in our studies [17–19], while most of the other researchers usually isolated mammalian VHCs II and OHCs by trypsin or collagenase IV [20,21]. So it was speculated that $\alpha 9$ nAChR/SK2 complex might undergo proteolysis during the collagenase IA isolation process. Indeed, the enzymes could alter physiological responses of $\alpha 9$ nAChR/SK2 complex. It has been reported that $\alpha 9$ nAChR sensitive SK2 current was well preserved

in frog saccular hair cells isolated by collagenase IV or trypsin, but this current in trypsinized cells could be totally abolished after brief exposure to protease VIII, protease XXIV or papain [16]. Moreover, Armstrong *et al.* had also showed that the electrical properties of frog saccular hair cells could be distorted by papain dissociation resulting in the absence of SK2 channels and the emergence of non-inactivating BK channels [22,23]. These results indicated that BK channel and SK2 channel may co-exist in the same saccular hair cells of frog.

Based on the above findings, we propose a hypothesis that m2mAChR and $\alpha 9$ nAChR may co-exist in the same VHCs II and mediate distinct channels respectively, while these differential cholinergic responses depend on the isolating preparation with different enzymes.

Therefore, the aim of the present study was to explore the correlation between m2mAChR mediated BK currents and $\alpha 9$ nAChR mediated SK2 currents in the same VHCs II of guinea pig. First, under the consistent experimental condition, we should demonstrate that these two cholinergic receptors $\alpha 9$ nAChR and m2mAChR mediated two distinct channels in VHCs II of guinea pig isolated with trypsin or collagenase IA respectively. Then, we should further prove that $\alpha 9$ nAChR and m2mAChR may co-exist in the same cell. Therefore, we designed a three-step experiment by whole cell patch-clamp recordings. Consequently we presented our results in three parts: (1) VHCs II isolated by collagenase IA routinely displayed the m2mAChR sensitive BK current; (2) VHCs II isolated by trypsin routinely displayed the $\alpha 9$ nAChR sensitive SK2 current; (3) After brief exposing the patched cells isolated by trypsin to collagenase IA for 3 min, the $\alpha 9$ nAChR sensitive SK2 current was abolished, while a non-inactivating BK current was activated by ACh. These findings provided evidences that the two distinct channels (BK and SK2) as well as the two ACh receptors ($\alpha 9$ nAChR and m2mAChR) may co-exist in the same mammalian VHCs II, and the alternative presence of these two ACh receptors sensitive currents was dependent on the isolating preparation with different enzymes.


2. Results and Discussion

2.1. m2mAChR Sensitive BK Currents Exist in VHCs II Isolated by Collagenase IA

Our previous finding showed that the BK currents activated by ACh in VHCs II isolated by collagenase IA were mediated by m2mAChR [19], which could be potently inhibited by the BK channel blocker IBTX [17,19] and m2mAChR antagonist methoctramine [19], but was insensitive to the SK2 channel blocker apamin [17] and $\alpha 9$ nAChR antagonist strychnine [19]. In order to further confirm these results in the same cell under the consistent experimental condition, we again demonstrated the properties of currents activated by ACh in the same VHCs II isolated by collagenase IA. As shown in Figure 1, the outward currents activated by ACh (100 μ M) were strongly inhibited by 200 nM IBTX and 100 nM methoctramine to $20.3\% \pm 7.1\%$ (control, 206.4 ± 55.2 pA; ACh + IBTX, 44 ± 24.5 pA; $p < 0.001$; $n = 6$) and $28.6 \pm 6.8\%$ (control, 185.2 ± 59 pA; ACh + methoctramine, 55 ± 26.1 pA; $p = 0.002$; $n = 6$) respectively, but insensitive to 200 nM strychnine (control, 207.6 ± 46.3 pA, strychnine + ACh, 206.8 ± 47.8 pA, $p = 0.981$; $n = 6$) and 300 nM apamin (control, 218.3 ± 66.5 pA, ACh + apamin, 213 ± 65.4 pA, $p = 0.914$; $n = 6$). Therefore, these results showed that ACh activated a non-inactivating outward current in VHCs II isolated by collagenase IA, which was

BK current and mediated by m2mAChR. As described in our previous findings [17,19], the SK2 channels and the $\alpha 9$ nAChR were not activated at the same time.

Figure 1. ACh-activated BK currents in collagenase IA isolated VHCs II. ACh routinely activated a non-inactivating outward current. (A) The current activated by 100 μ M ACh was insensitive to strychnine and apamin, respectively; (B) The current activated by 100 μ M ACh was potently inhibited by methoctramine and IBTX, respectively. All of these four results were obtained from the one same cell; (C) Bar histogram showed the effects of 200 nM strychnine, 300 nM apamin, 100 nM methoctramine and 200 nM IBTX on the current evoked by 100 μ M ACh. Each point represents the mean \pm SD of 5–6 cells. ($V_{\text{hold}} = -50$ mV).




2.2. $\alpha 9$ nAChR Sensitive SK2 Currents Exist in VHCs II Isolated by Trypsin

It has been reported that $\alpha 9$ nAChR sensitive SK2 currents were well preserved in frog saccular hair cells isolated by trypsin [16], which were sensitive to apamin and strychnine [14–16]. Here, we studied the properties of currents activated by ACh in VHCs II isolated by trypsin. As shown in Figure 2, the outward currents activated by 100 μ M ACh in one same cell were potently inhibited to $37.6\% \pm 10.5\%$ and $28.4\% \pm 10.6\%$ by 200 nM strychnine (control, 145.3 ± 6.2 pA; ACh + strychnine, 54.8 ± 16.5 pA; $p < 0.001$; $n = 6$) and 300 nM apamin (control, 129.8 ± 10.5 pA; ACh + apamin, 36.8 ± 13.4 pA; $p < 0.001$; $n = 6$), respectively, but insensitive to 100 nM methoctramine (control, 141.3 ± 20 pA; ACh + methoctramine, 140.3 ± 19.5 pA; $p = 0.945$; $n = 6$) and 200 nM IBTX (control, 140 ± 14.7 pA; ACh + IBTX, 137.5 ± 10.1 pA; $p = 0.789$; $n = 6$). Therefore, these results showed that ACh activated a

fast inactivating current, which was SK2 current and mediated by the $\alpha 9$ nAChR in VHCs II isolated by trypsin. Meanwhile, the BK channels and the m2mAChR were not activated at the same time.

Figure 2. ACh activated SK2 currents in trypsin-isolated VHCs II. ACh routinely activated a fast inactivating outward current. **(A)** The current activated by 100 μ M ACh was potently inhibited by strychnine and apamin, respectively; **(B)** The current activated by 100 μ M ACh was insensitive to methoctramine and IBTX, respectively. All of these four results were obtained from the one same cell; **(C)** Bar histogram showed the inhibition effects of 200 nM strychnine, 300 nM apamin, 100 nM methoctramine and 200 nM IBTX on the current evoked by 100 μ M ACh. Each point represents the mean \pm SD of 6 cells. ($V_{\text{hold}} = -50$ mV).



2.3. m2mAChR Sensitive BK Currents and $\alpha 9$ nAChR Sensitive SK2 Currents Co-Exist in VHCs II

The above results indicated that there presented two distinct ACh receptors activated currents in VHCs II isolated by different enzymes. We noted that certain enzymes could affect or even damage the physiological properties of hair cells [16,22,23]. Therefore, we investigated the proteolytic effect of collagenase IA on the trypsin-dissociated cells to verify the hypothesis that two distinct channels mediated by m2mAChR and $\alpha 9$ nAChR co-exist in VHCs II.

As illustrated in Figure 3A, the fast inactivating currents activated by 100 μ M ACh in one same trypsinized cell were dramatically reduced to $29.9\% \pm 5.8\%$ by 200 nM strychnine (control,

196 ± 53 pA; ACh + strychnine, 53.6 ± 9.6 pA; $p = 0.006$; $n = 6$), while insensitive to 100 nM methoctramine (control, 196 ± 53 pA; ACh + methoctramine, 194.8 ± 63.4 pA; $p = 0.936$; $n = 6$). Next, 0.8 mg/mL collagenase IA dissociation solution was applied to this patched cell for 3 min. There was a downward shift in the baseline (Figure 3B). Then, this patched cell was incubated with the normal external solution for 5 min. Finally, 100 μ M ACh was applied again. Interestingly, re-application of ACh activated a non-inactivating current, which was different from the previously recorded fast inactivating SK2 current. As shown in Figure 3C, this non-inactivating current was insensitive to 200 nM strychnine (control, 142.2 ± 36.4 pA; ACh + strychnine, 135.2 ± 37.5 pA; $p = 0.883$; $n = 6$), but could be potently inhibited by 100 nM methoctramine to $22.9\% \pm 6.7\%$ (control, 142.2 ± 38.7 pA; ACh + methoctramine, 32 ± 12.5 pA; $p = 0.002$; $n = 6$). Similar results were obtained from six cells. These results not only showed that the $\alpha 9$ nAChR responses were replaced by the m2mAChR responses after brief exposure to collagenase IA, but also suggested that these two distinct channels and their corresponding ACh receptors may co-exist in the same VHC II.

2.4. Discussion


In the present study, we verified the hypothesis that m2mAChR, $\alpha 9$ nAChR and the two distinct channels (BK and SK2) may co-exist in the same VHCs II of guinea pig. We showed that m2mAChR sensitive BK currents were located in VHCs II isolated by collagenase IA, and $\alpha 9$ nAChR sensitive SK2 currents were located in VHCs II isolated by trypsin. After brief exposing the patched cells isolated by trypsin to collagenase IA, $\alpha 9$ nAChR sensitive SK2 currents were abolished, while m2mAChR sensitive BK currents were activated. These results suggested that these two distinct channels and their respective ACh receptors may co-exist in the same VHCs II of guinea pig, and the alternative presence of these two ACh receptors-sensitive currents was dependent on the isolating preparation with different enzymes.

2.4.1. BK Channel and SK2 Channel Co-Exist in the Same VHCs II, but do Not Play Their Roles at the Same Time in Vestibular Efferent Inhibition as in OHCs

The inner ear comprises two main functional parts, cochlea and vestibule, which are responsible for sound perception and balance maintenance respectively. It has been established that ACh is the main efferent neurotransmitter in mammalian inner ear [6,24], which activates $\alpha 9$ nAChR with the subsequent opening of SK2 channel [14–16]. In the mammalian vestibular neuroepithelium, VHCs II are very small and rare sensory cells, with which the patch clamp experiments are very hard to manipulate, so the detailed mechanisms underlying cholinergic efferent inhibition in vestibule remain unclear, except for the well known $\alpha 9$ nAChR mediated SK2 channel [14–16]. However, there has been morphological evidence recently showing that another kind of calcium-activated potassium channel named BK channels are expressed in VHCs II of rat by immunohistochemical study [25]. Besides, our previous studies have also demonstrated that ACh activated BK currents are present in the VHCs II of guinea pig [17,18]. Further studies showed that this inhibition is mediated by m2mAChR [19]. In the present study, we again demonstrated that m2mAChR mediated BK channels in VHCs II isolated by collagenase IA, which was not activated by $\alpha 9$ nAChR (Figure 1). All these results indicate that BK channel is also involved in cholinergic efferent inhibition in vestibule. Moreover, after brief exposing

the patched cell isolated by trypsin to collagenase IA, $\alpha 9$ nAChR sensitive SK2 currents were abolished, while m2mAChR sensitive BK currents were activated (Figure 3). These results imply that BK channel and SK2 channel may co-exist in the same VHCs II, so do their respective upstream receptors $\alpha 9$ nAChR and m2mAChR.

Figure 3. Change of the currents after exposure to collagenase IA on one trypsin-isolated cell. **(A)** Before exposure to collagenase IA, the trypsin-isolating cell displayed a typical fast inactivating current, which was strongly inhibited by 200 nM strychnine, but insensitive to 100 nM methoctramine; **(B)** 0.8 mg/mL collagenase IA dissociation solution was applied to the patched cell for 3 min. Note that there was a downward shift in the baseline; **(C)** After exposure to collagenase IA and incubated with normal external solution for 5 min, the fast inactivating current was totally abolished, while a non-inactivating current was activated. The non-inactivating current was insensitive to 200 nM strychnine, but could be potentially inhibited by 100 nM methoctramine. All of these above results were obtained from one same cell; **(D)** Bar histogram showed the inhibition effects of 200 nM strychnine and 100 nM methoctramine on the current evoked by 100 μ M ACh. Each point represents the mean \pm SD of 6 cells. ($V_{\text{hold}} = -50$ mV).



Recently, in the OHC area of the adult rat cochlea, immunohistochemical studies have shown that BK-labeled plaques are observed beneath the nuclei of the OHCs except at the most apical cochlear region [26,27], and are located at the interface between presynaptic olivocochlear efferent terminals and the post-synaptic OHC membrane [26,28,29]. Since SK2 channel is almost expressed in all the apical, middle, and basal OHCs, double labeling experiments with antibodies against SK2 channel

and BK channel demonstrate that SK2 channel and BK channel co-exist in the middle and basal OHCs [26,27]. By whole cell patch clamp recordings, it has been reported that BK channels play important roles in cholinergic inhibition of high frequency cochlear OHCs [26]. Then, after electrically stimulating the olivocochlear bundle in the BK^{-/-} ears, the OC-mediated suppression is reduced, but not eliminated at all frequencies. The remaining suppression is blocked by strychnine, suggesting involvement of $\alpha 9/\alpha 10$ cholinergic receptors, which are coupled to activation of the remaining SK2 channels. These results indicate that the efferent inhibition in the OHCs is combined action of BK and SK2 channels throughout the cochlea [27], but the subtype of cholinergic receptors mediating BK channels in OHCs is subtler.

According to the results that the efferent inhibition in the OHCs is combined action of BK and SK2 channels throughout the cochlea, it is reasonable to suppose that this efferent inhibition in the vestibule may involve both BK and SK2 channels. In the present study, we showed that VHCs II isolated by trypsin routinely displayed the $\alpha 9$ nAChR sensitive SK2 current (Figure 2), but was insensitive to IBTX, a specific blocker of BK channels, while VHCs II isolated by collagenase IA routinely displayed the m2mAChR sensitive BK current (Figure 1), but was insensitive to apamin, a specific blocker of SK2 channels. According to these results, although the co-existence of BK and SK2 channels in VHCs II, we speculate that the efferent inhibition in the VHCs II might not be the combined action of these two channels. The reason may be as follow: we demonstrated that ACh activated BK currents were mediated by m2mAChR [19], while a recent study showed that m2mAChR was only present in olivocochlear neurons in cochlea, including their axons in the nerve trunk and the intraganglionic bundle and the osseous spiral lamina, as well as their terminals in the inner spiral bundle and underneath each of the three outer hair cell rows by immunohistochemistry. Besides, single cell RT-PCR also showed that m2mAChR was not present in OHCs [30]. ‘Knockout’ mice missing either $\alpha 9$ or $\alpha 10$ nAChRs failed to show suppression of distortion product otoacoustic emissions (DPOAEs) during efferent fiber stimulation [31,32] and their hair cells no longer responded to local application of ACh [32]. Therefore, these results indicated that the subtypes of AChR mediating BK channel in cochlea may be $\alpha 9/\alpha 10$ nAChR, which also mediated SK2 channel at the same time in OHCs. Nevertheless, our studies showed that BK channel mediated cholinergic inhibition in VHCs II by m2mAChR, which stimulated G $\beta\gamma$ -mediated excitation of AC/cAMP activities and led to the phosphorylation of Ca²⁺ channels, resulting in the influx of Ca²⁺ and opening of the BK channel [19]. Based on these kinetics features, due to the involvement of second messenger, the signal transduction of m2mAChR mediated BK channel would be activated slower than that of $\alpha 9$ nAChR mediated SK2 channel. In spite of the coexistence of these two receptors on the VHCs II, ACh would more quickly activate the $\alpha 9$ nAChR, which mediated a fast outward current SK2 and hyperpolarized membrane potential, thereby masking m2mAChR response in VHCs II.

2.4.2. The Mechanisms Underlying the Changes of the ACh Response in Trypsined Cells after Exposure to Collagenase IA

In order to isolate individual vestibular hair cells from mammalian vestibular epithelium, the enzymes are commonly used, including papain and several serine proteases [33–36]. It has been reported that some collagenases have been successfully used to isolate cells from the cochlea [37–40],

but they are rather ineffective in isolating mammalian VHCs II [16]. Nevertheless, our previous studies showed that ACh activated m2AChR mediated BK currents by using collagenase IA to isolate VHCs II, yet previous studies routinely demonstrated that ACh activated $\alpha 9$ nAChR mediated SK2 currents by using trypsin to isolate mammalian VHCs II. This failure to demonstrate routine ACh responses led to speculation that the $\alpha 9$ nAChR might undergo proteolysis during the collagenase IA isolation process and become non-functional. Indeed, there are several instances in the literature where enzymes have been shown to alter physiological responses of $\alpha 9$ nAChR/SK2 complex. Armstrong and Roberts [22] have recently showed that papain altered particular ion channels in frog saccular hair cells including SK2, and Holt JC [16] has also recently showed that perfusions with either protease or papain permanently abolished the $\alpha 9$ -nicotinic response in isolated saccular hair cells. According to these previous studies about the proteolytic effects on $\alpha 9$ nAChR/SK complex, we speculate that it maybe collagenase IA that permanently abolished the $\alpha 9$ -nicotinic response in trypsin isolated VHCs II. Therefore, we would record m2AChR mediated BK currents in collagenase IA isolated VHCs II.

Meanwhile, the possible mechanism of proteolysis of $\alpha 9$ nAChR/SK2 complex by collagenase IA but not trypsin may be as follow:

Collagenase IA, a crude enzyme containing 10 to 18 enzymes, including two specific collagenases, clostripain, trypsin and a neutral protease caseinase (Sigma Product information), and most of them are proteases. it has been showed that perfusions with protease could permanently abolished the $\alpha 9$ nAChR response in isolated saccular hair cells, while these inactivated enzymes became completely ineffective at abolishing the $\alpha 9$ nAChR response [16]. Therefore, we speculated that the protease of clostripain and caseinase contained in crude collagenase IA may exert the proteolytic effect on the $\alpha 9$ nAChR/SK2 complex as a result that clostripain specifically cleaves protein on the C-terminal of arginine residues [41] and caseinase has a broad range of amino acid specificity cleavage points, acting at the neutral pH.

However, trypsin is a polypeptide containing approximately 220 amino acid residues with 6 disulfide bridges and 1 calcium ion, which highly cleaves on the C-terminal side of lysine and arginine [42]. It has been reported that the voltage-sensitive Na^+ , K^+ and Ca^{2+} channels which have been shown to contain trypsin-sensitive inhibitory domains share some common features: (1) large subunit molecular weights; (2) organization into domains containing six membrane spanning helices and a number of conserved motifs [43]. Besides, SK2 channel seems to belong to the trypsin-sensitive ion channel family due to the features as below [44]: (1) it contains 553–580 KD amino acids and 6 putative transmembrane domains; (2) there are five possible hydrolysis sites along its four extracellular loops. However, the arginine-glutamate bond in the third loop of SK2 channel should slow or impair the proteolytic effect since it has been reported that the neighboring acid residues such as aspartate and glutamate can impair the proteolytic effect of trypsin [45]. Therefore, in the present study, we showed that trypsin has little damaging effect on the function of $\alpha 9$ nAChR in vestibular hair cells, which is in consistent with previous study [16].

Based on these reports, we speculated that the function of $\alpha 9$ nAChR/SK complex may be irreversibly abolished after brief exposure to collagenase IA. Whether the lost activation of the SK2 current is a result of the proteolysis of SK2 alone or both $\alpha 9$ nAChR and SK2 should be further investigated.

Meanwhile, as shown in Figure 3B, after exposing to collagenase IA, there was a downward shift in the baseline. It is well known that the resting membrane is mainly permeable to K^+ ; thus, K^+ channels

play an important role in regulating the resting membrane potential, including Kir channel [46] and K_V channel [47]. Armstrong *et al.* showed that Kir channel [48,49] and K_V channel [22] were altered by enzymatic dissociation and K_V current [22] was apparently eliminated. Therefore, the downward shift in the baseline observed after the application of collagenase IA might be due to alteration of K^+ channel by collagenase IA in VHCs II, which finally affects the resting membrane potential of the VHCs II.

3. Experimental Section

3.1. Animal Procedures and VHCs II Preparation

The Animal Care and Use Committee of Tongji Medical College approved these animal experiments. Briefly, 75 young guinea pigs (weighing 250–300 g, 6–10 weeks old) were deeply anesthetized by intramuscular injection with 0.3 mL of a mix of $\frac{1}{3}$ xylazine (2%, Rompum, Bayer, German) and $\frac{2}{3}$ ketamine hydrochlorate (50 mg/mL, Ketalar, Parke-Davis, France) and decapitated. Then, two methods of hair cells preparation involving different enzymes were applied independently.

3.2. VHCs II Isolated by Collagenase IA

Following decapitation, the vestibular epithelium was removed and incubated for 3 min at room temperature (20–24 °C) with 0.8 mg/mL collagenase IA in a low Ca^{2+} and Mg^{2+} -free balanced salt solution (137 mM NaCl, 5.4 mM KCl, 0.1 mM $CaCl_2$, 0.2 mM Na_2HPO_4 , 0.4 mM KH_2PO_4 , 10 mM glucose; pH = 7.2). Washing with normal external solution containing 2 mM $CaCl_2$ stopped the enzymatic action. Finally, hair cells were mechanically dissociated by gentle trituration and were left to settle on the bottom of the experimental chamber coated with rat tail collagen.

3.3. VHCs II Isolated by Trypsin

Following decapitation, the vestibular epithelium was removed and incubated for 3 min at room temperature (20–24 °C) with 0.8 mg/mL trypsin in normal external solution as described below. Next, the normal external solution containing trypsin inhibitor was used to stop the enzymatic action. Finally, hair cells were mechanically dissociated by gentle trituration and were left to settle on the bottom of the experimental chamber coated with rat tail collagen.

3.4. Electrophysiology

VHCs II were readily identified by the cylindrical shape and absence of a distinct neck region [1]. The current activated by ACh was recorded in a whole-cell configuration, using an Axon-200B patch clamp amplifier (Axon Instruments, Foster City, CA, USA). Patch electrodes were pulled from borosilicate glass capillaries, using a Model P-97 electrode puller (Sutter Instrument, Novato, CA, USA). Electrode resistances ranged from 3 to 6 M Ω when filled with the internal solution, as described below. Records were low-pass filtered at 5 kHz with a four-pole Bessel filter. After gigaseal formation onto the basolateral membrane of VHCs and membrane disruption, the membrane capacitance was 4.6 ± 1.3 pF on average (C_m , $n = 6$), and series resistance (R_s , 6–15 M Ω) was compensated for up to 80%. The drug was applied to a patched cell at a holding potential of –50 mV.

The external solution component was as follow: 150 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 1 mM MgCl₂, 5 mM glucose, and 10 mM HEPES (pH = 7.2). The internal solution component was as follow: 150 mM KCl, 2 mM MgCl₂, 0.1 mM CaCl₂, 3 mM Na₂ATP, 5 mM EDTA and 10 mM HEPES (pH = 7.2).

The test solutions were applied to cells by a gravity-delivered linear barrel microperfusion system described elsewhere [18]. The microperfusion system consisted of a series of fused silica tubes (eight tubes with outer diameter of 150 µm and internal diameter of 100 µm) connected to a series of independent reservoirs. The tip of the tube was positioned approximately 100 to 150 µm from the cells. This microperfusion system was manipulated by shifting the tubes horizontally with a Leitz micromanipulator (ACS01, Leitz Corp., Wetzlar, Germany).

3.5. Drugs

All drugs were purchased from Sigma (St. Louis, MO, USA). Trypsin inhibitor (T9128), ACh (A6625), apamin (A9459), iberiotoxin (IBTX, I5904), methoctramine (M105) and strychnine (S7001) were directly dissolved in the external solution.

3.6. Data Analysis

Data were analyzed and plotted using the pCLAMP8.1 Clampfit 8.1 software (Axon Instruments, Foster City, CA, USA) and SigmaPlot 9.0 (Systat Software, Richmond, CA, USA). Results are presented as the mean ± SD, and statistical significance was determined using the t test. Differences were considered to be significant if $p < 0.05$; all differences listed were statistically significant, unless stated otherwise.

4. Conclusions

In conclusion, our findings provide evidences that BK channel and SK channel, as well as their corresponding m2mAChR and α9nAChR may co-exist in the same VHCs II of guinea pig, but the efferent inhibition in the VHCs II is not combined action of BK and SK2 channels as in cochlea. The physiological significance of m2mAChR mediated a slower BK current in VHCs II and the co-existence of these two ACh receptors sensitive currents in the same mammalian VHCs II should be further elaborated. Moreover, the alternative presence of these two ACh receptors sensitive currents is dependent on the isolating preparation with different enzymes. The proteolytic mechanisms of different enzymes on the α9nAChR/SK2 complex need further elucidation.

Acknowledgments

This work was supported by grants from the Major State Basic Research Development Program of China (973 program) (No. 2011CB504504), the National Science and Technology Pillar Program during the Twelfth Five-year Plan Period (No. 2012BAI12B02) and National Nature Science Foundation of China (Nos. 81230021 and 81000409).

Conflict of Interest

The authors declare no conflict of interest.

References

1. Scarfone, E.; Ulfendahl, M.; Lofstrand, P.; Flock, A. Light- and electron microscopy of isolated vestibular hair cells from the guinea pig. *Cell Tissue Res.* **1991**, *266*, 51–58.
2. Godfrey, D.A.; Park, J.L.; Ross, C.D. Choline acetyltransferase and acetylcholinesterase in centrifugal labyrinthine bundles of rats. *Hear. Res.* **1984**, *14*, 93–106.
3. Kong, W.J.; Egg, G.; Hussl, B.; Spoendlin, H.; Schrott-Fischer, A. Localization of chat-like immunoreactivity in the vestibular endorgans of the rat. *Hear. Res.* **1994**, *75*, 191–200.
4. Kong, W.J.; Hussl, B.; Thumfart, W.F.; Schrott-Fischer, A. Ultrastructural localization of ChAT-like immunoreactivity in the human vestibular periphery. *Hear. Res.* **1998**, *119*, 96–103.
5. Kong, W.J.; Scholtz, A.W.; Hussl, B.; Kammen-Jolly, K.; Schrott-Fischer, A. Localization of efferent neurotransmitters in the inner ear of the homozygous Bronx waltzer mutant mouse. *Hear. Res.* **2002**, *167*, 136–155.
6. Guth, P.S.; Perin, P.; Norris, C.H.; Valli, P. The vestibular hair cells: Post-transductional signal processing. *Prog. Neurobiol.* **1998**, *54*, 193–247.
7. Caulfield, M.P. Muscarinic receptors—Characterization, coupling and function. *Pharmacol. Ther.* **1993**, *58*, 319–379.
8. Wess, J. Molecular biology of muscarinic acetylcholine receptors. *Crit. Rev. Neurobiol.* **1996**, *10*, 69–99.
9. Anderson, A.D.; Troyanovskaya, M.; Wackym, P.A. Differential expression of alpha2–7, alpha9 and beta2–4 nicotinic acetylcholine receptor subunit mRNA in the vestibular end-organs and Scarpa's ganglia of the rat. *Brain. Res.* **1997**, *778*, 409–413.
10. Wackym, P.A.; Chen, C.T.; Ishiyama, A.; Pettis, R.M.; Lopez, I.A.; Hoffman, L. Muscarinic acetylcholine receptor subtype mRNAs in the human and rat vestibular periphery. *Cell Biol. Int.* **1996**, *2*, 187–192.
11. Ishiyama, A.; Lopez, I.; Wackym, P.A. Molecular characterization of muscarinic receptors in the human vestibular periphery. Implications for pharmacotherapy. *Am. J. Otol.* **1997**, *18*, 648–654.
12. Li, G.Q.; Kevetter, G.A.; Leonard, R.B.; Prusak, D.J.; Wood, T.G.; Correia, M.J. Muscarinic acetylcholine receptor subtype expression in avian vestibular hair cells, nerve terminals and ganglion cells. *Neuroscience* **2007**, *146*, 384–402.
13. Yao, Q.; Cheng, H.; Guo, C.; Zhou, T.; Huang, X.; Kong, W. Muscarinic acetylcholine receptor subtype expression in type vestibular hair cells of guinea pigs. *J. Huazhong. Uni. Sci. Technol.* **2011**, *31*, 682–686.
14. Yoshida, N.; Shigemoto, T.; Sugai, T.; Ohmori, H. The role of inositol trisphosphate on ACh-induced outward currents in bullfrog saccular hair cells. *Brain. Res.* **1994**, *644*, 90–100.
15. Elgoyhen, A.B.; Vetter, D.E.; Katz, E.; Rothlin, C.V.; Heinemann, S.F.; Boulter, J. alpha10: A determinant of nicotinic cholinergic receptor function in mammalian vestibular and cochlear mechanosensory hair cells. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 3501–3506.
16. Holt, J.C.; Lioudyno, M.; Athas, G.; Garcia, M.M.; Perin, P.; Guth, P.S. The effect of proteolytic enzymes on the alpha9-nicotinic receptor-mediated response in isolated frog vestibular hair cells. *Hear. Res.* **2001**, *152*, 25–42.

17. Kong, W.J.; Guo, C.K.; Zhang, S.; Hao, J.; Wang, Y.J.; Li, Z.W. The properties of ACh-induced BK currents in guinea pig type II vestibular hair cells. *Hear. Res.* **2005**, *209*, 1–9.
18. Kong, W.J.; Guo, C.K.; Zhang, X.W.; Chen, X.; Zhang, S.; Li, G.Q.; Li, Z.W.; van Cauwenberge, P. The coupling of acetylcholine-induced BK channel and calcium channel in guinea pig saccular type II vestibular hair cells. *Brain. Res.* **2007**, *1129*, 110–115.
19. Guo, C.K.; Wang, Y.; Zhou, T.; Yu, H.; Zhang, W.J.; Kong, W.J. M2 muscarinic ACh receptors sensitive BK channels mediate cholinergic inhibition of type II vestibular hair cells. *Hear. Res.* **2012**, *285*, 13–19.
20. Kong, W.J.; Guo, C.K.; Zhang, S.; Zhang, X.W.; Wang, Y.J.; Li, Z.W. Fast cholinergic efferent inhibition in guinea pig outer hair cells. *Brain. Res.* **2006**, *1102*, 103–108.
21. Darbon, P.; Wright, D.J.; Evans, M.G. Conductance properties of the acetylcholine receptor current of Guinea pig outer hair cells. *J. Assoc. Res. Otolaryngol.* **2011**, *12*, 59–70.
22. Armstrong, C.E.; Roberts, W.M. Electrical properties of frog saccular hair cells: Distortion by enzymatic dissociation. *J. Neurosci.* **1998**, *18*, 2962–2973.
23. Armstrong, C.E.; Roberts, W.M. Rapidly inactivating and non-inactivating calcium-activated potassium currents in frog saccular hair cells. *J. Physiol.* **2001**, *536*, 49–65.
24. Eybalin, M. Neurotransmitters and neuromodulators of the mammalian cochlea. *Physiol. Rev.* **1993**, *73*, 309–373.
25. Schweizer, F.E.; Savin, D.; Luu, C.; Sulstemeier, D.R.; Hoffman, L.F. Distribution of high-conductance calcium-activated potassium channels in rat vestibular epithelia. *J. Comp. Neuro.* **2009**, *517*, 134–145.
26. Wersinger, E.; McLean, W.J.; Fuchs, P.A.; Pyott, S.J. BK channels mediate cholinergic inhibition of high frequency cochlear hair cells. *PLoS One* **2010**, *5*, e13836.
27. Maison, S.F.; Pyott, S.J.; Meredith, A.L.; Liberman, M.C. Olivocochlear suppression of outer hair cells *in vivo*: Evidence for combined action of BK and SK2 channels throughout the cochlea. *J. Neurophysiol.* **2013**, doi:10.1152/jn.00924.2012.
28. Engel, J.; Braig, C.; Rüttiger, L.; Kuhn, S.; Zimmermann, U.; Blin, N.; Sausbier, M.; Kalbacher, H.; Munkner, S.; Rohbock, K.; *et al.* Two classes of outer hair cells along the tonotopic axis of the cochlea. *Neuroscience* **2006**, *143*, 837–849.
29. Hafidi, A.; Beurg, M.; Dulon, D. Localization and developmental expression of BK channels in mammalian cochlear hair cells. *Neuroscience* **2005**, *130*, 475–484.
30. Maison, S.F.; Liu, X.P.; Vetter, D.E.; Eatock, R.A.; Nathanson, N.M.; Wess, J.; Liberman, M.C. Muscarinic signaling in the cochlea: presynaptic and postsynaptic effects on efferent feedback and afferent excitability. *J. Neurosci.* **2010**, *30*, 6751–6762.
31. Vetter, D.E.; Liberman, M.C.; Mann, J.; Barhanin, J.; Boulter, J.; Brown, M.C.; Saffioti-Kolman, J.; Heinemann, S.F.; Elgoyhen, A.B. Role of $\alpha 9$ nicotinic ACh receptor subunits in the development and function of cochlear efferent innervation. *Neuron* **1999**, *23*, 93–103.
32. Vetter, D.E.; Katz, E.; Maison, S.F.; Taranda, J.; Turcan, S.; Ballesteros, J.; Liberman, M.C.; Elgoyhen, A.B.; Boulter, J. The $\alpha 10$ nicotinic acetylcholine receptor subunit is required for normal synaptic function and integrity of the olivocochlear system. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 20594–20599.

33. Lewis, R.S.; Hudspeth, A.J. Voltage- and ion-dependent conductances in solitary vertebrate hair cells. *Nature* **1983**, *304*, 538–541.
34. Steinacker, A.; Rojas, L. Acetylcholine modulated potassium channel in the hair cell of the toadfish saccule. *Hear. Res.* **1988**, *35*, 265–269.
35. Housley, G.D.; Norris, C.H.; Guth, P.S. Cholinergically-induced changes in outward currents in hair cells isolated from the semicircular canal of the frog. *Hear. Res.* **1990**, *43*, 121–133.
36. Holt, J.R.; Eatock, R.A. Inwardly rectifying currents of saccular hair cells from the leopard frog. *Neurophysiology* **1995**, *73*, 1484–1502.
37. ErosteGUI, C.; Norris, C.H.; Bobbin, R.P. *In vitro* pharmacologic characterization of a cholinergic receptor on outer hair cells. *Hear. Res.* **1994**, *74*, 135–147.
38. Nenov, A.P.; Norris, C.; Bobbin, R.P. Acetylcholine response in guinea pig outer hair cells. I. Properties of the response. *Hear. Res.* **1996**, *101*, 132–148.
39. Sugawara, M.; ErosteGUI, C.; Blanchet, C.; Dulon, D. ATP activates a cation conductance and Ca^{2+} -dependent Cl^- conductance in Hensen cells of guinea pig cochlea. *Am. J. Physiol.* **1996**, *271*, C1817–1827.
40. Nenov, A.P.; Chen, C.; Bobbin, R.P. Outward rectifying potassium currents are the dominant voltage activated currents present in Deiters' cells. *Hear. Res.* **1998**, *123*, 168–182.
41. Gille, A.M.; Imhoff, J.M.; Keil, B. á-Clostripain. *J. Biol. Chem.* **1979**, *10*, 1462–1468.
42. Hermodson, M.A.; Ericsson, L.H.; Neurath, H.; Walsh, K.A. Determination of the amino acid sequence of porcine trypsin by sequenator analysis. *Biochemistry* **1973**, *12*, 3146–3153.
43. Catterall, W.A. Structure and function of voltage-sensitive ion channels. *Science* **1988**, *242*, 50–61.
44. Desai, R.; Peretz, A.; Idelson, H.; Lazarovici, P.; Attali, B. Ca^{2+} -activated K^+ channels in human leukemic Jurkat T cells. Molecular cloning, biochemical and functional characterization. *J. Biol. Chem.* **2000**, *275*, 39954–39963.
45. Smyth, D.G. Techniques in enzymic hydrolysis. *Methods Enzymol.* **1967**, *11*, 214–231.
46. Chilton, L.; Ohya, S.; Freed, D.; George, E.; Drohic, V.; Shibukawa, Y.; Maccannell, K.A.; Imaizumi, Y.; Clark, R.B.; Dixon, I.M.; *et al.* K^+ currents regulate the resting membrane potential, proliferation, and contractile responses in ventricular fibroblasts and myofibroblasts. *Am. J. Physiol. Heart. Circ. Physiol.* **2005**, *288*, H2931–H2939.
47. Yuan, X.J. Voltage-gated K^+ currents regulate resting membrane potential and $[\text{Ca}^{2+}]_i$ in pulmonary arterial myocytes. *Cir. Res.* **1995**, *77*, 370–378.
48. Armstrong, C.E.; Roberts, W.M. Papain alters the resonant frequency of frog saccular hair cells. *Biophys. J. Abstr.* **1996**, *70*, A346.
49. Armstrong, C.E.; Roberts, W.M. At least three K^+ currents are involved in resonance in frog saccular hair cells. *Biophys. J. Abstr.* **1997**, *72*, A355.