

Supplementary Information

Table S1. Top 10 Differentially expressed genes by microarray.

| Gene symbol | Gene name | Expression value | p value |
|-----------------|---------------------------------------|------------------|-------------------|
| <i>IFI27</i> | Interferon alpha-inducible protein 27 | 0.774 | <i>p</i> < 0.0001 |
| <i>CA1</i> | Carbonic anhydrase 1 | 0.705 | <i>p</i> < 0.0001 |
| <i>HBD</i> | Hemoglobin subunit delta | 0.640 | <i>p</i> < 0.0001 |
| <i>XK</i> | X-linked Kx blood group | 0.534 | <i>p</i> < 0.0005 |
| <i>RHCE/RHD</i> | Blood group Rh(CE) polypeptide | 0.507 | <i>p</i> < 0.0005 |
| <i>MS4A1</i> | B-lymphocyte antigen CD20 | -0.821 | <i>p</i> < 0.0001 |
| <i>IGHM</i> | Ig mu chain C region | -0.816 | <i>p</i> < 0.0001 |
| <i>PAX5</i> | Paired box protein Pax-5 | -0.791 | <i>p</i> < 0.0001 |
| <i>FCRLA</i> | Fc receptor-like A | -0.669 | <i>p</i> < 0.0001 |
| <i>TTC3</i> | Tetratricopeptide repeat protein 3 | -0.647 | <i>p</i> < 0.0005 |

© 2013 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).