

Article

Design of Phosphonium-Type Zwitterion as an Additive to Improve Saturated Water Content of Phase-Separated Ionic Liquid from Aqueous Phase toward Reversible Extraction of Proteins

Yoritsugu Ito^{1,2}, Yuki Kohno^{1,2}, Nobuhumi Nakamura^{1,2,3} and Hiroyuki Ohno^{1,2,3,*}

¹ Department of Biotechnology, Tokyo University of Agriculture and Technology, 2-24-16, Naka-cho, Koganei, Tokyo 184-8588, Japan; E-Mails: 50011641201@st.tuat.ac.jp (Y.I.); y-kohno@cc.tuat.ac.jp (Y.K.); nobu1@cc.tuat.ac.jp (N.N.)

² Functional Ionic Liquid Laboratories, Graduate School of Engineering, Tokyo University of Agriculture and Technology, 2-24-16, Naka-cho, Koganei, Tokyo 184-8588, Japan

³ Japan Science and technology Agency (JST), Core Research for Evolutional Science and Technology (CREST), Chiyoda, Tokyo 102-0076, Japan

* Author to whom correspondence should be addressed; E-Mail: ohnoh@cc.tuat.ac.jp; Tel./Fax: +81-42-388-7024.

Received: 8 August 2013; in revised form: 30 August 2013 / Accepted: 2 September 2013 /

Published: 5 September 2013

Abstract: We designed phosphonium-type zwitterion (ZI) to control the saturated water content of separated ionic liquid (IL) phase in the hydrophobic IL/water biphasic systems. The saturated water content of separated IL phase, 1-butyl-3-methylimidazolium bis(trifluoromethanesulfonyl)imide, was considerably improved from 0.4 wt% to 62.8 wt% by adding *N,N,N*-tripentyl-4-sulfonyl-1-butanephosphonium-type ZI (P₅₅₅C4S). In addition, the maximum water content decreased from 62.8 wt% to 34.1 wt% by increasing KH₂PO₄/K₂HPO₄ salt content in upper aqueous phosphate buffer phase. *Horse heart cytochrome c* (cyt.*c*) was dissolved selectively in IL phase by improving the water content of IL phase, and spectroscopic analysis revealed that the dissolved cyt.*c* retained its higher ordered structure. Furthermore, cyt. *c* dissolved in IL phase was re-extracted again from IL phase to aqueous phase by increasing the concentration of inorganic salts of the buffer solution.

Keywords: ionic liquid/water biphasic system; protein extraction; zwitterion; hydrophobicity; recovering

1. Introduction

Ionic liquids (ILs) are organic molten salts with very low melting temperature [1–3]. It is not so difficult to tune a wide variety of liquid properties including polarity [4], viscosity [5], and hydrophobicity [6] by appropriate combination of cation and anion species. There are a growing number of articles on aqueous biphasic systems (ABSs) based on ILs [7]. ABS offers an attractive alternative to conventional extraction methods for the separation of numerous kinds of materials such as biomolecules. Previous ABSs were mainly composed of two different polymers or a polymer and aqueous salt solution. These ABSs have been used for product recovery and downstream processing in biotechnology. It was shown that not only polymers like poly(ethylene glycol) and dextran but also ILs in combination with a kosmotropic inorganic salt are able to form ABS [8]. Some ILs having hydrophobic ions easily form liquid/liquid biphasic systems after mixing with water. These hydrophobic IL/water biphasic systems have been broadly studied and applied as novel energy-saving processes for extraction, separation, and condensation of biopolymers. More specifically, solubilization and stabilization of proteins in separated hydrophobic IL phase rather than aqueous phase would lead to the interesting application of ILs as continuous enzymatic reaction and/or product separation systems. It would be very convenient to separate enzymes from reaction products dissolved in an aqueous phase after the enzymatic reaction. However, most enzymes are not soluble in hydrophobic ILs. Some methods exist to dissolve proteins in hydrophobic ILs, such as modification of proteins with amphiphilic polymers [9,10], addition of crown ethers [11], and formation of aqueous microemulsion droplets in a hydrophobic IL [12].

According to our previous results on hydrated ILs [13–15], it would be essential to improve the hydrated state of these hydrophobic ILs for stable dissolution of proteins. However, there was no report on the successive control of water content in hydrophobic ILs until today. To control the water content of the hydrophobic ILs, one can design component ions of ILs with suitable hydrophilicity [16,17]. By this method, it is not easy to control the desirable amount of saturated water in the ILs. Other methods include simple mixing of some hydrated ILs. Mixing two or more ILs is much easier for the fine-tuning of the hydrophilicity of ILs than structural design of the ILs. However, there is always a fear of the ion exchange reaction among the mixed ions that would lead to unexpected change in the physico-chemical properties of pristine ILs. Recently, we proposed zwitterions (ZIs) as additives suitable for controlling the water content of hydrophobic ILs [18]. ZIs should prevent ion exchange in the IL mixtures, because their component cations and anions are both covalently tethered. In our previous study, saturated water content of [C₄mim][Tf₂N] was improved from 0.4 wt% to 17.8 wt% by adding hydrated 3-(1-butyl-3-imidazolium)propanesulfonate-type ZI (C₄Im3S) in appropriate amounts [18]. However the saturated water content of IL phase showed little increase after adding excess amount of water to obtain biphasic systems. This was because C₄Im3S was partitioned to aqueous phase due to high hydrophilicity of C₄Im3S. When suitably hydrated ZIs were partitioned selectively in

hydrophobic IL phase, the resulting ZI would be an effective additive to improve the water content in the separated IL phase. In the present study, we have designed ZIs with suitable hydrophilicity to accomplish both improving hydrated state and partitioning in the separated IL phase. The distribution of cyt.c in the hydrophobic IL/water mixture and re-extraction of cyt.c from IL phase to aqueous phase have been also analyzed.

2. Results and Discussion

2.1. Control of Saturated Water Content of Separated IL Phase by Mixing Phosphonium-Type ZI

Figure 1 shows both ZI and IL prepared and used in this study. *N,N,N*-Triphenyl-4-sulfonyl-1-butanephosphonium ($P_{555}C4S$) was mixed with equimolar $[C_4mim][Tf_2N]$, then pure water was added to $[C_4mim][Tf_2N]/P_{555}C4S$ mixture as 1:10 by volume. The resulting mixture was not phase-separated but homogeneously mixed. Instead, when phosphate buffer (100 mM, pH 7.0) was added to $[C_4mim][Tf_2N]/P_{555}C4S$ mixture as 1:10 by volume, the resulting mixture was found to form liquid/liquid phase separation. Then different amount of $P_{555}C4S$ was mixed with $[C_4mim][Tf_2N]$ and phosphate buffer (100 mM, pH 7.0). The resulting solutions were stirred vigorously and stored. The saturated water content of phase separated IL phase was then determined by Karl-Fischer titration method. The saturated water content of the $[C_4mim][Tf_2N]$ phase increased with the amount of the $P_{555}C4S$ as shown in Figure 2. This result suggests that the saturated water content of hydrophobic ILs can be controlled by the added amount of $P_{555}C4S$. As seen in Figure 2, addition of $P_{555}C4S$ improves the saturated water content of $[C_4mim][Tf_2N]$ from 0.4 wt% to 62.8 wt%. The water content of separated $[C_4mim][Tf_2N]$ phase without $P_{555}C4S$ was comparable to the previously reported value [19]. It follows that suitably hydrophobic $P_{555}C4S$ comprising phosphonium cation with long alkyl chains and sulfonate anions to increase saturated water content of hydrophobic IL phase.

Figure 1. Structure of ionic liquid (IL) and zwitterion (ZI) in this study; **left:** $[C_4mim][Tf_2N]$; **right:** $P_{555}C4S$.






Figure 2. Relation between water content of IL phase and fraction of ZI in IL/ZI mixtures.



2.2. Effect of K_2HPO_4/KH_2PO_4 Concentration in Aqueous Phase on the Water Content of IL Phase

We next studied the effect of concentration of K_2HPO_4/KH_2PO_4 salts in buffer solution added to mixture on the saturated water content of hydrophobic IL phase. An equimolar $P_{555}C4S$ was mixed with $[C_4mim][Tf_2N]$, and phosphate buffers with different concentration of inorganic salts (from 100 to 1000 mM, pH 7.0) were added to $[C_4mim][Tf_2N]/P_{555}C4S$ mixture as 1:10 by volume. The resulting solutions were stirred vigorously. The saturated water content of phase separated IL phase was also determined using the same procedure as describe above. The saturated water content of the $[C_4mim][Tf_2N]$ phase decreased with increasing the amount of inorganic salts (Figure 3). The decrease of saturated water content of IL phase attributed to concentrated inorganic salts strongly interacting water molecules. Rogers *et al.* reported the pioneering research on formation of ABS by the addition of inorganic salts to aqueous solution of hydrophilic ionic liquids [8]. They firstly demonstrated the formation of ABS between hydrophilic $[C_4mim]Cl$ -rich upper phase and inorganic-salt-rich bottom phase based on the salting-out phenomena of the inorganic salts [8]. Coutinho *et al.* reported effects of both IL structure and salting-out species (inorganic salts, organic salts, and so on) on the phase equilibria of ABSs [7,20–22]. They concluded that the necessary factor for formation of ABS comprising ionic liquids and conventional salts was the salting-out effect resulting in the creation of water-ion complexes that cause the dehydration of the solute and the increase in the surface tension of the cavity in the aqueous media. It is plausible that the decrease of the saturated water content of $[C_4mim][Tf_2N]$ by adding phosphate buffer was derived from the salting-out effect of K_2HPO_4/KH_2PO_4 salts.

Figure 3. Relationship between water content of IL phase and concentration of K_2HPO_4/KH_2PO_4 salts in buffer solution.



2.3. Extraction of Cytochrome c from Aqueous Phase to IL Phase by Mixing Phosphonium-Type ZI

Solubility of horse heart cytochrome c (cyt.c) in our $[C_4mim][Tf_2N]/P_{555}C4S$ /buffer mixture was analyzed to clarify the effect of ZI concentration. Both $[C_4mim][Tf_2N]$ and $P_{555}C4S$ were mixed as the mixing ratio of $[C_4mim][Tf_2N]/P_{555}C4S$ from 1.0:0.2 to 1.0:1.0. Cyt.c was dissolved into phosphate buffer (100 mM, pH 7.0) and added to $[C_4mim][Tf_2N]/P_{555}C4S$ mixture. The resulting solutions were stirred vigorously. Figure 4a shows a photograph of cyt.c in the mixtures. The distribution ratio (D) was calculated from the maximum absorbance of the Soret band around 408.0 nm of cyt.c in each phase of $[C_4mim][Tf_2N]/P_{555}C4S$ /buffer mixture. As seen in Figure 4b, the D value was found to

increase with increasing the saturated water content of IL phase by adding P₅₅₅C4S. When an equimolar amount of P₅₅₅C4S was mixed with [C₄mim][Tf₂N], the *D* value reached up to 94%. Since cyt.*c* was not soluble in [C₄mim][Tf₂N] neither pure nor buffer-saturated, it is confirmed that the P₅₅₅C4S facilitated the dissolution of cyt.*c* in the [C₄mim][Tf₂N]. Resonance Raman (RR) spectroscopy of cyt.*c* in the mixture was then undertaken in order to determine the environment of the heme vicinity. Several relevant modes observed in the RR spectra are sensitive both to axial coordination and the spin state of the iron ion of heme [23]. In the RR spectrum of cyt.*c* in buffer solution at pH 7.0, the ν_4 band around 1372 cm⁻¹ represents the valency of the heme iron. The ν_2 (1586 cm⁻¹) and ν_3 (1502 cm⁻¹) bands are both helpful in determining both the coordination state and the spin state, and these bands indicated that the heme iron ion was in a six-coordinated low spin state. Figure 4c shows the RR spectra of cyt.*c* in phosphate buffer (black line), and cyt.*c* in IL phase of [C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture (ratio of IL:ZI was 1:1) (red line). The RR spectrum of cyt.*c* in the [C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture included intense bands at 1372 cm⁻¹ (ν_4), 1502 cm⁻¹ (ν_3), 1586 cm⁻¹ (ν_2), and 1634 cm⁻¹ (ν_{10}), which indicate that the central iron ion of the heme complex is in a oxidized and six-coordinated low spin state. These characteristic bands are similar to these for cyt.*c* in buffer solution, suggesting that the dissolution of cyt.*c* in the [C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture did not give rise to a drastic change in the heme vicinity. Furthermore, spectroscopic analysis has been undertaken to determine the dissolved state and stability of cyt.*c* in the [C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture. Since cyt.*c* is known as a typical redox-active protein, the activity of the cyt.*c* in the IL phase is easily detectable by analyzing the redox response. The IL phase containing cyt.*c* was pipetted out and UV-vis spectra of the IL phase was measured. At the initial stage, both strong Soret band and broad Q-band were observed at 408 and 530 nm, respectively (Figure 5, black line). These bands indicate that cyt.*c* exists as an oxidized state in the IL phase similar to that in an aqueous buffer solution. When a small excess amount of sodium dithionite was added as a reducing agent to the IL phase, small amount of water was drained from the IL phase but cyt.*c* was still dissolved in the IL phase. Then UV-vis spectrum of the IL phase was also measured to find the reduced cyt.*c* (Figure 5, red line). The spectrum showed a sharp α - and β -band at 550 nm and 521 nm, respectively in a Q-band region and a Soret band at 415 nm, typically seen in reduced cyt.*c* in an aqueous buffer solution. Relatively large absorbance of the reduced cyt.*c* as shown in Figure 5 was due to the increased concentration of cyt.*c* by losing small amount of water as described above. This means that cyt.*c* retains the redox activity even after treated with the ILs. Both solubilization and stabilization of cyt.*c* in [C₄mim][Tf₂N] phase are accomplished by adding P₅₅₅C4S without damaging its functionality.

Figure 4. Extraction of cyt.c from aqueous to IL phase by mixing P₅₅₅C4S-type ZI. (a) Photograph of [C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture that different amount of P₅₅₅C4S added; (b) Relation between distribution ratio of cyt.c and mole fraction of ZI in IL/ZI mixtures; (c) Resonance Raman spectra of cyt.c in phosphate buffer (black line), and cyt.c in IL phase of [C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture (red line).





Figure 5. UV spectra of cyt.c extracted to IL phase before (black line) and after (red line) adding excess amount of dithionite. The [C₄mim][Tf₂N]/P₅₅₅C4S ratio was 1:1.




2.4. Effect of K₂HPO₄/KH₂PO₄ Concentration in Aqueous Phase on the Distribution Ratio of cyt.c

As mentioned above, the saturated water content of IL phase decreased upon increasing the amount of K₂HPO₄/KH₂PO₄ salts in an aqueous buffer solution phase (Figure 3). This result suggested that the *D* value of cyt.c was changed by the amount of the added inorganic salts. We then studied the effect of concentration of K₂HPO₄/KH₂PO₄ salts in buffer solution on the *D* value of cyt.c in the

[C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture. Both [C₄mim][Tf₂N] and P₅₅₅C4S were mixed with the mixing ratio of [C₄mim][Tf₂N]/P₅₅₅C4S of 1:1. Cyt.*c* was then dissolved in phosphate buffer with different concentration of inorganic salts (100, 200, and 400 mM, pH 7.0). These aqueous solutions (1 mg mL⁻¹) of cyt.*c* were added to [C₄mim][Tf₂N]/P₅₅₅C4S mixture as 1:10 by volume of IL:buffer solution. Figure 6a shows a photograph of cyt.*c* in the mixtures with different concentration of K₂HPO₄/KH₂PO₄ salts. The *D* value of cyt.*c* in the [C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture was determined by the same procedure as described above. The *D* value of cyt.*c* in the mixture decreased with the amount of K₂HPO₄/KH₂PO₄ salts of buffer solution (Figure 6b). In the case of 400 mM phosphate buffer solution, almost all cyt.*c* remained in the aqueous buffer solution phase rather than IL phase. This result suggested that the saturated water content of IL phase strongly affected to dissolution of cyt.*c* to IL phase.

Figure 6. Control of *D* value by mixing different amounts of K₂HPO₄/KH₂PO₄ salts. (a) Photograph of [C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture that different concentration of phosphate buffer used. Upper phase: aqueous buffer solution, lower phase: IL rich phase; (b) Relationship between distribution ratio of cyt.*c* in IL phase and concentration of buffer solution.




2.5. Construction of Reversible Extraction Process for cyt.*c* by [C₄mim][Tf₂N]/P₅₅₅C4S/Buffer Mixture

In previous sections, it was confirmed that the saturated water content of IL phase increased by adding P₅₅₅C4S and decreased by using high concentration buffer. Furthermore, the saturated water content of IL phase strongly affected to the *D* value of cyt.*c* in the [C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture. From the obtained results, we speculated that re-extraction of Cyt.*c* from IL phase to aqueous phase should be facilitated by adding concentrated inorganic salt solution. Then we investigated the partition behavior of cyt.*c* between IL phase and aqueous buffer phase after mixing different concentration of inorganic salt solutions. [C₄mim][Tf₂N] and P₅₅₅C4S were mixed as the mixing ratio of [C₄mim][Tf₂N]:P₅₅₅C4S were 1:1. Cyt.*c* was then dissolved into phosphate buffer (100 mM, pH 7.0). The aqueous solution of cyt.*c* (2 mg mL⁻¹) were added [C₄mim][Tf₂N]/P₅₅₅C4S mixture as 1:10 by volume of IL:buffer solution. After stirring the solution and being left until the phases became clear (Figure 7 left), 1.0 M buffer solution was then added to the solution to reach the concentration of upper inorganic salt solution to 400 mM. It is easily confirmed by naked eyes that cyt.*c* has transferred from IL phase to buffer phase by controlling the concentration of inorganic salt of buffer solution

(Figure 7 right). The environment of heme vicinity and redox activity of cyt.*c* extracted from IL phase to buffer phase in the mixture was also determined by the same method described above, and confirmed to retain their initial states in an aqueous buffer solution. Cyt.*c* was accordingly re-extracted from IL phase to buffer phase by controlling the saturated water content of IL phase with concentration of inorganic salt of buffer solution, without significant change of higher ordered structure.

In 2003, Rogers *et al.* reported the IL-based ABS by adding inorganic salts to aqueous solution of hydrophilic ILs [8]. Since then, considerable efforts have been made to construct various ABSs with hydrophilic ILs and inorganic salts as a salting-out agent [7]. However, very few reports can be found in the literature on the extraction of proteins and enzymes using hydrophobic ILs that easily undergo phase separation by adding water. This is probably due to the limited solubility of proteins in most hydrophobic ILs. In our proposed ABS systems, the water content of separated hydrophobic IL phase can be finely controlled by adding adequate amounts of ZI with suitable hydrophobicity. In addition, reversible partition of proteins was facilitated depending on the concentration of inorganic salts. This result strongly suggests that adding ZIs would provide novel bio-related engineering process including continuous enzymatic reaction systems and downstream processes by exploiting hydrophobic IL/water biphasic systems.

Figure 7. Reverse extraction process of cyt.*c* from IL phase to aqueous phase by controlling water content of IL phase.



3. Experimental Section

3.1. Materials

1-Methylimidazole, 1-bromobutane and 1,4-butanedisulfone were purchased from Tokyo Chemical Industry Co. Ltd. (Tokyo, Japan). Dipotassium hydrogen phosphate, acetonitrile, diethylether, dichloromethane, toluene, ethyl acetate, methanol were purchased from Kanto Chemical Co. Inc. (Tokyo, Japan). Potassium dihydrogen phosphate was purchased from Wako Pure Chemical Co. Ltd. (Osaka, Japan). Lithium bis(trifluoromethanesulfonyl)imide was the gift from Sumitomo 3M Ltd (Tokyo, Japan). Cytochrome *c* from horse heart and aluminum oxide were purchased from Sigma-Aldrich Japan Co. (Tokyo, Japan). Triphenylphosphine was donated from Hokko Chemical Industry Co. Ltd. (Tokyo, Japan). Both 1-methylimidazole and 1-bromobutane were distilled before use.

3.2. Preparation of ILs and ZI

3.2.1. 1-Butyl-3-Methylimidazolium Bromide ([C₄mim]Br)

A zwitterion (ZI) and an ionic liquid (IL) used in this study were synthesized as reported previously [24]. We prepared 1-butyl-3-methylimidazolium bromide ([C₄mim]Br) as follows. 1-Methylimidazole was dissolved into acetonitrile, and mixed with 1-bromobutane under dry nitrogen. The resulting solution was stirred for 24 h at room temperature. After removal of acetonitrile by evaporation, the residual liquid was repeatedly washed with excess amounts of anhydrous diethyl ether. After evaporation of the lower phase, resultant liquid was dried *in vacuo* at room temperature for 24 h. A structure of [C₄mim]Br was confirmed by ¹H-NMR spectroscopy. These spectra were observed using a JEOL ECX-400. ¹H-NMR (400MHz, CDCl₃, δ/ppm relative to TMS): 0.96 (3H, t, *J* = 14.6, N-(CH₂)₃CH₃), 1.39 (2H, m, *J* = 37.6, N-(CH₂)₂CH₂CH₃), 1.92 (2H, m, *J* = 30.2, N-CH₂CH₂CH₂CH₃), 4.14 (3H, s, N-CH₃), 4.36 (2H, t, *J* = 14.7, N-CH₂(CH₂)₂CH₃), 7.61 (1H, s, imidazolium cation), 7.73 (1H, s, imidazolium cation), 10.25 (1H, s, imidazolium cation).

3.2.2. 1-Butyl-3-Methylimidazolium Bis(trifluoromethanesulfonyl)imide ([C₄mim][Tf₂N])

We then prepared 1-butyl-3-methylimidazolium bis(trifluoromethanesulfonyl)imide ([C₄mim][Tf₂N]) as follows. First, [C₄mim]Br and lithium bis(trifluoromethanesulfonyl)imide were individually dissolved in water, and the resulting solutions were mixed under dry nitrogen gas atmosphere. The resulting solution was further stirred for 24 h at room temperature. After dissolution to dichloromethane, the resulting solution was repeatedly washed with excess amounts of water. After evaporation of the dichloromethane phase, resultant liquid was dried *in vacuo* at room temperature for 24 h. The structure of [C₄mim]Br was confirmed by ¹H-NMR spectra. These spectra were observed using a JEOL ECX-400. ¹H-NMR (400 MHz, CDCl₃, δ/ppm relative to TMS): 0.96 (3H, t, *J* = 15.1, N-(CH₂)₃CH₃), 1.36 (2H, m, *J* = 37.1, N-(CH₂)₂CH₂CH₃), 1.85 (2H, m, *J* = 30.2, N-CH₂CH₂CH₂CH₃), 3.93 (3H, s, N-CH₃), 4.17 (2H, t, *J* = 15.1, N-CH₂(CH₂)₂CH₃), 7.27 (1H, s, imidazolium cation), 7.31 (1H, s, imidazolium cation), 8.73 (1H, s, imidazolium cation).

3.2.3. *N,N,N*-Triphenyl-4-Sulfonyl-1-Butanephosphonium (P₅₅₅C4S)

We next prepared *N,N,N*-triphenyl-4-sulfonyl-1-butanephosphonium (P₅₅₅C4S) as follows. Triphenylphosphine and 1,4-butanediol were dissolved into toluene, and the resulting solutions were mixed under dry nitrogen gas atmosphere. The resulting solution was stirred for three days at 100 °C. After removal of toluene by evaporation, the residual liquid was repeatedly washed with excess amounts of anhydrous diethylether. A resultant solid was dissolved in dichloromethane, and the solution was passed through a column filled with aluminum oxide. It was then further purified by recrystallization from ethylacetate, and resultant white powder was dried *in vacuo* at room temperature for 24 h. Structure of P₅₅₅C4S was confirmed by ¹H-NMR spectra. ¹H-NMR (400 MHz, CDCl₃, δ/ppm relative to TMS): 0.92 (9H, t, P-(CH₂)₄CH₃), 1.37 (6H, m, P-(CH₂)₂CH₂CH₂CH₃), 1.48 (12H, m, P-CH₂CH₂CH₂CH₂CH₃), 1.76 (2H, m, P-CH₂CH₂(CH₂)₂SO₃), 2.00 (2H, t, P-CH₂(CH₂)₃SO₃), 2.22 (6H, m, P-CH₂(CH₂)₃CH₃), 2.47 (2H, m, P-(CH₂)₂CH₂CH₂SO₃), 2.87 (2H, t, P-(CH₂)₃CH₂SO₃).

3.3. Water Content of Hydrophobic IL Phase after Mixing with P₅₅₅C4S

Both [C₄mim][Tf₂N] and P₅₅₅C4S were mixed at the mixing ratio of [C₄mim][Tf₂N]/P₅₅₅C4S from 1:0.2 to 1:1. Phosphate buffer (100 mM, pH 7.0) was added to [C₄mim][Tf₂N]/P₅₅₅C4S mixture as 1:10 by volume, and the resulting solutions were stirred vigorously. The saturated water content of phase separated IL phases was determined by Karl-Fischer titration method (Kyoto Electronics; MKC-520N) after dilution of IL phase by dehydrated methanol.

3.4. Control of Water Content of Hydrophobic IL Phase by Adding Phosphate Buffers

P₅₅₅C4S was mixed with equimolar amount of [C₄mim][Tf₂N], then phosphate buffer with different concentration of K₂HPO₄/KH₂PO₄ salts (from 0.1 to 1.0 M, pH 7.0) were added to [C₄mim][Tf₂N]/P₅₅₅C4S mixture as 1:10 by volume of IL:buffer solution. The resulting solutions were stirred vigorously. The saturated water content of phase separated IL phases was determined by using the same procedure as describe above.

3.5. Distribution Ratio of cyt.c in Hydrophobic IL/Water Biphasic System

[C₄mim][Tf₂N] and P₅₅₅C4S were mixed at the mixing ratio of from 1:0.2 to 1.0:1.0. Cytochrome *c* from *horse heart* was dissolved into phosphate buffer with different concentration of K₂HPO₄/KH₂PO₄ salts (100, 200, and 400 mM, pH 7.0). These aqueous solutions (1 mg mL⁻¹) of cyt.*c* were added to [C₄mim][Tf₂N]/P₅₅₅C4S mixtures as 1:10 by volume of IL:buffer solution. The resulting solutions were stirred vigorously. The distribution ratio (*D*) was calculated from the absorbance of cyt.*c* in each phase of [C₄mim][Tf₂N]/P₅₅₅C4S/buffer mixture, with the equation; $D = (Abs_{IL} \times V_{IL}) / (Abs_{IL} \times V_{IL} + Abs_{buffer} \times V_{buffer})$, where *Abs*_{IL} and *Abs*_{buffer} corresponded to the absorbance of IL phase and buffer phase, respectively, and *V*_{IL} and *V*_{buffer} corresponded to the volume of IL phase and buffer phase, respectively. The maximum absorption of the Soret band around 408.0 nm was used as the absorbance. The absorbance of proteins was determined with UV-vis spectrometer (UV-2550, Shimadzu Co., Kyoto, Japan).

The environment of heme vicinity of cyt.*c* in the IL phase was determined by Resonance Raman (RR) Spectroscopy. The Raman spectra were obtained on a JASCO NRS-1000 spectrometer with a Kaiser Optical holographic notch-plus filter and a liquid N₂-cooled charge-coupled device (CCD) detector. Data were accumulated for 150 s with spectral resolution of 4.0 cm⁻¹. The excitation source was a Coherent Innova 90C Kr laser with a 20 mW beam at a 413.1 nm excitation wavelength.

The redox activity of cyt.*c* in the IL phase was determined by UV-vis spectrometer. An excess amount of sodium dithionite was added and stirred gently. The absorption of cyt.*c* in the IL phase before and after addition of the sodium dithionite was determined by UV-vis spectroscopy.

3.6. Extraction of cyt.c Dissolved in IL Phase to Aqueous Phase

[C₄mim][Tf₂N] and P₅₅₅C4S were mixed as the mixing ratio of [C₄mim][Tf₂N]/P₅₅₅C4S as 1:1. Cyt.*c* was dissolved into phosphate buffer (100 mM, pH 7.0). The aqueous solution of cyt.*c* (2 mg mL⁻¹) was added to [C₄mim][Tf₂N]/P₅₅₅C4S mixture as 1:10 by volume of IL:buffer solution. The resulting solutions were stirred vigorously. Then, the concentration of K₂HPO₄/KH₂PO₄ salts was set to 400 mM by adding high concentration buffer solution (1000 mM) to the mixture. The environment of heme

vicinity and redox activity of cyt.c extracted from IL phase to buffer phase in the mixture was determined by RR spectroscopy and UV-vis spectroscopy, respectively.

4. Conclusions

In conclusion, we have used 1-butyl-3-methylimidazolium bis(trifluoromethanesulfonyl)imide as a hydrophobic IL to form phase separated state with aqueous buffer solution. The saturated water content of the hydrophobic IL phase was considerably improved from 0.4 wt% to 62.8 wt% by adding *N,N,N*-tripentyl-4-sulfonyl-1-butanephosphonium-type zwitterion. The biphasic system containing 62.8 wt% water in IL phase successfully dissolved *horse heart* cytochrome *c* (cyt.c) without significant change of the higher ordered structure. The saturated water content of IL phase was controlled by the concentration of inorganic salt of buffer solution. Cyt.c dissolved in IL phase was extracted again from IL phase to aqueous phase by controlling the concentration of inorganic salt in the mixture without significant change of the higher ordered structure.

Acknowledgments

This study was supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (No. 21225007).

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Wilkes, J.S.; Zaworotko, M.J. Air and water stable 1-ethyl-3-methylimidazolium based ionic liquids. *J. Chem. Soc. Chem. Commun.* **1992**, 965–967.
2. Welton, T. Room-temperature ionic liquids. Solvents for synthesis and catalysis. *Chem. Rev.* **1999**, *99*, 2071–2083.
3. Wasserscheid, P.; Welton, T. *Ionic Liquids in Synthesis*; Wiley-VCH: Weinheim, Germany, 2003.
4. Crowhurst, L.; Mawdsley, P.R.; Perez-Arlandis, J.M.; Salter, P.A.; Welton, T. Solvent–solute interactions in ionic liquids. *Phys. Chem. Chem. Phys.* **2003**, *5*, 2790–2794.
5. Böhnhöte, P.; Dias, A.P.; Papageorgiou, N.; Kalyanasundaram, K.; Grätzel, M. Hydrophobic, highly conductive ambient-temperature molten salts. *Inorg. Chem.* **1996**, *35*, 1168–1178.
6. Seddon, K.R.; Stark, A.; Torres, M.J. Influence of chloride, water, and organic solvents on the physical properties of ionic liquids. *Pure Appl. Chem.* **2000**, *72*, 2275–2287.
7. Freire, M.G.; M Cláudio, A.F.; Araújo, J.M.M.; Coutinho, J.A.P.; Marrucho, I.M.; Lopes, J.N.C.; Rebelo, L.P.N. Aqueous biphasic systems: A boost brought about by using ionic liquids. *Chem. Soc. Rev.* **2012**, *41*, 4966–4995.
8. Gutowski, K.E.; Broker, G.A.; Willauer, H.D.; Huddleston, J.G.; Swatloski, R.P.; Holbrey, J.D.; Rogers, R.D. Controlling the aqueous miscibility of ionic liquids: Aqueous biphasic systems of water-miscible ionic liquids and water-structuring salts for recycle, metathesis, and separations. *J. Am. Chem. Soc.* **2003**, *125*, 6632–6633.

9. Kawahara, N.V.; Ohno, H. Stability of poly(ethylene oxide)-modified myoglobin in ion conductive polymers at high temperatures. *Solid State Ionics* **1998**, *114*, 161–166.
10. Nakashima, K.; Maruyama, T.; Kamiya, N.; Goto, M. Comb-shaped poly(ethylene glycol)-modified subtilisin carlsberg is soluble *and highly active in* ionic liquids. *Chem. Commun.* **2005**, 4297–4299.
11. Shimojo, K.; Nakagawa, H.; Kamiya, N.; Goto, M. Crown ether-mediated extraction and functional conversion of cytochrome c in ionic liquids. *Biomacromolecules* **2006**, *7*, 2–5.
12. Moniruzzaman, M.; Kamiya, N.; Nakashima, K.; Goto, M. Water-in-ionic liquid microemulsions as a new medium for enzymatic reactions. *Green Chem.* **2008**, *10*, 497–500.
13. Fujita, K.; MacFarlane, D.R.; Forsyth, M. Protein solubilising and stabilising ionic liquids. *Chem. Commun.* **2005**, 4804–4806.
14. Fujita, K.; Ohno, H. Enzymatic activity and thermal stability of metallo proteins in hydrated ionic liquids. *Biopolymers* **2010**, *93*, 1093–1099.
15. Fujita, K.; MacFarlane, D.R.; Forsyth, M.; Yoshizawa-Fujita, M.; Murata, K.; Nakamura, N.; Ohno, H. Solubility and stability of cytochrome c in hydrated ionic liquids: Effect of oxo acid residues and kosmotropicity. *Biomacromolecules* **2007**, *8*, 2080–2086.
16. Kohno, Y.; Ohno, H. Temperature-responsive ionic liquid/water interfaces: Relation between hydrophilicity of ions and dynamic phase change. *Phys. Chem. Chem. Phys.* **2012**, *14*, 5063–5070.
17. Kohno, Y.; Ohno, H. Ionic liquid/water mixtures: From hostility to conciliation. *Chem. Commun.* **2012**, *48*, 7119–7130.
18. Ito, Y.; Kohno, Y.; Nakamura, N.; Ohno, H. Addition of suitably-designed zwitterions improves the saturated water content of hydrophobic ionic liquids. *Chem. Commun.* **2012**, *48*, 11220–11222.
19. Huddleston, J.G.; Visser, A.E.; Reichert, W.M.; Willauer, H.D.; Broker, G.A.; Rogers, R.D. Characterization and comparison of hydrophilic and hydrophobic room temperature ionic liquids incorporating the imidazolium cation. *Green Chem.* **2001**, *3*, 156–164.
20. Freire, M.G.; Carvalho, P.J.; Silva, A.M.S.; Santos, L.M.N.B.F.; Rebelo, L.P.N.; Marrucho, I.M.; Coutinho, J.A.P. Ion specific effects on the mutual solubilities of water and hydrophobic ionic liquids. *J. Phys. Chem. B* **2009**, *113*, 202–211.
21. Freire, M.G.; Neves, C.M.S.S.; Silva, A.M.S.; Santos, L.M.N.B.F.; Marrucho, I.M.; Rebelo, L.P.N.; Shah, J.K.; Maginn, E.J.; Coutinho, J.A.P. ¹H NMR and molecular dynamics evidence for an unexpected interaction on the origin of salting-in/salting-out phenomena. *J. Phys. Chem. B* **2010**, *114*, 2004–2014.
22. Shahriari, S.; Neves, C.M.S.S.; Freire, M.G.; Coutinho, J.A.P. Role of the Hofmeister series in the formation of ionic-liquid-based aqueous biphasic systems. *J. Phys. Chem. B* **2012**, *116*, 7252–7258.
23. Scott R.A.; Mauk, A.G. *Cytochrome c: A Multidisciplinary Approach*; University Science Books: Sausalito, CA, USA, 1996.
24. Ueda, S.; Kagimoto, J.; Ichikawa, T.; Kato, T.; Ohno, H. Anisotropic proton-conductive materials formed by the self-organization of phosphonium-type zwitterions. *Adv. Mater.* **2011**, *23*, 3071–3074.