

**Supplementary material for:**

**Novel Thiazolo[5,4-b]phenothiazine Derivatives: Synthesis, Structural Characterization, and In Vitro Evaluation of Antiproliferative Activity against Human Leukaemia**



B. Brem<sup>1</sup>, E. Gal<sup>1</sup>, L. Găină<sup>1</sup>, L. Silaghi-Dumitrescu<sup>1</sup>, E. Fischer-Fodor<sup>2,3</sup>, C. I. Tomuleasa<sup>2,4</sup>, A. Grozav<sup>5</sup>, V. Zaharia<sup>5</sup> L. Filip<sup>5\*</sup> and C. Cristea<sup>1\*</sup>



Address. <sup>1</sup> Faculty of Chemistry and Chemical Engineering, Babes-Bolyai University, Cluj-Napoca, Romania. <sup>2</sup> Tumor Biology Department, I. Chiricuta Oncology Institute, Cluj Napoca, Romania. <sup>3</sup> Medfuture Research Center, I. Hatieganu University of Medicine and Pharmacy, Cluj Napoca, Romania <sup>4</sup> Research Center for Functional Genomics and Translational Medicine, I. Hatieganu University of Medicine and Pharmacy, Cluj Napoca, Romania. <sup>5</sup> Faculty of Pharmacy, "Harițeganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania.



\*Corresponding authors



**Table of contents**


<b>NMR-spectra of 3a-3e</b>	<b>S2-S5</b>
<b>2D NOESY of 3a</b>	<b>S6</b>
<b>HRMS spectra of 3a-3e</b>	<b>S7-S9</b>
<b>Frontier molecular orbital plots of 3a-3d</b>	<b>S10</b>
<b>In vitro cell metabolic activity in populations of THP-1, HL-60, and PBMC cell cultures</b>	<b>S11</b>














3a comp 170531163709 #18-22 RT: 0.43-0.55 AV: 5 NL: 1.71E7  
T: FTMS + p ESI Full ms [150.00-2000.00]





3b comp 170531163709 #7 RT: 0.16 AV: 1 NL: 2.04E7  
T: FTMS + p ESI Full ms [150.00-2000.00]




3c comp 170607194459 #1-6 RT: 0.01-0.16 AV: 6 NL: 4.46E5  
T: FTMS + p ESI Full ms [150.00-2000.00]








**Frontier MO plots**





**Figure S1.** In vitro cell metabolic activity in populations of THP-1, HL-60, and PBMC cell cultures treated with N-(phenothiazinyl)-thioamides **2a**, **2c**, and TAP TZ **3e**, monitored by Alamar Blue staining and measurement of fluorescence intensity at 620 nm