



Supplementary Materials

Epigallocatechin-3-Gallate Suppresses Human Herpesvirus 8 Replication and Induces ROS Leading to Apoptosis and Autophagy in Primary Effusion Lymphoma Cells

Ching-Yi Tsai ¹, Chang-Yu Chen ¹, Yee-Hsuan Chiou ², Huey-Wen Shyu ¹, Kuan-Hua Lin ¹,
Miao-Chen Chou ¹, Mei-Han Huang ¹ and Yi-Fen Wang ^{1,*}

¹ Department of Medical Laboratory Science and Biotechnology, Fooyin-University, Kaohsiung, Taiwan; a07646815@yahoo.com.tw (C.-Y.T.); chencyallen1689@gmail.com (C.-Y.C.); shyuhw@hotmail.com (H.-W.S.); su6473@mail2000.com.tw (K.-H.L.); miaochen119@gmail.com (M.-C.C.); hnmhmtw8@gmail.com (M.-H.H.)

² Department of Pediatrics, Kaohsiung Veterans General Hospital, Kaohsiung, Taiwan; chysn@ms6.hinet.net

* Correspondence: sc071@fy.edu.tw; Tel.: + ; Fax: +886-7-782-7162




Figure S1. Effects of necroptosis inhibitor (necrostatin) on EGCG induced cell death of PEL cells. BCBL-1 cells were pretreated with necrostatin-1 1 h before EGCG (20 μ g/ml) treatment. Cell viability was evaluated by trypan blue exclusion assay 24 h later. The cells in supplemented medium were used as control. The values represent mean \pm SE of three independent experiments and are presented as the percentage of control. * $p < 0.05$ indicates significant differences between the cells treated with EGCG and the cells co-treated with EGCG and necrostatin-1.




Figure S2. JNK inhibitor SP600125 enhanced the cytotoxicity of EGCG in HHV8 harboring PEL cells.