Table 1. Summary of ZnT8-KO mouse studies

	Deletion	Deletion	Genetic	Crystalization of	Proinsulin	Glucose	Glucose-stimulated	Insulin level in
Study	location	strategy	background	insulin granules	concentration	intolerance	insulin secretion	vivo
				granule/ rod-shaped		Impaired		
Nicolson et al. (2009)	Whole body	Exon 1	Mixed	cores	Unchanged	(only males)	Increased	Impaired
					Not			
Pound et al. (2009)	Whole body	Exon 3	Mixed	Not examined	examined	Normal	Decreased	Impaired
				Loss of dense core				
Lemaire et al. (2009)	Whole body	Exon 1	Mixed	granule	Unchanged	Normal	Unchanged	Unchanged
	specific			granule/ rod-shaped				
Wijesekara et al. (2010)	(Ins2Cre)	Exon 1	Mixed	cores	Increased	Impaired	Decreased	Unchanged
Pound et al. (2012)	Whole body	Exon 3	C57BL/6J	Unchanged	Decreased	Impaired	Decreased	Unchanged
	specific			Loss of dense core				
Tamaki et al. (2013)	(Ins2Cre)	Exon 5	C57BL/6J	granules	Increased	Impaired	Increased	Impaired
	specific			granule/ rod-shaped	Not			
Mitchell et al. (2016)	(Ins1Cre)	Exon 1	C57BL/6J	cores	examined	Impaired	Unchanged	Impaired