




Antisense Oligonucleotide-Based Splicing Correction

in Individuals with Leber Congenital Amaurosis due

to Compound Heterozygosity for the c.2991+1655A>G 3

Mutation in CEP290 4

- 5 Lonneke Duijkers 1, L. Ingeborgh van den Born 2, John Neidhardt 3,4, Nathalie M. Bax 5,6,
- 6 Laurence H.M. Pierrache ^{2,7}, B. Jeroen Klevering ^{5,6}, Rob W.J. Collin ^{1,6,*} and Alejandro Garanto ^{1,6}
- Department of Human Genetics, Radboud University Medical Center, 6525 GA Nijmegen, 8 The Netherlands; lonneke.duijkers@radboudumc.nl (L.D.); alex.garanto@radboudumc.nl (A.G.) 9
 - The Rotterdam Eye Hospital, 3011 BH Rotterdam, The Netherlands; born@oogziekenhuis.nl (L.I.v.d.B.); L.Pierrache@oogziekenhuis.nl (L.H.M.P.)
- 11 Human Genetics, Faculty of Medicine and Health Sciences, University of Oldenburg, 26129 Oldenburg, 12 Germany; john.neidhardt@uni-oldenburg.de 13
 - ⁴ Research Center Neurosensory Science, University Oldenburg, 26129 Oldenburg, Germany
 - ⁵ Department of Ophthalmology, Radboud University Medical Center, 6525 GA Nijmegen, The Netherlands; nathalie.bax@radboudumc.nl (N.M.B.); jeroen.klevering@radboudumc.nl (B.J.K.)
- ⁶ Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Center, 16 17 6525 GA Nijmegen, The Netherlands
 - Department of Ophthalmology, Erasmus Medical Center, 3015 CE Rotterdam, The Netherlands
- 19 Correspondence: rob.collin@radboudumc.nl; Tel.: +31-24-3613750; Fax: +31-24-3668752



20 21

22

23

24

25

10

14

15

18

Figure S1. Cilium staining in all fibroblast cell lines. Representative images of each fibroblast cell line used in the ciliation studies. Control (CON), LCA-affected (LCA) and carrier (HET1) cell lines were used for immunocytochemical analysis in the non-treated (NT) and AON-treated (AON) situations. DAPI was used to stain the nucleus (blue) and Acetylated Tubulin for cilium axoneme detection (red). Scale bar represents 20 µm.