Glycidamide Promotes the Growth and Migratory Ability of Prostate Cancer Cells by Changing the Protein Expression of Cell Cycle Regulators and Epithelial-to-Mesenchymal Transition (EMT)-Associated Proteins with Prognostic Relevance

Titus I. Ekanem^{1,2}, Chi-Chen Huang^{3,+}, Ming-Heng Wu^{4,+}, Ding-Yen Lin^{5,6}, Wen-Fu T. Lai^{7,8,9} and Kuen-Haur Lee^{6,10,*}

¹Ph.D. Program for Cancer Molecular Biology and Drug Discovery, College of Medical Science and Technology, Taipei Medical University and Academia Sinica ²Department of Hematology, University of Uyo, Uyo, Nigeria ³Graduate Institute of Neural Regenerative Medicine, College of Medical Science and Technology, Taipei Medical University, Taipei, Taiwan ⁴Graduate Institute of Translational Medicine, Taipei Medical University, Taipei, Taiwan. ⁵Department of Biotechnology and Bioindustry Sciences, College of Bioscience and Biotechnology, National Cheng Kung University, Tainan ⁶Graduate Institute of Cancer Biology and Drug Discovery, College of Medical Science and Technology, Taipei Medical University, Taipei, Taiwan ⁷McLean Imaging Center, McLean Hospital, Harvard Medical School, Belmont, MA, United States of America ⁸Department of Research, Taipei Medical University/Shuang-Ho Hospital, New Taipei City, Taiwan ⁹Department of Dentistry, Taipei Medical University/Shuang-Ho Hospital, New Taipei City, Taiwan ¹⁰Ph.D. Program for Cancer Molecular Biology and Drug Discovery, College of Medical Science and Technology, Taipei Medical University, Taipei, Taiwan


⁺These authors contributed equally to this work.

* Correspondence: Dr. Kuen-Haur Lee, Graduate Institute of Cancer Biology and Drug Discovery, College of Medical Science and Technology, Taipei Medical University, No. 250 Wu-Hsing Street, Taipei 11031, Taiwan. Tel: +886-2-27361661 ext. 7627, Fax: +886-2-66387537, E-mail: khlee@tmu.edu.tw


Supplementary Material

Figure S1. Survival analysis of prostate cancer patients with SurvExpress (n=140). Low expression of CCND1 (**A**), CDK4 (**B**), and TWIST1 (**C**) were correlated with good prognosis of prostate cancer patients. High expression of SNAI1 (**D**), SNAI2 (**E**), and CDH1 (**F**) were correlated with good prognosis of prostate cancer patients.

Figure S2. Kaplan–Meier curves according to any two-gene models. Clinical outcomes for the combinations of CDK4/TWIST1 (**A**), CDK4/SNAI2 (**B**), and TWIST1/SNAI2 (**C**) mRNA status of prostate cancer patients.



Supplementary Fig. 1



Supplementary Fig. 2