

Therapeutic Effect of an Antibody-Derived Peptide in a *Galleria mellonella* Model of Systemic Candidiasis

Emerenziana Ottaviano ¹, Elisa Borghi ¹, Laura Giovati ², Monica Falleni ¹, Delfina Tosi ¹, Walter Magliani ², Giulia Morace ¹, Stefania Conti ^{2*} and Tecla Ciociola ²

¹ Department of Health Sciences, University of Milan, 20142 Milan, Italy; emerenziana.ottaviano@unimi.it (E.O.); elisa.borghi@unimi.it (E.B.); monica.falleni@unimi.it (M.F.); delfina.tosi@unimi.it (D.T.); giulia.morace@unipr.it (G.M.)

² Department of Medicine and Surgery, University of Parma, 43126 Parma, Italy; laura.giovati@unipr.it (L.G.); walter.magliani@unipr.it (W.M.); tecla.ciociola@unipr.it (T.C.)

* Correspondence: stefania.conti@unipr.it; Tel.: +390521903492




Figure S1. Peptides' toxicity in *Galleria mellonella* larvae. Larvae were inoculated with peptides (15 $\mu\text{mol/kg}$, single injection 10 μl) or saline (control group). The survival curves of treated larvae were not significantly different from that of control group, as assessed by Mantel-Cox log-rank test.




Figure S2. Histology of an untouched *Galleria mellonella* larva. Larval tissues are well preserved and only a few hemocytes can be observed in the hemolymph near the fat body (circle). Ct: cuticle; Fb: fat body; GI: gastrointestinal tract; T: trachea. Bar = 250 μm.