

Supporting information


Cytotoxic Potential of Novel Quinoline Derivative 11-(1,4-Bisaminopropylpiperazinyl)-5-Methyl-5H-indolo[2,3-b]quinoline against Different Cancer Cell Lines via Activation and Deactivation of the Expression of Some Proteins

Sara Fathy Abd Elrahman ¹, Abdullah A. S. Ahmed ¹, Doaa Abd Elsatar ¹, Salma Elkady ¹, Amira Elgendy ¹, Fatma Alnakeeb ¹, Elshaymaa I. Elmongy ², Hanan A. Henidi ^{3,*}, Saad M. El-Gendy ⁴, Ibrahim El Tantawy El Sayed ^{1,*}, Ahmed A. El-Gokha ¹ and Mabrouk Attia Abd Eldaim ⁵

- ¹ Department of Chemistry, Faculty of Science, Menoufia University, Shibin El-Kom 32511, Egypt; sara.fathy.mostafa@gmail.com (S.F.A.E.); chemist_abdullah_2009@yahoo.com (A.A.S.A.); douaaabdelatar94@gmail.com (D.A.E.); salmaelkady16@yahoo.com (S.E.); amirajulita765@gmail.com (A.E.); tamtamalnakeeb@gmail.com (F.A.); aelgokha@yahoo.com (A.A.E.-G.)
- ² Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Helwan University, Cairo 11795, Egypt; shaymaa.taha@pharm.helwan.edu.eg
- ³ Research Department, Health Sciences Research Center, Princess Nourah bint Abdulrahman University, Riyadh 84428, Saudi Arabia
- ⁴ Department of Cancer Biology, National Cancer Institute, Cairo University, Cairo 11562, Egypt; saadelgendy2004@yahoo.com
- ⁵ Department of Biochemistry and Chemistry of Nutrition, Faculty of Veterinary Medicine, Menoufia University, Shibin El-Kom 32511, Egypt; mabroukattia@vet.menofia.edu.eg
- * Correspondence: hahenidi@pnu.edu.sa (H.A.H.); ibrahimtantawy@science.menofia.edu.eg (I.E.T.E.S.)



¹H NMR of 11(1,4-Bisaminopropylpiperazinyl) indoloquinoline (BAPPN)



¹³C NMR of 11(1,4-Bisaminopropylpiperazinyl) indoloquinoline (BAPPN)