

Table S1. Patient demographics.

No	Sex /Age	Duration	Type	Activity	Previous Treatment	Lesional Location and Sampling Sites
1	M/14	2 yrs	Non-segmental	Stable	Steroid+Excimer	Face, upper and <u>lower extremities</u> , dorsa of feet
2	M/15	Several yrs	Non-segmental	Active	None	<u>Face, upper extremities, back</u>
3	M/16	6 mos	Non-segmental	Stable	Steroid+NB-UVB	<u>Trunk</u> , inguinal, lower extremities
4	M/20	2 yrs	Non-segmental	Active	Steroid	Scalp, <u>forehead</u> , nose, perioral, neck, anterior chest, both upper extremities
5	M/31	9 yrs	Non-segmental	Active	Steroid+NB-UVB	Wrists, dorsa of feet, anterior chest, <u>abdomen</u>
6	F/18	1 yrs	Non-segmental	Active	None	<u>Arms</u> , trunk
7	F/18	>5 yrs	Non-segmental	Active	Steroid+NB-UVB	Hands, wrists, anterior chest, <u>abdomen</u> , inguinal
8	F/54	2 yrs	Non-segmental	Active	None	<u>Anterior chest, abdomen, both groins</u>
9	M/18	4 yrs	Segmental	Stable	Steroid+NB-UVB	<u>Lt forearm</u>
10	F/16	a few yrs	Segmental	Stable	Steroid+NB-UVB	<u>Lt breast</u> , anterior chest
11	F/51	a few yrs	Segmental	Active	None	<u>Rt. Buttock</u> , posterior thigh
12	F/52	2 yrs	Segmental	Stable	Tacrolimus+NB-UVB	Rt eyelid, nose, <u>forehead</u>
13	F/8	4 yrs	Mixed	Stable	Steroid+NB-UVB	Rt upper lip, chin, <u>neck</u> , scalp
14	M/15	>10 yrs	Mixed	Stable	None	<u>Rt forehead</u>
15	M/18	a few yrs	Localized	Stable	Steroid+Excimer	<u>Rt inner thigh</u>
16	F/9	5 yrs	Localized	Stable	Steroid+NB-UVB	Posterior neck, <u>anterior chest</u>

yrs: years, mos; months, steroid: topical steroid, tacrolimus: topical tacrolimus, Lt: left, Rt: right Sampling site of each patient was underlined.