

**Table S5:** Uncorrected p-distances, Tamura—Nei genetic distances within the studied Cheliferidae pseudoscorpions. DL: Delimited lineage, H: number of haplotypes within the lineage, S.E.: standard error.

Species	DL	H	p-distance [S.E.]	Tamura - Nei [S.E.]
<i>Chelifer cancroides</i>	A	2	0.02 [0.01]	0.02 [0.01]
<i>Chelifer cancroides</i>	B	3	0.04 [0.01]	0.05 [0.01]
<i>Chelifer cancroides</i>	C	16	0.01 [0.00]	0.01 [0.00]
<i>Dactylochelifer sp.</i>	A	1	N/A [N/A]	N/A [N/A]
<i>Dactylochelifer sp.</i>	B	5	0.02 [0.00]	0.02 [0.00]
<i>Dactylochelifer sp.</i>	C	1	N/A [N/A]	N/A [N/A]
<i>Dactylochelifer copiosus</i>	D	2	0.01 [0.00]	0.01 [0.00]
<i>Dactylochelifer sp.</i>	E	4	0.03 [0.00]	0.04 [0.01]
<i>Dactylochelifer latreillii</i>	F	10	0.01 [0.00]	0.01 [0.00]
<i>Dactylochelifer latreillii</i>	G	7	0.01 [0.00]	0.01 [0.00]
<i>Dactylochelifer latreillii</i>	H	11	0.01 [0.00]	0.02 [0.00]
<i>Rhacochelifer sp.</i>	A	1	N/A [N/A]	N/A [N/A]
<i>Rhacochelifer sp.</i>	B	2	0.01 [0.00]	0.01 [0.00]
<i>Rhacochelifer sp.</i>	C	2	0.02 [0.01]	0.02 [0.01]
<i>Rhacochelifer sp.</i>	D	21	0.03 [0.00]	0.03 [0.00]
<i>Hysterochelifer meridianus</i>	A	26	0.03 [0.00]	0.03 [0.00]
<i>Hysterochelifer tuberculatus</i>	B	7	0.02 [0.00]	0.02 [0.00]
<i>Hysterochelifer tuberculatus</i>	C	2	0.00 [0.00]	0.00 [0.00]
<i>Hysterochelifer tuberculatus</i>	D	2	0.03 [0.01]	0.03 [0.01]
<i>Hysterochelifer tuberculatus</i>	E	4	0.01 [0.00]	0.01 [0.00]
<i>Hysterochelifer tuberculatus</i>	F	8	0.01 [0.00]	0.01 [0.00]
<i>Hysterochelifer tuberculatus</i>	G	2	0.01 [0.00]	0.01 [0.00]
<i>Hysterochelifer tuberculatus</i>	H	3	0.01 [0.00]	0.01 [0.00]
<i>Hysterochelifer tuberculatus</i>	I	7	0.01 [0.00]	0.01 [0.00]