

A list of sampled species with their ecological classification

| Ecological classification | Number of species | Species | Family |
|-----------------------------|-------------------|--|----------------|
| TROGLOBIONT | 3 | <i>Catops tristis infernus</i> Szymczakowski, 1957 | Leiodidae |
| | | <i>Choleva lederiana gracilentia</i> Szymczakowski, 1957 | Leiodidae |
| | | <i>Speonomus hydrophilus</i> Jeannel, 1907 | Leiodidae |
| EUTROGLOPHILE | 4 | <i>Catops picipes</i> (Fabricius, 1787) | Leiodidae |
| | | <i>Omalius validum</i> Kraatz, 1857 | Staphylinidae |
| | | <i>Tachinus subterraneus</i> (Linnaeus, 1758) | Staphylinidae |
| | | <i>Quedius mesomelinus mesomelinus</i> (Marsham, 1802) | Staphylinidae |
| SUBTROGLOPHILE/ MSS SPECIES | 32 | <i>Anthobium atrocephalum</i> (Gyllenhal, 1810) | Staphylinidae |
| | | <i>Atheta sodalis</i> (Erichson, 1837) | Staphylinidae |
| | | <i>Catops fuscus</i> (Panzer, 1794) | Leiodidae |
| | | <i>Catops morio</i> (Fabricius, 1787) | Leiodidae |
| | | <i>Catops neglectus</i> E.G. Kraatz, 1852 | Leiodidae |
| | | <i>Catops nigricans</i> (Spence, 1813) | Leiodidae |
| | | <i>Catops fuliginosus</i> W.F. Erichson, 1837 | Leiodidae |
| | | <i>Catops kirbyi</i> (Spence, 1813) | Leiodidae |
| | | <i>Catops subfuscus</i> Kellner, 1846 | Leiodidae |
| | | <i>Choleva bicolor</i> Jeannel, 1923 | Leiodidae |
| | | <i>Choleva glauca</i> Britten, 1918 | Leiodidae |
| | | <i>Choleva agilis</i> (J.K.W. Illiger, 1798) | Leiodidae |
| | | <i>Choleva oblonga</i> Latreille, 1806 | Leiodidae |
| | | <i>Choleva cisteloides</i> (Frölich, 1799) | Leiodidae |
| | | <i>Choleva nivalis</i> (E.G. Kraatz, 1856) | Leiodidae |
| | | <i>Choleva paskoviensis</i> E. Reitter, 1913 | Leiodidae |
| | | <i>Choleva spadicea</i> (J. Sturm, 1839) | Leiodidae |
| | | <i>Choleva sturmii</i> Ch.N.F. Brisout de Barneville, 1863 | Leiodidae |
| | | <i>Cryptophagus distinguendus</i> J. Sturm, 1845 | Cryptophagidae |
| | | <i>Cryptophagus scutellatus</i> L. Newman, 1834 | Cryptophagidae |

A list of sampled species with their ecological classification

| | | | |
|------------|----|--|---------------|
| TROGLOXENE | | <i>Mycetaea subterranea</i> (Fabricius, 1801) | Endomychidae |
| | | <i>Nargus velox</i> (Spence, 1813) | Leiodidae |
| | | <i>Nargus wilkinii</i> (Spence, 1815) | Leiodidae |
| | | <i>Nemadus colonoides</i> (E.G. Kraatz, 1851) | Leiodidae |
| | | <i>Neuraphes talparum</i> Lokay, 1920 | Scydmenidae |
| | | <i>Omalium caesum</i> Gravenhorst, 1806 | Staphylinidae |
| | | <i>Omalium excavatum</i> Stephens, 1834 | Staphylinidae |
| | | <i>Omalium rivulare</i> (Paykull, 1789) | Staphylinidae |
| | | <i>Othius punctulatus</i> (Goeze, 1777) | Staphylinidae |
| | | <i>Othius subuliformis</i> Stephens, 1833 | Staphylinidae |
| | | <i>Ptomaphagus sericatus</i> Chaudoir, 1845 | Leiodidae |
| | | <i>Quedius limbatus</i> (Heer, 1839) | Staphylinidae |
| | | <i>Syntomium aeneum</i> (Müller, 1821) | Staphylinidae |
| | 66 | <i>Abax ovalis</i> (C. Duftschmid, 1812) | Carabidae |
| | | <i>Abax parallelus</i> (C. Duftschmid, 1812) | Carabidae |
| | | <i>Acidota cruentata</i> Mannerheim, 1830 | Staphylinidae |
| | | <i>Agathidium seminulum</i> (Linnaeus, 1758) | Leiodidae |
| | | <i>Amphicyllis globus</i> (Fabricius, 1792) | Leiodidae |
| | | <i>Anisotoma humeralis</i> (J.F.W. Herbst, 1791) | Leiodidae |
| | | <i>Anoplotrupes stercorosus</i> (Scriba, 1791) | Geotrupidae |
| | | <i>Anotylus inustus</i> (Gravenhorst, 1806) | Staphylinidae |
| | | <i>Anotylus sculpturatus</i> (Gravenhorst, 1806) | Staphylinidae |
| | | <i>Atheta crassicornis</i> (Fabricius, 1793) | Staphylinidae |
| | | <i>Atheta fungi</i> (Gravenhorst, 1806) | Staphylinidae |
| | | <i>Atheta gyllenhalii</i> (Thomson, 1856) | Staphylinidae |
| | | <i>Atheta ravilla</i> (Erichson, 1839) | Staphylinidae |
| | | <i>Athous vittatus</i> (Fabricius, 1792) | Elateridae |
| | | <i>Autalia longicornis</i> Scheerpeltz, 1947 | Staphylinidae |
| | | <i>Brachysomus echinatus</i> (Bonsdorff, 1785) | Curculionidae |
| | | <i>Bryaxis puncticollis</i> (Denny, 1825) | Staphylinidae |
| | | <i>Bryophacis crassicornis</i> (Mäklin, 1847) | Staphylinidae |
| | | <i>Bythinus macropalpus</i> Aubé, 1833 | Staphylinidae |

A list of sampled species with their ecological classification

| | | |
|--|--|----------------|
| | <i>Calathus micropterus</i> (C. Duftschmid, 1812) | Carabidae |
| | <i>Carabus arvensis</i> Herbst, 1784 | Carabidae |
| | <i>Carabus coriaceus</i> Linnaeus, 1758 | Carabidae |
| | <i>Cartodere nodifer</i> (J.O. Westwood, 1839) | Lathridiidae |
| | <i>Colenis immunda</i> (J. Sturm, 1807) | Leiodidae |
| | <i>Cryptophagus montanus</i> Ch.N.F. Brisout de Barneville, 1863 | Cryptophagidae |
| | <i>Cryptophagus pallidus</i> J. Sturm, 1845 | Cryptophagidae |
| | <i>Cryptophagus punctipennis</i> Ch.N.F. Brisout de Barneville, 1863 | Cryptophagidae |
| | <i>Cryptophagus reflexus</i> C. Rey, 1889 | Cryptophagidae |
| | <i>Cryptophagus saginatus</i> J. Sturm, 1845 | Cryptophagidae |
| | <i>Dinaraea angustula</i> (Gyllenhal, 1810) | Staphylinidae |
| | <i>Dinothenarus fossor</i> (Scopoli, 1771) | Staphylinidae |
| | <i>Domene scabricollis</i> (Erichson, 1840) | Staphylinidae |
| | <i>Euconnus</i> sp. C.G. Thomson, 1859 | Scydmaenidae |
| | <i>Gabrius breviventer</i> (Sperk, 1835) | Staphylinidae |
| | <i>Ischnosoma longicorne</i> (Mäklin, 1847) | Staphylinidae |
| | <i>Leiodes</i> sp. Latreille, 1797 | Leiodidae |
| | <i>Leistus ferrugineus</i> (Linnaeus, 1758) | Carabidae |
| | <i>Leistus rufomarginatus</i> (C. Duftschmid, 1812) | Carabidae |
| | <i>Lesteva longoelytrata</i> (Goeze, 1777) | Staphylinidae |
| | <i>Liogluta alpestris</i> (Heer, 1839) | Staphylinidae |
| | <i>Megasternum concinuum</i> (Th. Marsham, 1802) | Hydrophilidae |
| | <i>Molops piceus</i> (Panzer, 1793) | Carabidae |
| | <i>Mycetoporus clavicornis</i> (Stephens, 1832) | Staphylinidae |
| | <i>Ocalea picata</i> (Stephens, 1832) | Staphylinidae |
| | <i>Ocypus macrocephalus</i> (Gravenhorst, 1802) | Staphylinidae |
| | <i>Onthophilus punctatus</i> (O.F. Müller, 1776) | Histeridae |
| | <i>Otiorhynchus scaber</i> (Linnaeus, 1758) | Curculionidae |
| | <i>Oxypoda acuminata</i> (Stephens, 1832) | Staphylinidae |
| | <i>Oxypoda annularis</i> (Mannerheim, 1830) | Staphylinidae |

A list of sampled species with their ecological classification

| | | | |
|--|--|--|---------------|
| | | <i>Oxypoda praecox</i> Erichson, 1839 | Staphylinidae |
| | | <i>Philonthus decorus</i> (Gravenhorst, 1802) | Staphylinidae |
| | | <i>Phloeonomus punctipennis</i> Thomson, 1867 | Staphylinidae |
| | | <i>Proteinus brachypterus</i> (Fabricius, 1792) | Staphylinidae |
| | | <i>Pterostichus melanarius</i> (J.K.W. Illiger, 1798) | Carabidae |
| | | <i>Pterostichus niger</i> (Schaller, 1783) | Carabidae |
| | | <i>Pterostichus oblongopunctatus</i> (Fabricius, 1787) | Carabidae |
| | | <i>Ptinus fur</i> (Linnaeus, 1758) | Ptinidae |
| | | <i>Rhagonycha gallica</i> Pic, 1923 | Cantharidae |
| | | <i>Rhizophagus bipustulatus</i> (Fabricius, 1792) | Monotomidae |
| | | <i>Sepedophilus marshami</i> (Stephens, 1832) | Staphylinidae |
| | | <i>Serica brunnea</i> (Linnaeus, 1758) | Scarabaeidae |
| | | <i>Staphylinus erythropterus</i> Linnaeus, 1758 | Staphylinidae |
| | | <i>Stenichnus</i> sp. C.G. Thomson, 1859 | Scydmenidae |
| | | <i>Xantholinus laevigatus</i> Jacobsen, 1849 | Staphylinidae |
| | | <i>Xantholinus linearis</i> (Olivier, 1795) | Staphylinidae |
| | | <i>Xantholinus longiventris</i> Heer, 1839 | Staphylinidae |