

Review

## The Fungal Defensin Family Enlarged

Jiajia Wu, Bin Gao and Shunyi Zhu \*

Group of Animal Innate Immunity, State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute of Zoology, Chinese Academy of Sciences, 1 Beichen West Road, Chaoyang District, Beijing 100101, China

\* Author to whom correspondence should be addressed; E-Mail: Zhusy@ioz.ac.cn.

Received: 5 June 2014; in revised form: 5 August 2014 / Accepted: 8 August 2014 /

Published: 18 August 2014

---

**Abstract:** Fungi are an emerging source of peptide antibiotics. With the availability of a large number of model fungal genome sequences, we can expect that more and more fungal defensin-like peptides (fDLPs) will be discovered by sequence similarity search. Here, we report a total of 69 new fDLPs encoded by 63 genes, in which a group of fDLPs derived from dermatophytes are defined as a new family (fDEF8) according to sequence and phylogenetic analyses. In the oleaginous fungus *Mortierella alpine*, fDLPs have undergone extensive gene expansion. Our work further enlarges the fungal defensin family and will help characterize new peptide antibiotics with therapeutic potential.

**Keywords:** peptide antibiotic; gene duplication; exon-intron structure; cysteine-stabilized  $\alpha$ -helical and  $\beta$ -sheet motif

---

### 1. Introduction


Fungal defensin-like peptides (fDLPs) are emerging as attractive anti-infective agents due to their therapeutic efficacy, low toxicity and high serum stability [1,2]. On the basis of a combined analyses of sequence, structural, and phylogenetic data, we have identified seven fDLP families [2,3], in which three members (plectasin, micasin and eurocin), classified as ancient invertebrate-type defensins (AITDs) [1,2,4,5], have been structurally and functionally characterized. These fDLPs exhibit activity against several antibiotic-resistant clinical isolates with significant therapeutic potential [1,2,5,6]. Some efforts have been taken to improve antimicrobial efficacy and to reduce undesirable side effects of fDLPs. For example, an improved mutant of plectasin (NZ2114) is superior to two conventional antibiotics (vancomycin and daptomycin) in inhibiting methicillin-resistant *Staphylococcus aureus*

(MRSA) with even more enhanced serum stability and extended *in vivo* half-life [7–9]. In this work, we describe 69 new fDLPs in terms of their sequences, structural characteristics, and phylogenetic relationship. This provides an array of candidates for development of new anti-infective agents against antibiotic-resistant human pathogens.

## 2. Discovery of New fDLPs

The database search strategy used here has been described previously [3]. Through an exhaustive search of 26 fungal species, we retrieved a total of 69 new fDLPs. As previously stated, overall this class of molecules exhibits a taxa-specific distribution pattern in the fungus kingdom, of which 21 fDLPs are derived from *Ascomycota*, 39 from *Zygomycota*, eight from *Basidiomycota* and one from *Glomeromycota*. In the basal fungi (*Microsporidia* and *Chytridiomycota*), no typical fDLP has been identified (Figure 1). The general features of these peptides are listed in Tables 1 and 2. They can be grouped into six families based on sequence similarity, five of which are classified into the previously known families (fDEF1, fDEF2, fDEF3, fDEF4, and fDEF6) [3] (Figures 2 and 3). This grouping is consistent with the phylogenetic analysis supported by high bootstrap values (Figure 4).

**Figure 1.** Phylogenetic distribution of fDLPs. The left: A parsimony tree of fungal species, animalia is used as an outgroup. This tree is a modification of the SSU and LSU r-RNA analyses of Lutzoni *et al.* for the fungal kingdom [10]. The right: “+” means presence and “–” means absence.



All the fDLPs characterized here have a signal peptide located in the N-terminus. In comparison with fDEF1 and fDEF2 that possess a propeptide located between signal and mature peptides, fDEF6 and fDEF8 lack a propeptide. Five precursors (maglosin, beauvesin2, manisin, pochlasin2 and asosin) could release two defensins from a single precursor after the removal of a spacer propeptide (Figure 5). The malpisin family from *Mortierella alpine* exhibits two types of precursor organization: (1) the first type contains 10 members, all having a propeptides identified by its acidic feature and single or two basic amino acids at their ends as putative cleavage site of proprotein convertase [11]; (2) the second type contains 14 members that lack a propeptide and thus no further processing step is needed (Figure S1).

**Table 1.** Sources and characteristics of newly discovered non-*Mortierella* fDLPs.

| Name Accession No.           | Class  | Species (phylum: subphylum: class)  | Size  | MW   | NC   |
|------------------------------|--|---|---|------|------|
| Pyronesin1 CATG01000243 (G)  | fDEF1  | <i>Pyronema omphalodes</i> (Ascomycota: Pezizomycotina: Pezizomycetes)          | 40  | 4317 | +1.2 |
| Pyronesin2 CATG01000243 (G)  |  | <i>P. omphalodes</i>  | 40  | 4402 | +0.2 |
| Pyronesin3 CATG01000243 (G)  |  | <i>P. omphalodes</i>  | 40  | 4389 | +1.2 |
| Pyronesin4 CATG01000243 (G)  |  | <i>P. omphalodes</i>  | 40  | 4416 | +2.2 |
| Pyronesin5 CATG01000243 (G)  |  | <i>P. omphalodes</i>  | 40  | 4375 | +1.2 |
| Pyronesin6 CATG01000243 (G)  |  | <i>P. omphalodes</i>  | 40  | 4291 | +2.4 |
| Abisin1 AEOK01000166 (G)     |  | <i>Agaricus bisporus</i> (Basidiomycota: Agaricomycotina: Agaricomycetes)       | 40  | 4097 | -3.8 |
| Abisin2 AEOK01000166 (G)     |  | <i>A. bisporus</i>  | 40  | 4097 | -3.8 |
| Abisin3 AEOK01000166         |  | <i>A. bisporus</i>  | 39  | 3926 | -2.8 |
| Beauvesin1 ADAH01000714 (G)  |  | <i>Beauveria bassiana</i> (Ascomycota: Pezizomycotina: Sordariomycetes)         | 52  | 5475 | +2.9 |
| Pyrelysin GAJI01023341 (T)   |  | <i>Pyrenochaeta lycopersici</i> (Ascomycota: Pezizomycotina: Dothideomycetes)   | 55  | 5858 | +5.4 |
| Risin JAQX01005622           |  | <i>Rhizophagus irregularis</i> (Glomeromycota: Glomeromycetes)                  | 55  | 5972 | +6.1 |
| Trimensin FG132536 (E)       |  | <i>Trichophyton mentagrophytes</i> (Ascomycota: Pezizomycotina: Eurotiomycetes) | 38  | 4156 | +2.2 |
| Lecasin AWYC01000479         |  | fDEF2   | <i>Lecanosticta acicola</i> (Ascomycota: Pezizomycotina: Dothideomycetes)   | 42   | 4314 |
| Pochlasin1 AOSW01002431      | <i>Pochonia chlamydosporia</i> (Ascomycota: Pezizomycotina: Sordariomycetes)         |   | 43  | 4339 | -3.5 |
| Perisin AFRD01000258         | <i>Periglandula ipomoeae</i> (Ascomycota: Pezizomycotina: Sordariomycetes)           |   | 43  | 4080 | -1.5 |
| Masysin CANK01000016         | <i>Malassezia sympodialis</i> (Basidiomycota: Ustilaginomycotina: Exobasidiomycetes) |   | 35  | 3432 | +2.2 |
| Maglosin1N AAYY01000039 (G)  | <i>Malassezia globosa</i> (Basidiomycota: Ustilaginomycotina: Exobasidiomycetes)     |   | 40  | 3980 | +1.2 |
| Maglosin2N AAYY01000024 (G)  | <i>M. globosa</i>  |   | 40  | 4022 | +0.2 |
| Maglosin1C AAYY01000039 (G)  | <i>M. globosa</i>  |   | 41  | 3910 | +2.7 |
| Maglosin2C AAYY01000024 (G)  | <i>M. globosa</i>  |   | 40  | 3835 | +2.7 |
| Beauvesin2C ADAH01000123 (G) | <i>B. bassiana</i>   |   | 41  | 4243 | +0.9 |
| ManisinC ADNJ01000735        | fDEF3  |   | <i>Metarhizium anisopliae</i> (Ascomycota: Pezizomycotina: Sordariomycetes) | 41   | 4211 |
| Pochlasin2C AOSW01005877     |  | <i>P. chlamydosporia</i>  | 41  | 4381 | +0.2 |
| AsosinC BACA01000303         |  | <i>Aspergillus sojae</i> (Ascomycota: Pezizomycotina: Eurotiomycetes)           | 38  | 4002 | -1.0 |

Table 1. Cont.


| Name Accession No.           | Class   | Species (phylum: subphylum: class)                                      | Size   | MW   | NC    |
|------------------------------|---|---|--|------|-------|
| Beauvesin2N ADAH01000123 (G) | fDEF4   | <i>B. bassiana</i>  | 48   | 5067 | +2.9  |
| ManisinN ADNJ01000735        |   | <i>M. anisopliae</i>  | 46   | 4921 | +0.2  |
| Pochlasin2N AOSW01005877     |   | <i>P. chlamydosporia</i>  | 49   | 5185 | +2.9  |
| AsosinN BACA01000303         |   | <i>A. sojae</i>   | 49   | 5140 | -1.1  |
| Rhimisin1 ANKS01000620       | fDEF6   | <i>Rhizopus microsporus</i> (Zygomycota: Mucoromycotina: Mucorales)     | 45   | 4867 | +10.0 |
| Rhimisin2 ANKS01000620       |   | <i>R. microsporus</i>   | 44   | 4638 | +3.4  |
| Rhimisin3 ANKS01001486       |   | <i>R. microsporus</i>   | 44   | 4768 | +1.5  |
| Rhimisin4 ANKS01001486       |   | <i>R. microsporus</i>   | 45   | 4811 | +8.0  |
| Rhidesin1 AACW02000043       |   | <i>Rhizopus delemar</i> (Zygomycota: Mucoromycotina: Mucorales)         | 55   | 5885 | +10.4 |
| Rhidesin2 AACW02000259       |   | <i>R. delemar</i>   | 48   | 5270 | +0.5  |
| Mirresin AZYI01000143        |   | <i>Mucor irregularis</i> (Zygomycota: Mucoromycotina: Mucorales)        | 60   | 6424 | +13.4 |
| Mucisin AOCY01001156 (G)     |   | <i>Mucor circinelloides</i> (Zygomycota: Mucoromycotina: Mucorales)     | 53   | 5548 | +2.2  |
| Phycomysin EX863311 (E)      |   | <i>Phycomyces blakesleeanus</i> (Zygomycota: Mucoromycotina: Mucorales) | 50   | 5342 | +9.4  |
| TritoDLP ACPI01000196 (G)    |   | fDEF8   | <i>Trichophyton tonsurans</i> (Ascomycota: Pezizomycotina: Eurotiomycetes) | 41   | 4323  |
| TrequiDLP ABWI01000729 (G)   | <i>Trichophyton equinum</i> (Ascomycota: Pezizomycotina: Eurotiomycetes)    |   | 41   | 4323 | +3.7  |
| TriveDLP ACYE01000402        | <i>Trichophyton verrucosum</i> (Ascomycota: Pezizomycotina: Eurotiomycetes) |   | 42   | 4403 | +4.7  |
| ArgyDLP ABQE01000293         | <i>Arthroderma gypseum</i> (Ascomycota: Pezizomycotina: Eurotiomycetes)     |   | 41   | 4247 | +3.7  |
| ArbeDLP ABSU01000004         | <i>Arthroderma benhamiae</i> (Ascomycota: Pezizomycotina: Eurotiomycetes)   |   | 42   | 4493 | +4.7  |
| TriruDLP ACPH01000567 (G)    | <i>Trichophyton rubrum</i> (Ascomycota: Pezizomycotina: Eurotiomycetes)     |   | 42   | 4479 | +4.7  |
| MicaDLP ABVF01000093         | <i>Arthroderma otae</i> (Ascomycota: Pezizomycotina: Eurotiomycetes)        |   | 43   | 4745 | +3.2  |

Note: MW: molecular weight; NC (net charge) is estimated at pH 7.0 with protein calculation V3.4. “E” means peptides from the Expressed Sequence Tags (EST) database and “T” means peptides from the Transcriptome Shotgun Assembly (TSA) database. “G” means proteins currently annotated in the GenBank database as hypothetical proteins (<http://www.ncbi.nlm.nih.gov/>) [12].


**Table 2.** Sources and characteristics of the malpisin family.

| Name Accession No.        | Organism                             | Scaffold (Contig)                | Range       | Size   | MW            | NC   |      |      |
|---------------------------|--------------------------------------|----------------------------------|-------------|--|---------------|------|------|------|
| Malpisin1-1 AZCI01001104  | <i>Mortierella alpina</i> B6842      | jtg7180000084593f_7180000084594f | 55070–55405 | 41   | 4048          | –0.0 |      |      |
| Malpisin1-2 AZCI01001104  |                                      |                                  | 55870–56127 | 48   | 5166          | –3.3 |      |      |
| Malpisin1-3 AZCI01001104  |                                      |                                  | 56393–56635 | 45   | 5047          | +0.7 |      |      |
| Malpisin1-4 AZCI01001104  |                                      |                                  | 63869–64117 | 39   | 4117          | –1.0 |      |      |
| Malpisin1-5 AZCI01000882  |                                      | Contig 7180000084767             |             | 22045–22248  | 37            | 4259 | –2.8 |      |
| Malpisin1-6 AZCI01000882  |                                      |                                  |             | 25851–26051  | 39            | 4543 | –3.5 |      |
| Malpisin1-7 AZCI01000882  |                                      |                                  |             | 42456–42677  | 33            | 3624 | –1.0 |      |
| Malpisin1-8 AZCI01000882  |                                      |                                  |             | 43573–43800  | 47            | 5078 | +1.7 |      |
| Malpisin1-9 AZCI01000882  |                                      |                                  |             | 45037–45261  | 48            | 5203 | +3.0 |      |
| Malpisin1-10 AZCI01000882 |                                      |                                  |             | 45559–45738  | 35            | 3914 | –0.0 |      |
| Malpisin1-11 AZCI01000882 |                                      |                                  |             | 46707–46913  | 43            | 4941 | +5.4 |      |
| Malpisin1-12 AZCI01001135 |                                      |                                  |             | jtg7180000084204f_7180000084205f_7180000084206f                | 135437–135676 | 44   | 4722 | –0.0 |
| Malpisin1-13 AZCI01001084 |                                      |                                  |             | jtg7180000084699f_7180000084700f                               | 362415–362627 | 47   | 4919 | –1.8 |
| Malpisin1-14 AZCI01001006 |                                      |                                  |             | jtg7180000084769f_7180000084770f_7180000084771f_7180000084772f | 179488–179673 | 38   | 4188 | +0.2 |
| Malpisin2-1 ADAG01001070  | <i>Mortierella alpina</i> ATCC 32222 | Contig 1070                      | 9785–10114  | 39   | 4105          | +1.0 |      |      |
| Malpisin2-2 ADAG01001070  |                                      |                                  | 10532–10792 | 48   | 5187          | –2.3 |      |      |
| Malpisin2-3 ADAG01001070  |                                      |                                  | 11052–11297 | 44   | 4783          | –0.0 |      |      |
| Malpisin2-4 ADAG01001070  |                                      |                                  | 11773–12021 | 39   | 4052          | –1.8 |      |      |
| Malpisin2-5 ADAG01000791  |                                      | Contig 791                       | 4894–5097   | 37   | 4259          | –2.8 |      |      |
| Malpisin2-7 ADAG01000903  |                                      | Contig 903                       | 13145–13357 | 33   | 3899          | +1.2 |      |      |
| Malpisin2-8 ADAG01000903  |                                      |                                  | 14223–14450 | 47   | 5065          | +1.7 |      |      |
| Malpisin2-9 ADAG01000903  |                                      |                                  | 15634–15852 | 45   | 4917          | +4.0 |      |      |
| Malpisin2-10 ADAG01000903 |                                      |                                  | 16158–16337 | 35   | 3937          | +0.2 |      |      |
| Malpisin2-11 ADAG01000903 |                                      |                                  | 17264–17446 | 39   | 4531          | +5.0 |      |      |

**Figure 2.** Multiple sequence alignment of fDLPs. Cysteines are shadowed in cyan. Conserved glycines are highlighted in grey. Negatively (D and E) and positively (R, K and H) charged residues are boldfaced in red and blue, respectively. Introns are shown by arrows (phase 0) or small boxes (green: phase 1, yellow: phase 2). Functionally characterized fDLPs were indicated by “\*”. The N-terminal extension sequence in micaDLP belonging to the family fDEF8 is italicized. Defensins from *Pyronema omphalodes* have been predicted and investigated by RNA-seq [13]. Extra residues for C-terminal amidation are underlined once.




**Figure 3.** Multiple sequence alignment of malpispins. Color codes and symbol notes used here are the same as those in Figure 2. Pink box indicates the N-terminus of DLPs with variable length. Sequence identity (%) to micasin is shown on the right.



Peptides in fDEF8, all derived from dermatophytes, are characterized as a new family with a short N-terminus and an extra C-terminal extension rich in arginines, prolines and glycines (Figure 2). The C-terminal extension has been considered as a common mechanism for the complexity increase of some invertebrate antimicrobial peptides (AMPs). For example, the hymenopteran defensin-1 subfamily has an extended C-terminus relative to its ancestral defensin-2 subfamily by a so-called intron exonization-mediated mechanism [14,15]. It thus appears that fungal and invertebrate defensins both convergently evolved their C-termini. The extension of a C-terminal sequence via convergent evolution was also recently observed in interleukin 6 (IL-6), a class-I helical cytokine, of two leporids (*Oryctolagus* and *Pentalagus*) [16]. The presence of C-terminal Gly-Arg or Gly-Arg-Arg in some dermatophyte-derived fDLPs suggest that they may be amidated, as previously observed in some animal toxins, e.g., the *Mesobuthus*  $\alpha$ -toxins [17]. Interestingly, the mature peptide of micaDLP is larger in size than that of other members in this family, as identified by an N-terminal extension of 38 amino acids (Figure 2). High content of glycines together with a cationic characteristic hints a putative antimicrobial role of this extended unit.



**Figure 5.** Comparison of precursor organization and exon-intron structures between fDLPs and animal defensins. (A) fDLPs; (B) Animal defensins. Signal, pro- and mature peptides are shown in pink, grey and blue, respectively. Intron phases are shown in the same colors as Figure 2. Representative animal defensins are derived from *Branchiostoma floridae*, *Drosophila melanogaster*, *Anopheles arabiensis*, *Apis mellifera*, *Ixodes scapularis*, *M. martensii*, *Crassostrea gigas*, *Caenorhabditis remanei*, and *C. brenneri*.



*M. alpine* is a saprophytic species of *Mucoromycotina*, known as an oleaginous fungus [18]. The draft genome sequences of two *M. alpina* isolates (B6842 and ATCC 32222) [18,19] provide a possibility to undertake comparative study of their fDLPs. We found that the *M. alpine* B6842 genome encodes 14 fDLPs (Figure 3) but only 10 were found in *M. alpine* ATCC 32222. The failure to detect the four homologs (*i.e.*, malpisin1-6, 1-12, 1-13, 1-14) in *M. alpine* ATCC 32222 could be due to the incompletely-assembled genome sequences. Our phylogenetic analysis divides all malpisisins into fDEF1 and fDEF6 (Figure 4). Some malpisin members of fDEF1 extended their N-termini with diverse sequences and variable lengths (Figure 3).


### 3. Gene Duplication of fDLPs

Gene duplication extensively occurs in antimicrobial peptides from insects to humans [15,20,21]. In fungi, initial annotation of defense molecules of *Pyronema omphalodes* also identified gene duplication as a minor multigene family of fDLPs (herein termed pyronesin1 to pyronesin6) [13]. These fDLPs are highly similar to plectasin (Figure 6A). Our studies revealed new gene duplication event in other fungal species. Malpisin is a representative example of gene duplication. As mentioned previously, there are 14 and 10 members in *M. alpine* B6842 and *M. alpine* ATCC 32222, respectively. Malpisin1-1, 1-2, 1-3 and 1-4 are tandem located on one contig (jtg7180000084593f\_7180000084594f), and malpisin1-5 to malpisin1-11 on another contig (Contig 7180000084767). In addition, malpisin1-12, malpisin 1-13 and malpisin 1-14 reside on other three contigs, as shown in Figure 6B. In *M. alpine* ATCC 32222, malpisin2-1 to malpisin2-4 are located on contig 1070 and malpisin2-7 to malpisin2-11 on contig 903. Only malpisin2-5 is located on contig 791.

In the widely cultivated mushroom *Agaricus bisporus*, there are three paralogous fDLPs (abisin1 to abisin3) (Figure 6C), two of which (abisin1 and abisin3) share completely identical amino acid sequences in the mature peptide region but exhibit four synonymous substitutions at the nucleotide level. In the *Pochonia chlamydosporia* paralogues, pochlasin1 is highly similar to CITDs and pochlasin2 possesses two defensin-domains. In addition, a putative pseudogene (herein named pochlasin-pseu) was also identified in scaffold 1191 and assigned to AITDs in view of its high sequence similarity to micasin in the first exon. Pochlasin1 and pochlasin-pseu share a conserved phase 0 intron within the  $\alpha$ -helical region. The loss of the last two exons (2 and 3) results in the lack the last four cysteines involved in the Cs $\alpha$  $\beta$  folding of a mature peptide (Figure 2).

Gene duplication also occurs in the *Mucorales*-derived fDLPs, which leads to four and two gene copies in *Rhizopus microsporus* (Figure 6D) and *R. delemar*, respectively. In a Neighbor-Joining (NJ) tree, rhimisin1 and rhimisin4 (*R. microsporus*) constitutes a single clade clustering with the other three fDLPs (rhidesin1 from *R. delemar*, phycomycin from *Phycomyces blakesleeanus* and mirresin from *Mucor irregularis*) whereas rhimisin2 and rhimisin3 (*R. microsporus*) cluster with rhidesin2 (*R. delemar*) and mucisin (*M. circinelloides*) (Figure 4), suggesting that the gene duplication event could have occurred in the ancestor of the *Mucorales* prior to their speciation.

**Figure 6.** The arrangement of defensin genes in chromosomes. Color arrows refer to different orientation of the genes. **A** to **D** represent the genome location of defensins in four species: *Pyronema omphalodes*, *Mortierella alpine*, *Agaricus bisporus* and *Rhizopus microsporus*. Malpisinins in *M. alpine* B6842 is indicated in red and blue while in pink and green in *M. alpine* ATCC 32222. Pseudogenes of pyronesins are shown in gradient blue.



#### 4. Variable Gene Structures of fDLPs

Analysis of the exon-intron structures of the newly-discovered fDLPs revealed their variability that can be described as follows: (1) all the fDLPs retain the integrity of the signal peptide except risin (*Rhizophagus irregularis*) and malpisin1-1 (or malpisin2-1) that have a phase 1 or phase 0 intron disrupting their signal peptides; (2) all of the genes in fDEF8 and three genes in fDEF2 (*i.e.*, lecasin, pochlasin1 and perisin) have the same gene organization as previously identified dermatophytic defensins (micasin, arbesin, trivesin, tritosin and trirusin) and they contain two introns: the first intron (phase 0) disrupting the  $\alpha$ -helical region; the second intron (phase 2) disrupting the c-loop; (3) the pyronesin and abisin multi-gene family in fDEF1 have only one intron disrupting either the  $\alpha$ -helical or the c-loop region; (4) In addition to these intron-containing fDLP genes, there are some members without introns (Figures 2 and 5).

The highly variable gene structures in fDLPs are reminiscent of invertebrate defensins that also exhibit diverse gene structures [22,23] (Figures 5 and S2). Compared with invertebrate defensins of 5'-biased intron positions, introns of fDLPs occur preferentially in the 3'-end of the precursor-coded sequences. Because all eukaryotic C $\alpha$  $\beta$ -type defensins are hypothesized to be originated from a common bacterial ancestor [24], it is reasonable to infer that considerable intron gains might have occurred in defensins from some eukaryotic lineages, and later they differentially lost in some specific species. Such a dynamic intron evolution thus shapes the biased intron location pattern between fDLPs and animal DLPs after the animal-fungi split. It is also worth mentioning that some recognizable orthologues of defensins in *Branchiostoma floridae* [25,26], the basal chordate amphioxus, also contain a phase 0 intron located in their c-loop (Figures 5 and S2). Given a remote evolutionary distance between fungi and amphioxus, their intron position conservation could be a consequence of convergent insertion in a similar position due to the existence of "protosplice sites" [27,28]. However, the evolution via ancestral origin can be not completely ruled out in the case of the lack of gene structure information in many animal defensins from different lineages.

#### 5. Conclusions

It is estimated that there are as many as 1.5 million species of fungi in this world. However, only a small fraction has been described and even fewer have been sequenced. To date, only about six hundred genomes were being sequenced or completely sequenced. Fungal genome project (FGP) allows us to systematically exploit peptide antibiotics instead of accidental discovery or complicated biochemical screening. This work sheds light on the persistent discovery of fDLPs from model fungal genome data. Despite this, in the lack of experimental data, it cannot be stated that all these fDLPs possess antibacterial function because in fact a classical insect-type fungal defensin - pechrysin was found to lack antibacterial activity [29] likely due to the absence of cationic residues on its molecular surface. In addition, anisin1, a DLP from *Aspergillus giganteus*, was found to be involved in the fitness of the species by linking stress signaling with developmental regulation [30]. Recent studies have also shown that although some peptides of fungal origin contain a similar defensin structure, they exhibit diverse or alternative biological functions beyond antimicrobial activity. An interesting

overview is given by Hegedüs and Marx [31]. Therefore, further biochemical characterization of these newly-discovered fDLPs will help evaluate their potential as human medicines.

### Acknowledgements

This work was supported by the National Basic Research Program of China (2010CB945300), the National Natural Science Foundation of China (31221091), and the State Key Laboratory of Integrated Management of Pest Insects and Rodents (Grant No. ChineseIPM1307).

### Author Contributions

J.W. and S.Z. discovered all fDLP genes described here. J.W. and S.Z. wrote the paper. B.G. revised the paper.

### Conflicts of Interest

The authors declare no conflict of interest.

### References

1. Mygind, P.H.; Fischer, R.L.; Schnorr, K.M.; Hansen, M.T.; Sonksen, C.P.; Ludvigsen, S.; Raventos, D.; Buskov, S.; Christensen, B.; De Maria, L.; *et al.* Plectasin is a peptide antibiotic with therapeutic potential from a saprophytic fungus. *Nature* **2005**, *437*, 975–980.
2. Zhu, S.; Gao, B.; Harvey, P.J.; Craik, D.J. Dermatophytic defensin with antiinfective potential. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 8495–8500.
3. Zhu, S. Discovery of six families of fungal defensin-like peptides provides insights into origin and evolution of the CS $\alpha\beta$  defensins. *Mol. Immunol.* **2008**, *45*, 828–838.
4. Schneider, T.; Kruse, T.; Wimmer, R.; Wiedemann, I.; Sass, V.; Pag, U.; Jansen, A.; Nielsen, A.K.; Mygind, P.H.; Raventos, D.S.; *et al.* Plectasin, a fungal defensin, targets the bacterial cell wall precursor Lipid II. *Science* **2010**, *328*, 1168–1172.
5. Oemig, J.S.; Lynggaard, C.; Knudsen, D.H.; Hansen, F.T.; Norgaard, K.D.; Schneider, T.; Vad, B.S.; Sandvang, D.H.; Nielsen, L.A.; Neve, S.; *et al.* Eurocin, a new fungal defensin: Structure, lipid binding, and its mode of action. *J. Biol. Chem.* **2012**, *287*, 42361–42372.
6. Hara, S.; Mukae, H.; Sakamoto, N.; Ishimoto, H.; Amenomori, M.; Fujita, H.; Ishimatsu, Y.; Yanagihara, K.; Kohno, S. Plectasin has antibacterial activity and no affect on cell viability or IL-8 production. *Biochem. Biophys. Res. Commun.* **2008**, *374*, 709–713.
7. Xiong, Y.Q.; Hady, W.A.; Deslandes, A.; Rey, A.; Fraisse, L.; Kristensen, H.H.; Yeaman, M.R.; Bayer, A.S. Efficacy of NZ2114, a novel plectasin-derived cationic antimicrobial peptide antibiotic, in experimental endocarditis due to methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **2011**, *55*, 5325–5330.
8. Ostergaard, C.; Sandvang, D.; Frimodt-Moller, N.; Kristensen, H.H. High cerebrospinal fluid (CSF) penetration and potent bactericidal activity in CSF of NZ2114, a novel plectasin variant, during experimental pneumococcal meningitis. *Antimicrob. Agents Chemother.* **2009**, *53*, 1581–1585.

9. Andes, D.; Craig, W.; Nielsen, L.A.; Kristensen, H.H. *In vivo* pharmacodynamic characterization of a novel plectasin antibiotic, NZ2114, in a murine infection model. *Antimicrob. Agents Chemother.* **2009**, *53*, 3003–3009.
10. Lutzoni, F.; Kauff, F.; Cox, C.J.; McLaughlin, D.; Celio, G.; Dentinger, B.; Padamsee, M.; Hibbett, D.; James, T.Y.; Baloch, E.; *et al.* Assembling the fungal tree of life: Progress, classification, and evolution of subcellular traits. *Am. J. Bot.* **2004**, *91*, 1446–1480.
11. Seidah, N.G.; Chrétien, M. Proprotein and prohormone convertases: A family of subtilases generating diverse bioactive polypeptides. *Brain Res.* **1999**, *848*, 45–62.
12. Benson, D.A.; Karsch-Mizrachi, I.; Lipman, D.J.; Ostell, J.; Wheeler, D.L. GenBank. *Nucleic Acids Res.* **2008**, *36*, D25–30.
13. Traeger, S.; Altegoer, F.; Freitag, M.; Gabaldon, T.; Kempken, F.; Kumar, A.; Marcet-Houben, M.; Poggeler, S.; Stajich, J.E.; Nowrousian, M. The genome and development-dependent transcriptomes of *Pyronema confluens*: A window into fungal evolution. *PLoS Genet.* **2013**, *9*, e1003820.
14. Tian, C.; Gao, B.; Fang, Q.; Ye, G.; Zhu, S. Antimicrobial peptide-like genes in *Nasonia vitripennis*: A genomic perspective. *BMC Genomics* **2010**, *11*, 187.
15. Zhang, Z.; Zhu, S. Comparative genomics analysis of five families of antimicrobial peptide-like genes in seven ant species. *Dev. Comp. Immunol.* **2012**, *38*, 262–274.
16. Neves, F.; Abrantes, J.; Pinheiro, A.; Almeida, T.; Costa, P.P.; Esteves, P.J. Convergent evolution of IL-6 in two leporids (*Oryctolagus* and *Pentalagus*) originated an extended protein. *Immunogenetics* **2014**, doi:10.1007/s00251-014-0787-0.
17. Zhu, S.; Peigneur, S.; Gao, B.; Lu, X.; Cao, C.; Tytgat, J. Evolutionary diversification of *Mesobuthus*  $\alpha$ -scorpion toxins affecting sodium channels. *Mol. Cell. Proteomics* **2012**, doi:10.1074/mcp.M111.012054.
18. Wang, L.; Chen, W.; Feng, Y.; Ren, Y.; Gu, Z.; Chen, H.; Wang, H.; Thomas, M.J.; Zhang, B.; Berquin, I.M.; *et al.* Genome characterization of the oleaginous fungus *Mortierella alpina*. *PLoS ONE* **2011**, *6*, e28319.
19. Etienne, K.A.; Chibucos, M.C.; Su, Q.; Orvis, J.; Daugherty, S.; Ott, S.; Sengamalay, N.A.; Fraser, C.M.; Lockhart, S.R.; Bruno, V.M. Draft genome sequence of *Mortierella alpina* isolate CDC-B6842. *Genome Announc.* **2014**, *2*, e01180–13.
20. Schutte, B.C.; Mitros, J.P.; Bartlett, J.A.; Walters, J.D.; Jia, H.P.; Welsh, M.J.; Casavant, T.L.; McCray, P.B., Jr. Discovery of five conserved  $\beta$ -defensin gene clusters using a computational search strategy. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 2129–2133.
21. Semple, C.A.; Rolfe, M.; Dorin, J.R. Duplication and selection in the evolution of primate  $\beta$ -defensin genes. *Genome Biol.* **2003**, *4*, R31.
22. Froy, O.; Gurevitz, M. Arthropod and mollusk defensins - evolution by exon-shuffling. *Trends Genet.* **2003**, *19*, 684–687.
23. Rodriguez de la Vega, R.C.; Possani, L.D. On the evolution of invertebrate defensins. *Trends Genet.* **2005**, *21*, 330–332.
24. Gao, B.; Rodriguez Mdel, C.; Lanz-Mendoza, H.; Zhu, S. AdDLP, a bacterial defensin-like peptide, exhibits anti-*Plasmodium* activity. *Biochem. Biophys. Res. Commun.* **2009**, *387*, 393–398.

25. Yu, J.K.; Wang, M.C.; Shin-I, T.; Kohara, Y.; Holland, L.; Satoh, N.; Satou, Y. A cDNA resource for the cephalochordate amphioxus *Branchiostoma floridae*. *Dev. Genes Evol.* **2008**, *218*, 723–727.
26. Zhu, S.; Peigneur, S.; Gao, B.; Umetsu, Y.; Ohki, S.; Tytgat, J. Experimental conversion of a defensin into a neurotoxin: implications for origin of toxic function. *Mol. Biol. Evol.* **2014**, *31*, 546–559.
27. Sverdlov, A.V.; Rogozin, I.B.; Babenko, V.N.; Koonin, E.V. Reconstruction of ancestral protosplice sites. *Curr. Biol.* **2004**, *14*, 1505–1508.
28. Rogozin, I.B.; Sverdlov, A.V.; Babenko, V.N.; Koonin, E.V. Analysis of evolution of exon-intron structure of eukaryotic genes. *Brief. Bioinformatics* **2005**, *6*, 118–134.
29. Wu, Y.; Gao, B.; Zhu, S. Fungal defensins, an emerging source of anti-infective drugs. *Chin. Sci. Bull.* **2014**, *59*, 931–935.
30. Eigentler, A.; Pocsí, I.; Marx, F. The anisin1 gene encodes a defensin-like protein and supports the fitness of *Aspergillus nidulans*. *Arch. Microbiol.* **2012**, *194*, 427–437.
31. Hegedus, N.; Marx, F. Antifungal proteins: more than antimicrobials? *Fungal Biol. Rev.* **2013**, *26*, 132–145.

© 2014 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).