

Antioxidative activity of *Ginkgo*, *Echinacea*, and *Ginseng* tinctures

Rūta Masteikova, Jan Muselik, Jurga Bernatoniene¹, Rūta Bernatoniene²

Department of Pharmaceutics, University of Veterinary and Pharmaceutical Sciences, Czech Republic,

¹Department of Drugs Technology and Social Pharmacy, ²Department of Pharmaceutical Chemistry and Pharmacognosy, Faculty of Pharmacy, Kaunas University of Medicine, Lithuania

Key words: antioxidant activity; *Ginkgo*; *Echinacea*; *Ginseng*.

Summary. The aim of this study was to determine the amount of phenol compounds in tinctures prepared from *Ginkgo* leaves, *Echinacea* plant, and *Ginseng* roots and to evaluate the antioxidative activity of these preparations. We studied the antioxidative activity using the standard 2,2-diphenyl-1-picrylhydrazyl (DPPH•) radical cation scavenging and tyrosine nitration inhibition tests. The obtained findings showed that the amount of phenol compounds in the studied tinctures differed and ranged between 114 to 340±29 gallic acid equivalents (GAE) mg/100 mL. We found that the amount of phenol compounds in *Ginkgo* tincture was statistically significantly greater than that in *Echinacea* or *Ginseng* tinctures. The effectiveness of *Ginkgo* tincture was by 52.7% ($P<0.01$) lower (from 1343±11 µmol catechin/100 mL solution to 637±64 catechin/100 mL solution), compared to *Echinacea* tincture. *Ginseng* tincture was the weakest scavenger of free radicals – only 8±1 µmol catechin/100 mL solution. The inhibition of tyrosine nitration was by 34% ($P<0.01$) greater in *Echinacea* tincture, compared to *Ginkgo* tincture (from 892±36 µmol catechin/100 mL solution to 588±17 µmol catechin/100 mL solution). *Ginseng* tincture was the weakest inhibitor of tyrosine nitration – only 20±8 µmol catechin/100 mL solution, which was by 44.6 times less, compared to *Echinacea* tincture.

Tests on DPPH• radical cation scavenging and inhibition of nitration showed that the antioxidative activity of *Echinacea* tincture was statistically significantly greater compared to that of *Ginkgo* or *Ginseng* tinctures. This allows us to conclude that antioxidative activity is determined not only by phenol compounds, but also by a complex of other components of medicinal raw material.

Introduction

The cause of coronary heart diseases, cancers, and aging processes is oxidative stress that is caused by free radicals (1, 2). One of the ways to prevent the aforementioned pathologies is application of antioxidants. Antioxidants are compounds that protect cells against the damaging effects of reactive oxygen species (3). Experimental studies were performed on three plants that have antioxidative properties – *Ginkgo biloba*, *Panax ginseng*, and *Echinacea purpurea* (4). *Ginkgo biloba* preparations are used in the treatment of Parkinson's and Alzheimer's diseases caused by oxidative stress. Studies have shown that oxidative stress is related to cancer, aging, atherosclerosis, ischemic injury, inflammation, and neurodegenerative diseases (5). Studies by Bulgarian researchers showed that *Panax ginseng* preparations had antioxidant activity and were used for chemotherapy prophylaxis (6).

In the treatment of cancer, the human organism has to deal with cancer-combative strategies such as chemotherapy and radiation, which impede the immune defenses (7). Preparations of *Echinacea* are widely used to stimulate the immune system (8) and as alternative remedies for the prevention of common cold and infections in the upper respiratory tract (9, 10). It has been found that *Echinacea* preparations have antioxidative (4) and free radical scavenging properties (11).

The aim of this study was to determine the amount of phenol compounds in tinctures prepared from *Ginkgo* leaves, *Echinacea* plant, and *Ginseng* roots and to evaluate the antioxidative activity of these preparations.

Materials and methods

Reagents and samples

The chemicals and samples were available commercially and were used as received: tyrosine (BDH

Chemicals, England), 3-nitrotyrosine (Aldrich), DPPH (Sigma), (+)-catechin, gallic acid, and Folin-Ciocalteu reagent (Fluka). All other chemicals used were of analytical grade.

Peroxynitrite solution was prepared according to methods indicated in literature (12). Aliquots of 0.6 M H_2O_2 (in 0.5 M HClO_4) and 0.5 M NaNO_2 (in water) (10 mL each) were pre-cooled to -2°C and mixed rapidly. The reaction was immediately quenched by addition of 5 mL -2°C cold 3.5 M NaOH. Non-reacted H_2O_2 was removed by treatment with excess of MnO_2 , and the solution was filtered. The ONOO concentrations were determined spectrophotometrically in alkaline solution using $\epsilon_{302}=1670 \text{ M}^{-1}\cdot\text{cm}^{-1}$. Stock solutions were stored frozen at -80°C for several months without noticeable decomposition.

Equipment

High performance liquid chromatography (HPLC) system HP 1100 (Agilent Technologies) consisted of a quaternary pump and an autosampler, and DAD detector was used for HPLC separation; UV spectra were measured using HP 8453 (Agilent Technologies).

Inhibition of tyrosine nitration

From the 2.5 mM peroxynitrite solution in 0.05 M NaOH, a volume of 8 mL was drawn and mixed rapidly in the injector of HPLC autosampler with 42 μL 1.0 mM tyrosine solution in 0.15 M KH_2PO_4 – Na_2HPO_4 buffer (pH 7.4) containing suitably diluted plant tincture. The reaction mixture was injected directly onto the HPLC column (Supelcosil ABZ+Plus 250 \times 4.6 mm, 5 μL); the mobile phase consisted of 90% 40 mM HCOOH and 10% CH_3CN (v/v); flow rate was 1 mL/min. The chromatograms were measured at 276 nm. The activity of the tested compounds was calculated relative to the measured peak of 3-nitrotyrosine of control measurement. Quantification was done on the basis of the standard curve of catechin. Results were expressed as micromoles of catechin equivalent per 100 mL of tincture.

Measurement of DPPH quenching activity

Solution (25 μL) of plant tincture suitably dissolved in methanol was adjusted to the volume of 2.0 mL with 0.1 mM DPPH solution in methanol, and the absorbance was measured at 517 nm for 5 min after mixing. The activity of the tested compounds was calculated relative to the measured absorbance of the control measurement. Quantification was done on the basis of the standard curve of catechin. Results were expressed as micromoles of catechin equivalent per 100 mL of tincture.

Statistical analysis

Statistical analysis was performed using statistical software package Statistica 5.5. The data were presented as means \pm S.E.M. Statistical analysis was performed using Student's *t* test, and $P<0.05$ was used as the level of significance.

Results and discussion

Experimental studies were performed on three tinctures: *Ginkgo* 1:5 (extrahent 70% ethanol), *Ginseng* 1:5 (extrahent 70% ethanol), and *Echinacea* 1:5 (extrahent 50% ethanol) (7, 13). According to literature data, phenol compounds in plants are the largest group of compounds with antioxidative properties (10, 14). For this reason we found it expedient to determine the total amount of phenol compounds in the investigated tinctures (Fig. 1). The total amount of phenol compounds was determined using the regression equation of the calibration curve: $y=10.738x+0.061$, $R^2=0.98$. The obtained findings showed that the amount of phenol compounds in the studied tinctures differed and ranged between 11 ± 4 to 340 ± 29 GAE mg/100 mL.

Further on, we studied the antioxidative activity of *Ginkgo*, *Echinacea*, and *Ginseng* tinctures, using the standard DPPH radical cation scavenging and tyrosine nitration inhibition tests. Total antioxidant capacity assay, such as DPPH method, is most common for antioxidant activity for large-scale examination (1). We found that an antioxidant releases hydrogen, thus inactivating free radicals that consequently become stable compounds of DPPH-H type (1). Our findings showed that *Echinacea* tincture was most effective in scavenging DPPH radicals (Fig. 2). The effectiveness of *Ginkgo* tincture was by 52.7% ($P<0.01$) lower (from 1343 ± 11 μmol catechin/100 mL solution to 637 ± 64 μmol catechin/100 μL solution) compared to *Echinacea* tincture. *Ginseng* tincture was the




Fig. 1. The amount of phenol compounds (GAE mg/100 mL) in the investigated tinctures (n=5)




Fig. 2. Antioxidative activity of *Echinacea*, *Ginkgo*, and *Ginseng* tinctures, determined using DPPH and tyrosine nitration techniques (n=6)

*P<0.01 vs. *Ginseng* tinctures; #P<0.01 vs. *Ginkgo* tinctures.

weakest scavenger of free radicals – only 8 ± 1 μmol catechin/100 mL solution.

Peroxynitrite is a potent nitrating and oxidizing agent that is formed by a rapid reaction of nitric oxide with superoxide anion. Nitric oxide is a simple inorganic radical exhibiting diverse physiological functions, including the regulation of neurotransmission and vascular tone (12, 15). Peroxynitrite, a biological toxin, produced *in vivo* by the nearly diffusion-controlled reaction of nitrogen monoxide with superoxide, can nitrate and oxidize various biomolecules, such as thiols, lipids, carbohydrates, and nucleic acids (16). The inhibition of tyrosine nitration was by 34% ($P < 0.01$) greater in *Echinacea* tincture, compared to *Ginkgo* tincture (from 892 ± 36 μmol catechin/100 mL solution to 588 ± 17 μmol catechin/100 mL solution). *Ginseng* tincture was the weakest inhibitor of tyrosine nitration – only 20 ± 8 μmol catechin/100 mL solution, which was by 44.6 times lower, compared to *Echinacea* tincture.

According to literature, there is a proven relationship between antioxidant activity and total phenolic content in herbs, vegetables, and fruits (1, 17). The results of our study showed that *Ginkgo* tincture had the greatest amount of phenol compounds, and *Echinacea* tincture had the strongest properties of

free radical scavenging and nitration inhibition.

The study showed that tinctures produced from all three investigated plants – *Echinacea purpurea* plant, *Ginkgo biloba* leaves, and *Panax ginseng* roots – had antioxidative activity. The obtained findings allow estimation of the ability of these tinctures to inactivate free radicals. Although the greatest amount of phenol compounds was found in *Ginkgo biloba* tincture, *Echinacea* tincture had the greatest antioxidative activity. This allows stating that antioxidative activity is not directly related to the amount of phenol compounds detected in plants.

Conclusions

1. We found that the amount of phenol compounds in *Ginkgo* tincture was statistically significantly greater than that in *Echinacea* or *Ginseng* tinctures.

2. Tests on DPPH radical cation scavenging and inhibition of nitration showed that the antioxidative activity of *Echinacea* tincture was statistically significantly greater compared to that of *Ginkgo* or *Ginseng* tinctures. This allows us to conclude that antioxidative activity is determined not only by phenol compounds, but also by a complex of other components of medicinal raw material.

Ginkmedžių, ežiulių ir ženšenų tinktūrų antioksidacinis aktyvumas

Rūta Masteikova, Jan Muselik, Jurga Bernatienė¹, Rūta Bernatienė²

Veterinarijos ir farmacijos mokslų universitetas, Farmacijos katedra, Čekijos Respublika,

Kauno medicinos universiteto, ¹Vaistų technologijos ir socialinės farmacijos katedra,

²Farmacinės chemijos ir farmakognozijos katedra, Lietuva

Raktažodžiai: antioksidacinis aktyvumas, ginkmedis, ežiulė, ženšenis.

Santrauka. Darbo tikslas. Nustatyti fenolinių junginių kiekį tinktūrose, pagamintose iš ginkmedžių lapų, rausvažiedžių ežiulių žolės bei ženšenų šaknų, ir įvertinti šių preparatų antioksidacinį aktyvumą. Antioksidacinį aktyvumą tyrėme naudodami modelinius 2,2-difenil-1-pikrilhidrazilo (DPPH•) radikalų katijono sujungimo ir tirozino nitrinimo inhibicijos testus. Gauti rezultatai parodė, kad fenolinių junginių kiekis tirtose tinktūrose skiriasi. Jis svyravo nuo 11±4 iki 340±29 GAE (galo rūgšties ekvivalentai) mg/100 ml. Nustatėme, kad ginkmedžių tinktūroje fenolinių junginių kiekis yra statistiškai reikšmingai didesnis nei ežiulių ar ženšenų tinktūrose. Ginkmedžių tinktūros antioksidacinis efektyvumas buvo 52,7 proc. ($p<0,01$) mažesnis (nuo 1343±11 μmol katechino/100 ml tirpalo iki 637±64 μmol katechino/100 ml tirpalo) palyginti su ežiulių tinktūra. Silpniausia radikalų gaudyklė buvo ženšenų tinktūra – tik 8±1 μmol katechino/100 ml tirpalo. Tirozino nitrinimo inhibicija buvo 34 proc. ($p<0,01$) didesnė ežiulių ekstrakto palyginti su ginkmedžių (nuo 892±36 μmol katechino/100 ml tirpalo iki 588±17 μmol katechino/100 ml tirpalo). Mažiausia tirozino nitrinimo inhibicija pasižymėjo ženšenų tinktūra – tik 20±8 μmol katechino/100 ml tirpalo ir tai buvo 44,6 kartus mažiau palyginti su ežiulių tinktūra.

DPPH• radikalų katijono sujungimo ir nitrinimo inhibicijos testai parodė, kad ežiulių tinktūros antioksidacinis aktyvumas statistiškai reikšmingai didesnis palyginti su ginkmedžių ar ženšenų tinktūromis. Taigi galima daryti išvadą, kad antioksidacinį aktyvumą lemia ne tik fenoliniai junginiai, bet ir kompleksas kitų medžiagų, įeinančių į vaistinės žaliavos sudėtį.

Adresas susirašinti: J. Bernatienė, KMU Vaistų technologijos ir socialinės farmacijos katedra, A. Mickevičiaus 9, 44307 Kaunas. El. paštas: jurgabernatieniene@yahoo.com

References

1. Cai YZ. Structure-radical scavenging activity relationships of phenolic compounds from traditional Chinese medicinal plants. *Life Sci* 2005;30:15-20.
2. Kasparavičienė G, Briedis V. Juodųjų serbentų ir juodavaisių aronijų uogų sulčių stabilumas ir antioksidacinis aktyvumas. (Stability and antioxidant activity of black currant and black aronia berry juices.) *Medicina (Kaunas)* 2003;39:Suppl 2 65-9.
3. Jayaprakasha GK. Phenolic constituents in the fruits of *Cinnamomum zeylanicum* and their antioxidant activity. *J Agric Food Chem* 2006;54:1672-6.
4. Dalby-Brown L. Synergistic antioxidative effects of alkaloids, caffeic acid derivatives, and polysaccharide fractions from *Echinacea purpurea* on *in vitro* oxidation of human low-density lipoproteins. *J Agric Food Chem* 2005;53:9413-23.
5. Mielke MM. Hematologic risk factors of vascular disease and their relation to dementia. *Dement Geriatr Cogn Disord* 2006;21:335-8.
6. Lee TK. Radioprotective potential of ginseng. *Mutagenesis* 2005;20:237-9.
7. Jurkštienė V, Kondrotas AJ, Kėvelaitis E. Imuninės sistemos kompensacinės reakcijos ir rausvažiedės ežiulės (*Echinacea purpurea* (L.) Moench) preparatų poveikis. (Compensatory reactions of immune system and action of purple coneflower (*Echinacea purpurea* (L.) Moench) preparations). *Medicina (Kaunas)* 2004;40:657-62.
8. Chow G. Dietary *Echinacea purpurea* during murine pregnancy: effect on maternal hemopoiesis and fetal growth. *Biol Neonate* 2006;89:133-6.
9. Baranauskas A, Bernatienė J, Radžiūnas R, Bernatoni D. Imuninę sistemą gerinančio ekstrakto technologijos ypatybės. (Technology of extract improving the immune system.) *Medicina (Kaunas)* 2005;41:693-7.
10. Pettinari A. Effect of polyphenolic compounds on the proteolytic activities of constitutive and immuno-proteasomes. *Antioxid Redox Signal* 2006;8:121-6.
11. Hu C. Studies on the antioxidant activity of *Echinacea* root tincture. *J Agric Food Chem* 2000;48:1466-8.
12. Marcondes S. Cyclic GMP-independent mechanisms contribute to the inhibition of platelet adhesion by nitric oxide donor: a role for {alpha}-actinin nitration. *Proc Natl Acad Sci* 2006;103:3434-9.
13. Bernatienė J, Savickas A, Malinauskas F, Bernatienė R. Ginkmedžių tinktūros technologija ir analizė. (Technology and analysis of *Ginkgo* tincture). *Medicina (Kaunas)* 2002;38:1220-3.
14. Yokota J. Scavenging of reactive oxygen species by *Eriobotrya japonica* seed tincture. *Biol Pharm Bull* 2006;29:467-75.
15. Jarašiūnienė D, Šimaitis A. Oksidacinis stresas ir endotelio disfunkcija. (Oxidative stress and endothelial dysfunction). *Medicina (Kaunas)* 2003;39:1151-7.
16. Kasina S, Rizwani W, Radhika KV, Singh SS. Nitration of profilin effects its interaction with poly (L-proline) and actin. *J Biochem (Tokyo)* 2005;138:687-9.
17. Matsunaga H. Relationship between antiproliferative activity of acetylenic alcohol, panaxydol, and its affinity for target cell membrane. *Gan To Kagaku Ryoho* 1994;21:2585-9.

Received 28 June 2006, accepted 29 August 2006

Straipsnis gautas 2006 06 28, priimtas 2006 08 29