Two Furanosesterterpenes from the sponge *Luffariella variabilis*

Peni Ahmadi¹, Masahiro Higashi¹, Nicole J. de Voogd², Junichi Tanaka^{1,*}

¹ Department of Chemistry, Biology and Marine Science, University of the Ryukyus, Nishihara, Okinawa, 903-0213, Japan.

² Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, the Netherlands.

.

Contents

Figure S1. ¹ H-NMR spectrum of compound 1	. 2
Figure S2. ¹³ C-NMR spectrum of compound 1	. 2
Figure S3. HSQC spectrum of compound 1	. 3
Figure S4. HMBC spectrum of compound 1	. 3
Figure S5. COSY spectrum of compound 1	.4
Figure S6. ¹ H-NMR spectrum of compound 3	.4
Figure S7. ¹ H-NMR spectrum of compound 2	. 5
Figure S8. ¹³ C-NMR spectrum of compound 2	. 5
Figure S9. HSQC spectrum of compound 2	. 6
Figure S10. HMBC spectrum of compound 2	. 6
Figure S11. COSY spectrum of compound 2	. 7
Figure S12. HSQC-TOCSY spectrum of compound 2	. 7
Figure S13. ECD spectrum of compound 2	. 8
Figure S14. NOESY spectrum of compound 2	. 8



Figure S1. ¹H-NMR spectrum of compound **1** (CDCl₃, 500 MHz).











Figure S6. ¹H-NMR spectrum of compound **3** (CDCl₃, 500 MHz).



Figure S7. ¹H-NMR spectrum of compound **2** (CDCl₃, 500 MHz).











Figure S12. HSQC-TOCSY spectrum of compound 2 (CDCl₃, 500 MHz).

