Supplementary Material:

New Naphthoquinone Terpenoids from Marine Actinomycete, Streptomyces sp. CNQ-509

Jin-Soo Park and Hak Cheol Kwon *

Natural Constituents Research Center, Korea Institute of Science and Technology (KIST), Gangneung, Gangwon-do 25451, Republic of Korea; jinsoopark@kist.re.kr(J.-S.P.); hkwon@kist.re.kr(H.C.K.);

* Correspondence: hkwon@kist.re.kr; Tel.: +82-33-650-3504

Table of Contents

- Figure S1. ¹H spectrum of **1** in CDCl₃
- Figure S2. ¹³C NMR spectrum of **1** in CDCl₃
- Figure S3. COSY spectrum of 1 in CDCl₃
- Figure S4. HSQC NMR spectrum of 1 in CDCl₃
- Figure S5. HSQC NMR spectrum of 1 in CDCl₃
- Figure S6. NOESY NMR spectrum of 1 in CDCl₃
- Figure S7. ¹H spectrum of **2** in CDCl₃
- Figure S8. ¹³C NMR spectrum of 2 in CDCl₃
- Figure S9. ¹H spectrum of **1a** in CDCl₃
- Figure S10. ¹H spectrum of *epi-1a* in CDCl₃
- Figure S11. ¹H spectrum of *epi-1b* in pyridine-*d*⁵
- Figure S12. ¹H spectrum of *epi-*1c in pyridine-*d*⁵
- Figure S13. On-line-ABTS⁺ assay of Trolox (a), **1** (b) and **2** (c).



Figure S3. COSY spectrum of 1 in CDCl₃



Figure S4. HSQC NMR spectrum of 1 in CDCl₃



Figure S5. HSQC NMR spectrum of 1 in CDCl₃



Figure S6. NOESY NMR spectrum of $\mathbf{1}$ in CDCl₃











Figure S11. ¹H spectrum of *epi-1b* in pyridine- d_5



Figure S13. On-line-ABTS⁺ assay of Trolox (a), **1** (b) and **2** (c). Ten milliliters were analyzed by HPLC with a DAD at 210 nm and 268 nm prior to reaction with the ABTS⁺ radical and the analysis of antioxidant potential at 734 nm.





