

Supplementary Materials

Rhabdastrellosides A and B, isomalabaricane glycosides from the marine sponge *Rhabdastrella globostellata*, and their cytotoxic and cytoprotective effects

Anastasia B. Kozhushnaya^{1,1}, Sophia A. Kolesnikova^{1,1,*}, Ekaterina A. Yurchenko¹, Ekaterina G. Lyakhova¹, Alexander S. Menshov¹, Anatoly I. Kalinovsky¹, Roman S. Popov¹, Pavel S. Dmitrenok¹ and Natalia V. Ivanchina¹



¹ G.B. Elyakov Pacific Institute of Bioorganic Chemistry, Far Eastern Branch of Russian Academy of Sciences, Pr. 100-let Vladivostoku 159, 690022 Vladivostok, Russia; kozhushnaia.ab@mail.ru (A.B.K.); eyurch@piboc.dvo.ru (E.A.Y.); elyakhova@inbox.ru (E.G.L.); menshov90@piboc.dvo.ru (A.S.M.); kaaniw@piboc.dvo.ru (A.I.K.); prs_90@mail.ru (R.S.P.); paveldmt@piboc.dvo.ru (P.S.D.); ivanchina@piboc.dvo.ru (N.V.I.) ¹ A.B.K. and S.A.K. contributed equally.

* Correspondence: sovin81@inbox.ru (S.A.K.); Tel.: +7-423-231-1168


Contents:

- S1** (–)HRESIMS of Rhabdastrelloside A (**1**) and MS/MS spectrum of its [M – H][–] precursor ion at *m/z* 838
- S2** HRESIMS (Positive Ion Mode) of Rhabdastrelloside A (**1**)
- S3** ¹H NMR Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)
- S4** ¹³C NMR Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)
- S5** HSQC Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)
- S6** HMBC Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)
- S7** COSY Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)
- S8** Expanded COSY Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)
- S9** 2D TOCSY Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)
- S10** Expanded 2D TOCSY Spectrum for disaccharide moiety in Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)
- S11** ROESY Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)
- S12** UV and ECD Spectra of Rhabdastrelloside A (**1**) in EtOH
- S13** (–)HRESIMS of Rhabdastrelloside B (**2**) and MS/MS spectrum of its [M – H][–] precursor ion at *m/z* 822
- S14** HRESIMS (Positive Ion Mode) of Rhabdastrelloside B (**2**)
- S15** ¹H NMR Spectrum of Rhabdastrelloside B (**2**) in CD₃OD (500 MHz)
- S16** ¹³C NMR Spectrum of Rhabdastrelloside B (**2**) in CD₃OD (500 MHz)
- S17** HSQC Spectrum of Rhabdastrelloside B (**2**) in CD₃OD (500 MHz)
- S18** HMBC Spectrum of Rhabdastrelloside B (**2**) in CD₃OD (700 MHz)
- S19** COSY Spectrum of Rhabdastrelloside B (**2**) in CD₃OD (500 MHz)
- S20** ROESY Spectrum of Rhabdastrelloside B (**2**) in CD₃OD (500 MHz)
- S21** 1D TOCSY Spectra of Rhabdastrelloside B (**2**) in CD₃OD (500 MHz)
- S22** UV and ECD Spectra of Rhabdastrelloside B (**2**) in EtOH
- S23** Determination of Absolute Configurations of Monosaccharides: GC-MS profiles.
- S24** Determination of Absolute Configurations of Monosaccharides: GC-MS profiles for sum of Standard L-Glc and D-Glc, and Standard L-2-NHAcGlc.
- S25** Determination of Absolute Configurations of Monosaccharides: EIMS spectra of detected D-Glc and Standard D-Glc.
- S26** Determination of Absolute Configurations of Monosaccharides: EIMS spectra of detected D-Glc and Standard D-Glc (Continuation).
- S27** Determination of Absolute Configurations of Monosaccharides: EIMS spectra of detected D-2-NHAcGlc and Standard D-2-NHAcGlc.
- S28** The Isolation Scheme


S1 (-)HRESIMS of Rhabdastrelloside A (**1**) and MS/MS spectrum of its $[M - H]^-$ precursor ion at m/z 838




S2 HRESIMS (Positive Ion Mode) of Rhabdastrelloside A (1**)**




S3 ^1H NMR Spectrum of Rhabdastrelloside A (**1**) in CD_3OD (700 MHz)




S4 ^{13}C NMR Spectrum of Rhabdastrelloside A (**1**) in CD_3OD (700 MHz)




S5 HSQC Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)




S6 HMBC Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)




S7 COSY Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)




S8 Expanded COSY Spectrum of Rhabdastrelloside A (**1**) in CD_3OD (700 MHz)




S9 2D TOCSY Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)





S10 Expanded 2D TOCSY Spectrum for disaccharide moiety in Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)





S11 ROESY Spectrum of Rhabdastrelloside A (**1**) in CD₃OD (700 MHz)




S12 UV and ECD Spectra of Rhabdastrelloside A (**1**) in EtOH




S13 (-)HRESIMS of Rhabdastrelloside B (**2**) and MS/MS spectrum of its $[M - H]^-$ precursor ion at m/z 822




S14 HRESIMS (Positive Ion Mode) of Rhabdastrelloside B (**2**)




S15 ^1H NMR Spectrum of Rhabdastrelloside B (**2**) in CD_3OD (500 MHz)




S16 ^{13}C NMR Spectrum of Rhabdastrelloside B (**2**) in CD_3OD (500 MHz)




S17 HSQC Spectrum of Rhabdastrelloside B (**2**) in CD₃OD (500 MHz)




S18 HMBC Spectrum of Rhabdastrelloside B (**2**) in CD₃OD (700 MHz)




S19 COSY Spectrum of Rhabdastrelloside B (**2**) in CD₃OD (500 MHz)





S20 ROESY Spectrum of Rhabdastrelloside B (**2**) in CD₃OD (500 MHz)



S21 1D TOCSY Spectra of Rhabdastrellatoside B (**2**) in CD₃OD (500 MHz)




S22 UV and ECD Spectra of Rhabdastrelloside B (**2**) in EtOH




S23 Determination of Absolute Configurations of Monosaccharides: GC-MS profiles.

Sample Name: G2.6.3.13
 Misc Info : 90 mkl 1 mkl
 Vial Number: 1




Sample Name: DGlcNAc R(-)
 Misc Info : 300 mkl 0.6 mkl
 Vial Number: 1




Acid hydrolysis of compound **2** (1.4 mg) was carried out as described in the text of the article (See part 4.5) The acetylated 2-octylglycosides were analyzed by GC-MS using the corresponding authentic samples prepared by the same procedure.

GC-MS analyses were carried out on a Hewlett Packard HP6890 GC System, with a HP-5MS capillary column (30 m x 0.25 mm, 0.25 μ m). The carrier gas was helium (flow rate 1.0 mL/min), the injector temperature was 270 °C, the ionizing voltage was 70 eV and the temperature program was 100 °C (0.5 min) –5 °C/min –250 °C (10 min).

Sample Name: DGlc(-)
 Misc Info : 0.2 mg 300 mkl 0.4 mkl
 Vial Number: 1




S24 Determination of Absolute Configurations of Monosaccharides: GC-MS profiles for sum of Standard L-Glc and D-Glc, and Standard L-2-NHAcGlc.




According to the retention times (t_R) and corresponding mass-spectra, L-monosaccharides were not found among acetylated 2-octylglycosides of hydrolysate products.

S25 Determination of Absolute Configurations of Monosaccharides: EIMS spectra of detected D-Glc and Standard D-Glc.


Sample Name: G2.8.3.13
Misc Info : 50 μ l 1 μ l
Vial Number: 1




Sample Name: DGlc(-)
Misc Info : 0,2 mg 300 μ l 0,4 μ l
Vial Number: 1





Sample Name: G2.8.3.13
Misc Info : 50 μ l 1 μ l
Vial Number: 1



Sample Name: DGlc(-)
Misc Info : 0,2 mg 300 μ l 0,4 μ l
Vial Number: 1




S26 Determination of Absolute Configurations of Monosaccharides: EIMS spectra of detected D-Glc and Standard D-Glc (Continuation).




S27 Determination of Absolute Configurations of Monosaccharides: EIMS spectra of detected D-2-NHAcGlc and Standard D-2-NHAcGlc.


Sample Name: G2.8.3.13
Misc Info : 50 mkl 1 mkl
Vial Number: 1



Sample Name: DGlcNAc R(-)
Misc Info : 300 mkl 0.6 mkl
Vial Number: 1



S28 The Isolation Scheme



Cyclobutastelletolides A and B together with known globostelletins M, K, E, F were reported by us in 2019 (*J. Nat. Prod.*, <https://doi.org/10.1021/acs.jnatprod.9b00824>), and series of stellettins Q-V, along with known globostellettin N, was described in 2021 (*Molecules*, <https://doi.org/10.3390/molecules26030678>).