



Article

# The Association between Ambient Air Pollution and Allergic Rhinitis: Further Epidemiological Evidence from Changchun, Northeastern China

Bo Teng <sup>1,\*</sup>, Xuelei Zhang <sup>2,3,\*</sup>, Chunhui Yi <sup>4</sup>, Yan Zhang <sup>5</sup>, Shufeng Ye <sup>1</sup>, Yafang Wang <sup>1</sup>, Daniel Q. Tong <sup>3,6</sup> and Binfeng Lu <sup>1,7</sup>

<sup>1</sup> Department of Otolaryngology Head and Neck Surgery, The Second Hospital, Jilin University, Changchun, 130041, China; yeshufeng16@163.com (S.Y.); wangyafangwang@163.com (Y.W.); binfeng@pitt.edu (B.L.)

<sup>2</sup> Key Laboratory of Wetland Ecology and Environment, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun 130102, China

<sup>3</sup> Center for Spatial Information Science and Systems, George Mason University, Fairfax, VA 22030, USA; tongquansong@neigae.ac.cn

<sup>4</sup> Department of Pathology, Mount Sinai West, New York City, NY 10019, USA; yich1@hotmail.com

<sup>5</sup> Department of Otolaryngology Head and Neck Surgery, The First Hospital, Jilin University, Changchun 130021, China; zhangyan993300@163.com

<sup>6</sup> U.S. NOAA Air Resources Laboratory, College Park, MD 20740, USA

<sup>7</sup> Department of Immunology, University of Pittsburgh School of Medicine, Pittsburgh, PA 15261, USA

\* Correspondence: tengbo1975@163.com (B.T.); zhangxuelei@neigae.ac.cn (X.Z.)

Academic Editor: Paul B. Tchounwou

Received: 16 December 2016; Accepted: 13 February 2017; Published: 23 February 2017

**Abstract:** With the continuous rapid urbanization process over the last three decades, outdoors air pollution has become a progressively more serious public health hazard in China. To investigate the possible associations, lag effects and seasonal differences of urban air quality on respiratory health (allergic rhinitis) in Changchun, a city in Northeastern China, we carried out a time-series analysis of the incidents of allergic rhinitis (AR) from 2013 to 2015. Environmental monitoring showed that PM<sub>2.5</sub> and PM<sub>10</sub> were the major air pollutants in Changchun, followed by SO<sub>2</sub>, NO<sub>2</sub> and O<sub>3</sub>. The results also demonstrated that the daily concentrations of air pollutants had obvious seasonal differences. PM<sub>10</sub> had higher daily mean concentrations in spring (May, dust storms), autumn (October, straw burning) and winter (November to April, coal burning). The mean daily number of outpatient AR visits in the warm season was higher than in the cold season. The prevalence of allergic rhinitis was significantly associated with PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub> and NO<sub>2</sub>, and the increased mobility was 10.2% (95% CI, 5.5%–15.1%), 4.9% (95% CI, 0.8%–9.2%), 8.5% (95% CI, –1.8%–19.8%) and 11.1% (95% CI, 5.8%–16.5%) for exposure to each 1-Standard Deviation (1-SD) increase of pollutant, respectively. Weakly or no significant associations were observed for CO and O<sub>3</sub>. As for lag effects, the highest Relative Risks (RRs) of AR from SO<sub>2</sub>, NO<sub>2</sub>, PM<sub>10</sub> and PM<sub>2.5</sub> were on the same day, and the highest RR from CO was on day 4 (L4). The results also indicated that the concentration of air pollutants might contribute to the development of AR. To summarize, this study provides further evidence of the significant association between ambient particulate pollutants (PM<sub>2.5</sub> and PM<sub>10</sub>, which are usually present in high concentrations) and the prevalence of respiratory effects (allergic rhinitis) in the city of Changchun, located in Northeastern China. Environmental control and public health strategies should be enforced to address this increasingly challenging problem.

**Keywords:** air pollution; allergic rhinitis; significant association; seasonal effect; lag effects; Changchun

## 1. Introduction

Along with the rapid economic growth and urbanization, the severe and deteriorating regional haze has smothered the eastern region of China. The health effects caused by outdoor air pollution have become a sensitive topic for the public, media and even the government of China and adjacent countries. The need for a better comprehension to the role of ambient air pollution on human health and implementing suitable protective policies has fueled related studies in the past decade.

A number of adverse health effects, including non-accidental death, respiratory diseases (such as rhinitis, asthma, tracheitis, pneumonia), cardiovascular diseases (such as stroke, arrhythmia, ischemic heart disease, cerebrovascular disease), cardiopulmonary diseases (chronic obstructive pulmonary disease, COPD) and, more rarely, conjunctivitis, dermatological disorders, skin allergy and exacerbated cough are associated with ambient air pollution [1].

Most studies regarding respiratory diseases have addressed asthma and COPD, and only limited studies have focused on allergic rhinitis (AR) in China. As a typical respiratory illness, AR affects 20%–40% of the population worldwide, although the prevalence varies with age and region [2,3]. Although it is usually a minor respiratory disease, AR frequently presented with symptoms that affect work performance and quality of daily life, and consumes health recourses [4]. According to the Allergies in Asia-Pacific Survey, one of the largest studies of AR on adults and children in Asia, the prevalence of AR was 8.7% in Asia. The prevalence of self-reported AR in adults is much lower in China than in many Western and developed/developing countries (such as Japan and Korea). The age- and gender-adjusted incidence of AR was approximately 14% in China, ranging from 8.7% (Beijing) to 24.1% (Urumqi) in Figure 1. The prevalence of AR for adults was 11.2% and 15.7% in Changchun and Shenyang of northeastern China, respectively [5,6]. According to a cross-sectional questionnaire survey during 2010–2012, the prevalence of rhinitis in the 10 cities varied from 2.2% to 23.9% (mean 8.5%) [7].


Globally, studies have shown associations between vehicle and industrial emissions and increased risk of AR [8,9]. Individual pollutants responsible for the increased risk of allergic disease were nitrogen oxides ( $\text{NO}_x$ ), sulphur dioxide ( $\text{SO}_2$ ), ozone ( $\text{O}_3$ ), particulates with an aerodynamic diameter of 10  $\mu\text{m}$  or less ( $\text{PM}_{10}$ ), and particulates with an aerodynamic diameter of 2.5  $\mu\text{m}$  or less ( $\text{PM}_{2.5}$ ) [10,11]. Contradictory findings also have been found for the  $\text{SO}_2$ ,  $\text{NO}_2$ ,  $\text{O}_3$  and  $\text{PM}_{10}$  levels in some other studies for children and the elderly [7,12]. Both cross-sectional and cohort studies have shown obvious associations between traffic  $\text{NO}_2$  pollution and AR in children [8,13]. Ozone was also associated with AR in children who reside in industrial areas [9]. Air pollutants, such as  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$ ,  $\text{O}_3$ ,  $\text{CO}$ ,  $\text{NO}$ ,  $\text{NO}_2$ , and  $\text{SO}_2$  had positive correlation with AR incidents, when compared with meteorological factors in a heavy industry area of northern Taiwan [14]. There was no association between mean level of pollutants ( $\text{SO}_2$ ,  $\text{NO}_2$ ,  $\text{O}_3$ ) and symptoms of Ear, Nose, and Throat (acute rhinitis, 12-month rhinitis, fever, rhinoconjunctivitis, and hay fever) in children. Furthermore, several studies have also demonstrated that air pollution can promote and exaggerate response to allergens in the nasal cavity by increasing the allergenicity and bioavailability of airborne pollen allergens [15].

However, for the ambient pollutants, only six studies reports on the influence of outdoor air pollutant and the prevalence of AR in Asia [16]. A nationwide cross-sectional study covering Taiwanese schoolchildren showed that the prevalence of AR was associated with levels of  $\text{SO}_2$ ,  $\text{CO}$ , and  $\text{NO}_x$ , but not with levels of  $\text{O}_3$  and  $\text{PM}_{10}$  [17]. Another Taiwanese study reported that children's AR was associated with non-summer warmth and traffic-related air pollutant levels, including  $\text{CO}$ ,  $\text{NO}_x$  and  $\text{O}_3$  [18]. For Asia adults, a cross-sectional population-based study in Singapore found that outdoor air pollution was a significant environmental risk factor of AR [19]. A time-series study identified an association between ambient air pollutant levels and daily outpatient visits for AR among 1506 patients (96% adults) in Beijing [20].  $\text{NO}_2$  and  $\text{SO}_2$  concentration, but not  $\text{PM}_{10}$ , were associated with increased prevalence of AR among kindergarten children in seven cities in Liaoning Province during 2007–2008 [21]. In addition, a study in Changsha (China) showed that the prevalence of AR in

children was significantly positively correlated with age-related accumulative personal exposure of PM<sub>10</sub>, SO<sub>2</sub>, and NO<sub>2</sub> [22].

Moreover, the short-term effects of air pollutants on human health showed seasonal variations with the change of human activity and meteorological factors [23]. Several studies analyzing seasonal effects of air pollutants were focused on major mortality/morbidity [23,24]. There was only one study conducted on the seasonal effect of air pollutants to the AR patients [25]. Therefore, time series data on air pollution and daily number of outpatient for AR is needed to fill in the blanks.

In this article, we discussed the lag effects of air pollutants (specifically focused on particulate matter) on AR in Northeastern China. We also looked into the seasonal effects of air pollutants on the daily number of outpatients with AR.



**Figure 1.** Comprehensive published prevalences of AR in adults and children in different Chinese cities.

## 2. Materials and Methods

### 2.1. Air Pollution and Meteorological Data

Air quality data for the daily PM<sub>2.5</sub>, PM<sub>10</sub>, O<sub>3</sub>, CO, SO<sub>2</sub>, and NO<sub>2</sub> concentrations between 1 January 2013 and 31 December 2015 were provided by the Changchun Municipal Environmental Protection Monitoring Center. The daily data was obtained as average values derived from the hourly data of 10 state-controlled monitoring stations distributed across Changchun, except for O<sub>3</sub> with a running 8-h mean concentrations (which are averaged with specific hour and the preceding 7 h and the averaging period is stepped forward by one hour for each value). Meteorological factors, such

as hourly and daily temperatures, humidity, were obtained from the Weather Underground website ([www.wunderground.com](http://www.wunderground.com)).

## 2.2. Daily Number of AR Outpatients

Daily numbers of outpatients for AR symptoms between 2013 and 2015 were obtained from the Departments of Otolaryngology-Head and Neck Surgery, First and Second Hospitals, Jilin University. Both hospitals are Class-Three, Grade A-level tertiary university hospitals located in the central districts in Changchun. They are both comprehensive teaching and researching medical centers.

The study population includes all outpatients examined by general practitioners and specialties during the study period. AR was defined as symptoms of sneezing or a running, itchy or blocked nose without a cold or flu. The principal diagnosis of allergic rhinitis (ICD-9 code 477) was based on medical history, a physical examination, a standardized questionnaire, and the relevant test (such as, skin prick test).

In order to avoid repeated counting, only one visit per individual patient per day was used as daily visit counts. Subsequent follow-ups within 30 days of the initial visit were excluded. This project was approved by the Ethics Review Board of the two hospitals (2016106). All patients have given written consent to participate in the study. All medical interviewers (general practitioners or nurses) were trained to use uniform examination protocols. Medical records and the respective results were confirmed by the supervisors at each hospital.

## 2.3. Data Analysis

To investigate relationships between AR and ambient air pollution levels, the generalized additive model (GAM) with penalized splines were used to analyze the association of AR with air pollution, adjusting for potential confounders including meteorological factors, time trends, and day of the week. Due to the counted daily outpatients number for AR was small and approximately followed a Poisson distribution [26,27], the core analysis used a GAM with log link and Poisson error that accounted for smooth fluctuations in daily AR patients number.

Two basic steps needed to be conducted before conducting the model analyses, i.e., development of the best base model without any pollutants and the main model with pollutants. The latter is built by adding the air pollution variables to the final cause-specific best base model, assuming there is a linear relationship between the air pollutant concentration and logarithmic outpatient number.

We initially constructed the basic pattern of outpatients excluding the air pollutants by incorporating smoothed spline functions of time and weather conditions. This makes a flexible modeling tool to include non-monotonic and non-linear links between outpatient visits and time/weather conditions. Next, we considered adding the pollutant variables and further analyzed their effects on AR. To compare the relative quality of the outpatient predictions across these non-nested models, Akaike's Information Criterion (AIC) was used as a measure of how well the model fitted the data. Smaller AIC values indicate the preferred model. The following formula (log-linear GAM) is fitted to estimate the pollution log-relative rate  $\beta$ :

$$\log [E(Y_t)] = \alpha + \sum_{i=1}^q \beta_i(X_i) + \sum_{j=1}^p f_j(Z_j, df) + W_t(\text{week})$$

where  $E(Y_t)$  represents the expected outpatient visit number for AR at day  $t$ ;  $\beta$  represents the log-relative risk of outpatient visit associated with an unit increase of air pollutants;  $X_i$  indicates the concentrations of pollutants at day  $t$ ;  $\sum_{j=1}^p f_j(Z_j, df)$  is the non-parametric spline function of calendar time, humidity, temperature, wind speed and barometric pressure;  $W_t(\text{week})$  is the dummy variable for day of the week. More detailed introduction to the GAM has been previously described [27,28].

For the basic models, we also conducted a sensitivity analysis referring to Qian's method [29] and Welty's method [30]. We initialized the  $df$  as 8  $df$  per year for time, 3  $df$  for temperature, humidity, wind speed, and barometric pressure.

We further examined the effect of air pollutants with different lag (L) structures of single-day lag (distributed lag; from L0 to L7) and multi-day lag (moving average lag; L01 to L07). In this study, a lag of 0 days (L0) means the current-day pollution, and a lag of 1 day corresponds to the previous-day pollution. In multi-day lag models, L03 refers to a 4-day moving average of pollutant concentration of the current and previous 3 days [31]. The meteorological variables used in the lag models were the current day's data.

For seasonal analysis, seasonality was differentiated on the basis of heating/non-heating periods. In Changchun, the cold (heating) season is from October to April, and the warm (non-heating) season is from May to September. The majority of the heating in Changchun is provided by a central heating of the city from coal burning power plants. In order to avoid the effects of pollen, we classified the warm season as from mid-May to September, and the cold season as from November to mid-April. Air pollution load during the heating season increases significantly compared to non-heating season.

All statistical analyses were conducted using R version 3.1.2 (mgcv package) (all the related data and code are open-source distributed in the Supplementary Materials). Relative risk (RR) was estimated as  $e^{\beta \times \Delta C}$ , where  $\Delta C$  is the increased amount of air pollutants. In this study, we used standardized deviation (SD) as the  $\Delta C$ . We also calculated percent change in the number of consultations for AR patients by  $(RR-1) \times 100\%$ . A  $p < 0.05$  was considered as statistical significant. All  $p$  values were 2-sided.

### 3. Results

#### 3.1. General Statistical Analysis

Table 1 summarizes the data for the daily number of AR outpatients and meteorological and air pollution variables in Changchun during 2013–2015. There were 23,344 AR outpatients recorded, with a daily mean admission of 21.7 over this 3-year time-series study period. Age distribution and gender of AR outpatients are also summarized in Table S1 of the Supplementary Materials.

**Table 1.** Summary of environmental variables and daily number of outpatients for AR in Changchun, 2013–2015.

Variables	Mean	SD	Max.	Min.	Median	IQR
Number of AR Patients	21.7	24.5	177	0	15	12
PM <sub>2.5</sub> ( $\mu\text{g}/\text{m}^3$ )	66.5	59	495	2	47.3	48.4
PM <sub>10</sub> ( $\mu\text{g}/\text{m}^3$ )	114.4	74.4	642.4	19.5	96.5	70.4
SO <sub>2</sub> ( $\mu\text{g}/\text{m}^3$ )	37	36.9	191.3	2.4	19.1	50.6
CO ( $\text{mg}/\text{m}^3$ )	0.93	0.4	3.3	0.1	0.8	0.46
NO <sub>2</sub> ( $\mu\text{g}/\text{m}^3$ )	43.6	16.1	113.5	11.2	40.9	20.4
O <sub>3</sub> ( $\mu\text{g}/\text{m}^3$ )	71.1	37	332.2	14.2	62.3	44.4
MAXT ( $^{\circ}\text{C}$ )	12	14.5	35	−22	15	26
AVET ( $^{\circ}\text{C}$ )	6.3	14.3	28	−26	9	27
MINT ( $^{\circ}\text{C}$ )	0.7	14.5	24	−30	2	26
Dew ( $^{\circ}\text{C}$ )	−1.5	13.9	22	−31	−2	25
Press (hPa)	1015	9.5	1040	984	1014	14
Wind (km/h)	10.9	5.1	32	2	10	8
AVEH (%)	58.5	15.3	90	13	60	22

MAXT: maximum temperature; AVEH: mean temperature; MINT: minimum temperature; Dew: dew point; Press: sea level press; AVEH: mean humidity.

The results showed the daily mean concentrations of PM<sub>10</sub> and PM<sub>2.5</sub> were 114.4  $\mu\text{g}/\text{m}^3$  to 66.5  $\mu\text{g}/\text{m}^3$  respectively, with both exceeding the yearly concentrations of national level II (70  $\mu\text{g}/\text{m}^3$  and 35  $\mu\text{g}/\text{m}^3$ ). NO<sub>2</sub> and SO<sub>2</sub> had a similar trend of monthly mean concentrations, with the

concentration of NO<sub>2</sub> coming in at 43.6 µg/m<sup>3</sup>, near the national level standard (40 µg/m<sup>3</sup>), whereas the concentration of SO<sub>2</sub> (37.0 µg/m<sup>3</sup>) was also twofold higher than the national level standard (20 µg/m<sup>3</sup>). During the study period, the primary polluting agent in Changchun was the particulate matter emitted from natural and anthropogenic sources, with 27.9% and 20.8% of days PM<sub>2.5</sub> and PM<sub>10</sub> pollution, respectively. NO<sub>2</sub> was second, but with only 3.7% of days above the national standard. During the heating season (November to mid-April) in Figure 1, the concentration of SO<sub>2</sub> was higher. Concentrations of all air pollutants showed obvious seasonal differences.

The mean daily temperature and humidity were 6.3 °C and 58.5%, respectively. The mean daily temperature and humidity ranged from −26 °C to 28 °C, and 13% to 90%, reflecting the northern temperate continental monsoon climate of Changchun. Furthermore, extreme lower temperature and atmospheric boundary layer in winter are significant features in northeastern China.

Pearson correlation coefficients of air pollutants and meteorological variables are shown in Table 2. SO<sub>2</sub>, NO<sub>2</sub>, PM<sub>10</sub>, PM<sub>2.5</sub> and CO had significant positive correlations with each other in both cold and warm seasons ( $p < 0.05$ ), whereas O<sub>3</sub> had a significant negative correlation with the other five air pollutants in the cold season and only negative correlation with NO<sub>2</sub> in the warm season. This correlation was consistent with the stationary fossil fuel combustion-related pollutants (SO<sub>2</sub> and PM) and the traffic-related pollutant NO<sub>2</sub> [32]. Daily wind speed, daily mean temperature and daily dew point were negatively correlated with all air pollutants except for O<sub>3</sub> in the cold season. Daily atmospheric pressure shows a positive relationship with all air pollutants except for O<sub>3</sub> in both cold and warm seasons. Humidity was negatively correlated with all pollutants except for CO in the warm season, and positively correlated with all pollutants except for PM<sub>10</sub> and O<sub>3</sub> in the cold season.

PM<sub>10</sub> and PM<sub>2.5</sub> were highly correlated (correlation coefficient  $r = 0.89$ ) in both the cold and warm season. NO<sub>2</sub> and SO<sub>2</sub> were moderately correlated with PM<sub>2.5</sub> ( $r = 0.85$  and  $r = 0.64$ ) in cold season, but not in the warm season. CO was highly correlated with PM<sub>2.5</sub> ( $r = 0.93$  and  $r = 0.84$ ) in both the cold and warm season due to their acting as byproducts of incomplete combustion, but O<sub>3</sub> was poorly correlated with PM<sub>2.5</sub> ( $-0.23$ ) in cold season and moderately correlated in warm season, which indicated strong photochemical reactions in summer.

### 3.2. Temporal Patterns of Outpatients and Air Pollutants

Figure 2 shows temporal patterns of daily outpatients and daily concentrations of air pollutants in Changchun during 2013–2015. Figure 2 depicts the inter-annual variation of the daily number of AR patients (which ranged from 0 to 177). The mean daily number of AR outpatients was higher in warm than in the cold season. Consistent with the statistical distribution, this number had significant peaks in the warm season, especially around July and September. The number was smaller in other months. This phenomenon could be explained as seasonal allergic rhinitis, which is caused by pollens from weeds and trees [33], such as the pollens emitted from the *Artemisia* and *Ambrosia* in Changchun.


According to the statistical results, there are obvious seasonal differences for SO<sub>2</sub>, NO<sub>2</sub>, O<sub>3</sub> and CO. Although no clear temporal patterns of PM<sub>2.5</sub> and PM<sub>10</sub> concentration are evident in Figure 2, there were seasonal differences under further interpretation. PM<sub>10</sub> had higher daily mean concentrations in spring (May, from dust storms), autumn (October, from straw burning) and winter (November to April, from house heating).

**Table 2.** Pearson’s correlation coefficients among environmental variables in cold season and warm season, Changchun 2013–2015.

	PM <sub>2.5</sub>	PM <sub>10</sub>	SO <sub>2</sub>	CO	NO <sub>2</sub>	O <sub>3</sub>	MAXT	AVET	MINT	DEWP	Press	Wind	AVEH
PM <sub>2.5</sub>	1	0.89 **	0.64 **	0.93 **	0.85 **	−0.23 **	−0.17 *	−0.23	−0.27 **	−0.23 **	0.35 **	−0.14 **	0.09 **
	<i>1</i>	<i>0.89 **</i>	<i>0.17 *</i>	<i>0.84 **</i>	<i>0.21 **</i>	<i>0.48 **</i>	<i>0.34 **</i>	<i>0.36</i>	<i>0.35 **</i>	<i>0.30 *</i>	<i>0.06 **</i>	<i>0.01 **</i>	<i>−0.00 **</i>
PM <sub>10</sub>		1	0.38 **	0.80 **	0.69 **	−0.018	0.073	0.01 **	−0.06	−0.09 **	0.28 **	−0.03 **	−0.15
		<i>1</i>	<i>0.39 **</i>	<i>0.81 **</i>	<i>0.47 **</i>	<i>0.42 **</i>	<i>0.46 **</i>	<i>0.38 **</i>	<i>0.28</i>	<i>0.18 **</i>	<i>0.19 **</i>	<i>−0.10 **</i>	<i>−0.26 **</i>
SO <sub>2</sub>			1	0.68 **	0.68 **	−0.58 **	−0.64 **	−0.67 **	−0.65 **	−0.53	0.41 **	−0.28 **	0.46 **
			<i>1</i>	<i>0.22 **</i>	<i>0.52 **</i>	<i>0.26 **</i>	<i>0.05</i>	<i>−0.08 *</i>	<i>−0.19 **</i>	<i>−0.42</i>	<i>0.10 **</i>	<i>0.08 **</i>	<i>−0.64 **</i>
CO				1	0.90 **	−0.30 **	−0.19 **	−0.24 **	−0.27 **	−0.25 **	0.36	−0.12 **	0.07 **
				<i>1</i>	<i>0.35 **</i>	<i>0.42 **</i>	<i>0.32 **</i>	<i>0.31 **</i>	<i>0.28 **</i>	<i>0.31 **</i>	<i>0.21</i>	<i>−0.17 **</i>	<i>0.10 **</i>
NO <sub>2</sub>					1	−0.36 **	−0.18 **	−0.23 **	−0.27 **	−0.27 **	0.35 **	−0.12	0.01 **
					<i>1</i>	<i>−0.14</i>	<i>0.22 **</i>	<i>0.09 **</i>	<i>−0.05 **</i>	<i>−0.12 **</i>	<i>0.29 **</i>	<i>−0.49</i>	<i>−0.34</i>
O <sub>3</sub>						1	0.40 **	0.38 **	0.34	0.31 **	−0.15 **	0.13 **	−0.22
						<i>1</i>	<i>0.54 **</i>	<i>0.60 **</i>	<i>0.60 **</i>	<i>0.40 **</i>	<i>−0.31 **</i>	<i>0.51</i>	<i>−0.18</i>
MAXT							1	0.97 *	0.89	0.83 **	−0.41 *	0.44 *	−0.54 **
							<i>1</i>	<i>0.93 **</i>	<i>0.77 **</i>	<i>0.65</i>	<i>−0.17 **</i>	<i>−0.02 **</i>	<i>−0.22 **</i>
AVET								1	0.97	0.87 **	−0.48 **	0.50 **	−0.51 **
								<i>1</i>	<i>0.94 **</i>	<i>0.82</i>	<i>−0.35 **</i>	<i>0.10</i>	<i>0.02 **</i>
MINT									1	0.84 **	−0.54 **	0.53 **	−0.46 **
									<i>1</i>	<i>0.90 *</i>	<i>−0.48 **</i>	<i>0.19</i>	<i>0.23 **</i>
DEWP										1	−0.53 **	0.38 **	−0.06 **
										<i>1</i>	<i>−0.37 **</i>	<i>−0.01</i>	<i>0.56 **</i>
Press											1	−0.51 **	0.10
											<i>1</i>	<i>−0.39 **</i>	<i>−0.17 **</i>
Wind												1	−0.39
												<i>1</i>	<i>−0.17 **</i>
AVEH													1
													<i>1</i>

\* Correlation is significant at the 0.05 level (2-tailed); \*\* Correlation is significant at the 0.01 level (2-tailed); Italic values are correlation coefficients for the warm season.


Ranges of PM<sub>2.5</sub>, PM<sub>10</sub>, NO<sub>2</sub>, and O<sub>3</sub>, were wide, with the maximum many times higher than the Class 2 limits of GB3095-2012, according to the national standards of China: GB3095-2012 and HJ633-2012. PM<sub>2.5</sub> was the major air pollutant in Changchun, and there were a total of 306 days that had heavy fine particulate pollution with daily concentrations higher than 75 µg/m<sup>3</sup>. PM<sub>10</sub> was the secondary air pollutant, and there were 228 days and 13 days that had heavy PM<sub>10</sub> and SO<sub>2</sub> pollutions, with daily concentrations exceeding 150 µg/m<sup>3</sup>, respectively. NO<sub>2</sub> ranged from 11.2 to 113.5 µg/m<sup>3</sup>, and the annual mean level was 43.6 µg/m<sup>3</sup>. A total of 40 days had heavy NO<sub>2</sub> pollution with daily concentrations exceeding 80 µg/m<sup>3</sup>. Around 33 days had heavy O<sub>3</sub> pollution with daily maximum 8-h mean concentrations exceeding 160 µg/m<sup>3</sup>, most of which were during June–October. No polluted days for CO were observed during the study period.



**Figure 2.** Temporal variations of daily numbers for AR patients and ambient air pollutants (PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub>, CO, NO<sub>2</sub> and O<sub>3</sub>) in Changchun during 2013–2015.

### 3.3. Exposure-Response Associations


Figure 3 shows the exposure-response relationships for air pollutants with outpatient visits for AR. In this study, we found generally linear relationships (monotonic trends) for AR hospital visits associated with PM<sub>2.5</sub>, PM<sub>10</sub> and CO. Moreover, we observed basically monotonic increased relative risk for both SO<sub>2</sub> and NO<sub>2</sub> within these ranges of concentrations (Figure 3), indicating that SO<sub>2</sub> and NO<sub>2</sub> are significantly associated with increased hospital visits of AR.



**Figure 3.** Smoothed plots of exposure-response relations between air pollutants (current day) and outpatient visits for AR in Changchun, China (2012–2015). The solid line presents log relative risk of outpatient visits for AR, while the dashed lines present 95% confidence interval (CI) of the log relative risk.

Figure 4 shows the exposure-response relationships for meteorological factors with hospital visits for AR. The exposure-response relationships associated with humidity and atmospheric

pressure were essentially linear with monotonic increases, respectively. The relationships of minimal and maximal temperature were non-linear with higher positive dose-response functions at lower temperatures ( $<0\text{ }^{\circ}\text{C}$ ).



**Figure 4.** Smoothed plots of exposure-response relations between meteorological factors and outpatient visits for AR in Changchun, China (2012–2015). The solid line presents log relative risk of outpatient visits for AR, while the dashed lines present 95% confidence interval (CI) of the log relative risk.

We observed a positive exposure–response relationship between lower wind speeds ( $<10\text{ km/h}$ ) and AR relative risk, which is the result of static wind condition that not suitable for the diffusion of atmospheric pollutants.

Table 3 illustrates the associations between air pollution and the prevalence of AR in Changchun. It is obvious that high level of PM<sub>2.5</sub>, PM<sub>10</sub>, NO<sub>2</sub> and SO<sub>2</sub> were associated with increased hospital visits of AR. Among these pollutants, NO<sub>2</sub> and PM<sub>2.5</sub> showed stronger influences on outpatient visits than SO<sub>2</sub> and PM<sub>10</sub>. Each 1-SD increase of PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub> and NO<sub>2</sub> exposure was associated with 10.2% (95% CI, 5.5%–15.1%), 4.9% (95% CI, 0.8%–9.2%), 8.5% (95% CI, –1.8%–19.8%) and 11.1% (95% CI, 5.8%–16.5%) increase of AR prevalence, respectively. No significant associations were observed for CO and O<sub>3</sub>.

**Table 3.** Associations between each 1-SD increase of current day air pollution and allergic rhinitis.

Pollutant	Coefficient $\beta$ (95% CI)	Relative Risk RR (95% CI)	Percent Change, % (95% CI)
PM <sub>2.5</sub>	0.097 (0.053, 0.140)	1.102 (1.055, 1.151)	10.2 (5.5, 15.1)
PM <sub>10</sub>	0.048 (0.008, 0.088)	1.049 (1.008, 1.092)	4.9 (0.8, 9.2)
SO <sub>2</sub>	0.081 (–0.018, 0.181)	1.085 (0.982, 1.198)	8.5 (–1.8, 19.8)
NO <sub>2</sub>	0.105 (0.057, 0.153)	1.111 (1.058, 1.165)	11.1 (5.8, 16.5)
CO	–0.023 (–0.098, 0.052)	0.977 (0.907, 1.053)	–2.3 (–9.3, 5.3)
O <sub>3</sub>	–0.007 (–0.060, 0.047)	0.993 (0.941, 1.048)	–0.7 (–5.9, 4.8)

### 3.4. Lag Effects of Air Pollutants on AR

The lag effects of air pollutants on the daily number of AR outpatient visits are further analyzed to identify the possible delayed health effects of air pollutants. Given the high correlations between air pollutants, multiple air pollutant exposure effects on AR in one model were not considered in this study.

Table 4 shows the change of RRs in the number of outpatients for AR with each 1-SD increase in pollutants for single-day, 1–7 days prior to the outpatient visit (L0–L6), and moving average measures from day 0 and day 1 to day 7 prior to the visit. For each pollutant, the cumulative measure is mean (lags 0–1; L0–1), mean (lags 0–2; L0–2), mean (lags 0–3; L0–3), mean (lags 0–4; L0–4), mean (lags 0–5; L0–5) and mean (lags 0–6; L0–6), respectively.

The largest RRs of PM<sub>2.5</sub> for single-day lags were found for the current day (L0), and then decreased to 1.07–1.08 in the following 7 days. The effect magnitude of PM<sub>10</sub> showed a decreasing trend from L0 to L7. The effect magnitude of SO<sub>2</sub> for both single-day and multi-day lags showed a decreasing trend from the current day to the third day, but then showed an increasing trend from the third day to the sixth day. The greatest RRs were found in the 1-day cumulative measures (L0–1). RRs of NO<sub>2</sub> had similar time trends to that of PM<sub>2.5</sub>, and the largest associations were for L0. Then, the RRs decreased to around 1.05 over the next 5 days. Increasing trends for moving average day effects from L0–1 to L0–6 were observed for PM<sub>2.5</sub>, PM<sub>10</sub>, NO<sub>2</sub>, and CO, whereas there was no association between the current day and lag day of O<sub>3</sub> with AR.

## 4. Discussion

This study utilized environmental and epidemiological data to focus on the air quality with temporal variation and its health effects on daily outpatient visits for AR in Changchun during 2013–2015. In contrast to the self-report study conducted in China [34], our study was based on medical records from the two major hospitals in Changchun. We analyzed the daily number of outpatients for AR and attempted to establish time-series relationships between air pollution, metrological factors, and daily outpatient visits for AR.

**Table 4.** Associations between each 1-SD increase of air pollution with different lag periods and allergic rhinitis prevalence.

Lag	PM <sub>2.5</sub>		PM <sub>10</sub>		SO <sub>2</sub>		NO <sub>2</sub>		CO		O <sub>3</sub>	
	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI
L0	1.102	1.055–1.151	1.049	1.008–1.092	1.085	0.982–1.198	1.111	1.058–1.165	0.977	0.907–1.053	0.993	0.941–1.048
L1	1.071	1.027–1.117	1.029	0.991–1.069	1.036	0.940–1.142	1.048	1.007–1.091	1.008	0.938–1.083	0.975	0.934–1.017
L2	1.085	1.042–1.130	1.048	1.011–1.087	1.004	0.915–1.103	1.043	1.004–1.083	1.026	0.956–1.101	0.983	0.946–1.023
L3	1.081	1.039–1.125	1.049	1.012–1.087	0.984	0.8899–1.078	1.047	1.009–1.086	1.024	0.954–1.099	0.996	0.958–1.035
L4	1.08	1.038–1.123	1.049	1.012–1.088	1.017	0.929–1.113	1.049	1.011–1.087	1.044	0.971–1.121	1.005	0.967–1.044
L5	1.081	1.039–1.126	1.039	1.002–1.078	1.021	0.931–1.120	1.05	1.012–1.089	1.006	0.933–1.085	0.994	0.956–1.033
L6	1.078	1.035–1.121	1.037	1.000–1.076	1.024	0.934–1.124	1.025	0.988–1.064	0.993	0.920–1.073	1.006	0.967–1.045
L0–1	1.113	1.060–1.169	1.053	1.006–1.102	1.086	0.969–1.218	1.109	1.052–1.168	0.992	0.912–1.079	0.971	0.915–1.030
L0–2	1.138	1.080–1.200	1.073	1.021–1.127	1.075	0.950–1.217	1.115	1.055–1.178	1.008	0.920–1.105	0.965	0.908–1.026
L0–3	1.156	1.094–1.222	1.089	1.033–1.149	1.059	0.929–1.207	1.129	1.066–1.197	1.02	0.924–1.125	0.968	0.908–1.032
L0–4	1.171	1.105–1.241	1.104	1.044–1.168	1.065	0.930–1.220	1.143	1.076–1.214	1.037	0.934–1.150	0.974	0.911–1.041
L0–5	1.186	1.117–1.260	1.114	1.049–1.182	1.072	0.931–1.235	1.158	1.088–1.233	1.037	0.929–1.157	0.971	0.906–1.040
L0–6	1.199	1.127–1.277	1.122	1.054–1.194	1.082	0.936–1.252	1.164	1.089–1.243	1.033	0.920–1.160	0.973	0.905–1.046

Compared to former studies conducted in Beijing [20], which only focused on PM<sub>10</sub>, SO<sub>2</sub> and NO<sub>2</sub>, all the officially monitored parameters of air pollution and metrology were included in this study (especially adding PM<sub>2.5</sub>, CO and O<sub>3</sub> as regularly monitored pollutants). Another important difference is that the study period in this paper was extended to 3 years in comparison to former studies, which lasted for only several months to one year [25]. The result shows that daily concentrations of air pollutants had clear seasonal differences. O<sub>3</sub> levels were higher in the warm season than in the cold season. However, concentrations of the other five pollutants were higher in the cold season than in the warm season. Similar pollution characteristics were reported in northern cities of China, such as Beijing [35] and Shenyang [36]. From June to August, the air quality is relatively better due to the dilution effect of the meteorological conditions. Whereas in the heating period in winter, the air quality is comparatively poor because of the prevailing weather conditions and the lowest PBL height. Although the sources of air pollution sources in the mega cities have gradually changed from conventional coal combustion to a mixture of combustion as well as motor vehicle emissions [37], biomass burning sources in northeastern China still have a tremendous contribution on the heavy air pollution [38]. Moreover, during the periods of March to May and August to October, the concentration of air pollen peaks due to a high pollen release [33].

Many studies have reported health effects of PM<sub>2.5</sub> [39], but O<sub>3</sub> also has a strong potentially adverse health effects on various respiratory symptoms such as, dyspnea, upper airway irritation, coughing, and chest tightness [40]. However, no significant association between O<sub>3</sub> and the prevalence of AR was observed in Changchun. Research on harmful effects of O<sub>3</sub> is rare in China, so further studies are needed. De Marco et al. [41] have reported that outdoor NO<sub>2</sub> interacted with climate, and increased the risk for AR in high stable temperatures. Hajat et al. [26] found a strong association between 4-day lag SO<sub>2</sub> and O<sub>3</sub> and the number of consultations for AR in London. However, PM<sub>10</sub> and PM<sub>2.5</sub> are less significant associated with AR in London. Villeneuve et al. [42] reported that there was no statistically significant association between daily levels of air pollution and the number of physician visits for rhinitis among the elderly in Toronto. They only studied SO<sub>2</sub> and NO<sub>2</sub> exposure on the same day of the outpatient visits during the winter period. Similar to other time series studies (Table 5), the present work also revealed a statistically significant association between air pollution and AR. RR percentage changes (RRs) in the number of AR outpatients with a 10 µg/m<sup>3</sup> increase of air pollutants were 2.3%, 6.9%, 1.7% and 0.7% for SO<sub>2</sub>, NO<sub>2</sub>, PM<sub>2.5</sub> and PM<sub>10</sub>, respectively. In contrast with lag effects of air pollutants in other studies, the highest RRs for SO<sub>2</sub>, NO<sub>2</sub>, PM<sub>10</sub>, and PM<sub>2.5</sub> were for the current day and L04 for CO and O<sub>3</sub>. This may be attributed to the differences of exposure risk factors between cities, including environmental factors (geographical location, climatic type and air quality), population (children, adults and elderly) and allergen exposure (time, dose and number). More in-depth studies should be done to clarify the relationship between specific chemical components of particulates (e.g., ions, dust, black carbon, heavy metals, VOCs) and the provenance of AR.

It is of interest and importance to understand the shape of exposure-response association in investigating potential health effects of air pollutants and meteorological factors. In contrast, and like other studies in China [17,20,25], we found nonlinear associations for SO<sub>2</sub> and NO<sub>2</sub> that the relative risk increased at the lower SO<sub>2</sub> and NO<sub>2</sub> concentrations, but attenuated or even turned negative at higher concentrations (Figure 3). One possible reason is that saturation mechanism, when underlying biochemical and cellular processes become saturated with small doses [43]. Another reason is the small sample size at higher concentrations as can be reflected by the wider confidence intervals. Nonetheless, it should be noted that the ranges of air pollutant concentrations in our study were rather wide, compared with that of the U.S. and Europe. This figure also obviously demonstrated that the current air quality standard of national level II for PM<sub>2.5</sub> and PM<sub>10</sub> (daily concentration: 70 µg/m<sup>3</sup> and 150 µg/m<sup>3</sup>) can effectively reduce the effects to health of AR patients in Changchun. Both the cross-sectional studies and time-series studies (Table 5) have firstly established the significant association between AR and traffic-related NO<sub>x</sub> emissions.

**Table 5.** RR in the number of outpatients for AR with a 10 µg/m<sup>3</sup> increase in pollutants for lag effects.

Variables	Toronto [42]		London [26]		Taiwan [17]		Beijing [25]		Changchun	
	Concentration	RR	Concentration	RR	Concentration	RR	Concentration	RR	Concentration	RR
SO <sub>2</sub> (µg/m <sup>3</sup> )	12.3 (7.3)	1.041	21.2 (7.8)	−0.003	9.2 (5.23)	1.43	44.1 (21.0)	−0.022	37.0 (36.9)	1.023
NO <sub>2</sub> (µg/m <sup>3</sup> )	47.8 (14.5)	1.018	63.2 (19.7)	1.017	51.9 (15.7)	1.11	52.7 (22.5)	4.804	43.6 (16.1)	1.069
CO (mg/m <sup>3</sup> )	1.3 (0.5)	1	0.9 (0.5)	1.008	0.7 (0.2)	1.05	-	-	1.5 (0.4)	−0.023
O <sub>3</sub> (µg/m <sup>3</sup> )	59.1 (30.3)	−0.002	34.5 (22.7)	−0.005	45.5 (6.5)	1.05	-	-	71.1 (37.0)	−0.007
PM <sub>2.5</sub> (µg/m <sup>3</sup> )	7.7 (3.3)	−0.001	12.7 (7.9)	1.037	-	-	-	-	66.5 (59.0)	1.017
PM <sub>10</sub> (µg/m <sup>3</sup> )	16.4 (4.9)	−0.004	28.5 (13.7)	1.016	55.6 (16.6)	1	116.1 (67.7)	1.367	114.4 (74.4)	1.007
Temperature	8.4 (10.5)		11.9 (5.0)		23.7 (0.8)		12.0 (12.3)		6.3 (14.3)	
Humidity	70.2 (12.1)		70.4 (10.9)		74.0 (3.0)		53.3 (19.3)		58.5 (15.3)	
Period	1995–2000		1992–1994		2000		2009.03–2010.03		2012–2015	
Population	Elderly (≥64 years)		Adults (15–64 years)		Children (6–15 years)		General (all aged)		General (all aged)	

A controlled exposure experiment was conducted for seasonal AR patients to exposure in air and NO<sub>x</sub> (400 ppb, 6 h), and the results illustrated an increase in the number and/or activation state of eosinophils in the nasal mucosa, which in turn could enhance the upper airway response to the inhaled allergen and also exacerbate the disease [44]. Glück and Gebbers [45] also demonstrated that non-specific toxic damage of the nasal epithelium was possibly caused by air pollutants besides grass pollen. The effects of O<sub>3</sub> on AR were still in debate and were only shown by a few studies [9,11,17]. However, a human exposure experiment suggested that short-term exposure to ozone (250 ppb, 3 h) can increase the bronchial allergen responsiveness in mild allergic asthma or rhinitis patients. Likewise, Peden et al. [46] also reported enhanced nasal inflammatory responses in AR patients after O<sub>3</sub> exposure. According to Hwang and Jaakkola [47], co-exposure to O<sub>3</sub> and NO<sub>2</sub> significantly promoted the release of the cytotoxic protein in the nasal lavage after allergen exposure. No study was concerned with SO<sub>2</sub> exposure. This reveals that more exposure studies need to be conducted with considerations for different concentration gradients of gaseous pollutants to fix the bottom response concentrations for acute exposure.

More and more studies demonstrated the particulates would exacerbate the AR symptoms, and the pathological studies only focus on the diesel exhaust particles (DEP) and transported desert dust particles [48,49]. However, more detailed chemical components (ions, elements, black carbon, organic carbon, etc.) and their effects on AR need to be further investigated, the results of which could be used to propose reasonable control measures for corresponding emission sources. As mentioned above, there are two major mechanisms may account for the increased incidence of AR in industrialized areas. Firstly, increased concentration of pollutants lead to airway sensitization and responsiveness to allergens. Next, airway responsiveness to allergens may subsequently aggravate symptoms of AR [50].

## 5. Conclusions

The current study provided evidence of the adverse effect of ambient air pollution on AR in northeastern China. We examined the lag effects of air pollutants and the possible differences of warm seasons vs. cold seasons of air pollutants on the daily number of outpatients that were consulting the hospital for treatment for AR over a period of 3 years. The results showed that daily concentrations of air pollutants had obvious seasonal differences. The air pollutants positively correlated with the daily number of AR outpatients, and had lag effects on the daily number of AR outpatients. RR in the number of AR outpatients increased with air pollution level. There are also some limitations to our study. With the continuous rapid urbanization process, more people are becoming exposed to high levels of air pollution. Environmental control and public health strategies should be enforced by the health service policy makers to address this increasingly challenging problem. During haze events, both the health care provider and the public should be given real-time alerts on air quality and other allergy indexes. The affected population should be appropriately advised and treated. Autumn is the most important season when alerts need to be raised regarding the cost of health systems for pollen allergies and air pollution from straw burning in northeastern China.

**Supplementary Materials:** The following are available online at [www.mdpi.com/1660-4601/14/3/226/s1](http://www.mdpi.com/1660-4601/14/3/226/s1), Table S1: Age distribution and gender of AR outpatients in Changchun; All the related data and code for statistical analyses are open-source distributed in the Supplementary Materials.

**Acknowledgments:** This work was supported by the National Natural Science Foundation of China (Grant No. 21407148 and 41571063), the Education Department of Jilin Province in twelfth Five years of scientific and technological research project (2015514) and 135 Breeding Project of Chinese Academy of Sciences (Northeast Institute of Geography and Agroecology No. Y6H2081001).

**Author Contributions:** Bo Teng, Daniel Q. Tong, Xuelei Zhang and Binfeng Lu proposed and implemented the study; Xuelei Zhang and Chunhui Yi and Yan Zhang performed the experiments; Shufeng Ye and Yafang Wang performed the data collection and statistical analysis; Bo Teng and Xuelei Zhang wrote the manuscript.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Zhang, Q.; Qiu, M.; Lai, K.; Zhong, N. Cough and environmental air pollution in China. *Pulm. Pharmacol. Ther.* **2015**, *35*, 132–136. [[CrossRef](#)] [[PubMed](#)]
2. Greiner, A.N.; Meltzer, E.O. Overview of the treatment of allergic rhinitis and nonallergic rhinopathy. *Proc. Am. Thorac. Soc.* **2011**, *8*, 121–131. [[CrossRef](#)] [[PubMed](#)]
3. Adriana, I.D.; Valero, A.L.; Mullol, J. Comparative analysis of allergic rhinitis in children and adults. *Curr. Allergy Asthma Rep.* **2013**, *13*, 142–151.
4. Schoenwetter, W.F.; Dupclay, L.; Appajosyula, S.; Botteman, M.F.; Pashos, C.L. Economic impact and quality-of-life burden of allergic rhinitis. *Curr. Med. Res. Opin.* **2004**, *20*, 305–317. [[CrossRef](#)] [[PubMed](#)]
5. Katelaris, C.H.; Lee, B.W.; Potter, P.C.; Maspero, J.F.; Cingi, C.; Lopatin, A.; Saffer, M.; Xu, G.; Walters, R.D. Prevalence and diversity of allergic rhinitis in regions of the world beyond Europe and North America. *Clin. Experiment. Allergy* **2012**, *42*, 186–207. [[CrossRef](#)] [[PubMed](#)]
6. Zhang, Y.; Zhang, L. Prevalence of allergic rhinitis in China. *Allergy Asthma Immunol. Res.* **2014**, *6*, 105–113. [[CrossRef](#)] [[PubMed](#)]
7. Zhang, Y.P.; Li, B.Z.; Huang, C.; Yang, X.; Qian, H.; Deng, Q.; Zhao, Z.H.; Li, A.G.; Zhao, J.N.; Sundell, J.; et al. Ten cities cross-sectional questionnaire survey of children asthma and other allergies in China. *Chin. Sci. Bull.* **2013**, *58*, 4182–4189. [[CrossRef](#)]
8. Lindgren, A.; Strohm, E.; Nihlén, U.; Montnémery, P.; Axmon, A.; Jakobsson, K. Traffic exposure associated with allergic asthma and allergic rhinitis in adults. A cross-sectional study in southern Sweden. *Int. J. Health Geogr.* **2009**, *8*, 25. [[CrossRef](#)] [[PubMed](#)]
9. Kim, B.; Kwon, W.; Seo, H.; Kim, H.B.; Lee, S.Y.; Park, K.S.; Yu, J.; Kim, H.C.; Leem, J.H.; Sakong, J.; et al. Association of ozone exposure with asthma, allergic rhinitis, and allergic sensitization. *Ann. Allergy Asthma Immunol.* **2011**, *107*, 214–219. [[CrossRef](#)] [[PubMed](#)]
10. Pénard-Morand, C.; Charpin, D.; Raherison, C.; Kopferschmitt, C.; Caillaud, D.; Lavaud, F.; Annesi-Maesano, I. Long-term exposure to background air pollution related to respiratory and allergic health in schoolchildren. *Clin. Exp. Allergy* **2005**, *35*, 1279–1287. [[CrossRef](#)] [[PubMed](#)]
11. Kim, H.H.; Lee, C.S.; Jeon, J.M.; Yu, S.D.; Lee, C.W.; Park, J.H.; Shin, D.C.; Lim, Y.W. Analysis of the association between air pollution and allergic diseases exposure from nearby sources of ambient air pollution within elementary school zones in four Korean cities. *Environ. Sci. Pollut. Res.* **2013**, *20*, 4831–4846. [[CrossRef](#)] [[PubMed](#)]
12. Ramadour, M.; Burel, C.; Lanteaume, A.; Vervloet, D.; Charpin, D.; Brisse, F.; Dutau, H.; Charpin, D. Prevalence of asthma and rhinitis in relation to long-term exposure to gaseous air pollutants. *Allergy* **2000**, *55*, 1163–1169. [[CrossRef](#)] [[PubMed](#)]
13. Nordling, E.; Berglind, N.; Melén, E.; Emenius, G.; Hallberg, J.; Nyberg, F.; Pershagen, G.; Svartengren, M.; Wickman, M.; Bellander, T. Traffic-related air pollution and childhood respiratory symptoms, function and allergies. *Epidemiology* **2008**, *19*, 401–408. [[CrossRef](#)] [[PubMed](#)]
14. Wang, K.Y.; Chau, T.T. An association between air pollution and daily outpatient visits for respiratory disease in a heavy industry area. *PLoS ONE* **2013**, *8*, e75220. [[CrossRef](#)] [[PubMed](#)]
15. D'amato, G.; Cecchi, L.; D'amato, M.; Liccardi, G. Urban air pollution and climate change as environmental risk factors of respiratory allergy: An update. *J. Investig. Allergol. Clin. Immunol.* **2010**, *20*, 95–102. [[PubMed](#)]
16. Leung, T.F.; Wai-san Ko, F.; Wong, G.W. Roles of pollution in the prevalence and exacerbations of allergic diseases in Asia. *J. Allergy Clin. Immunol.* **2012**, *129*, 42–47. [[CrossRef](#)] [[PubMed](#)]
17. Hwang, B.F.; Jaakkola, J.J.; Lee, Y.L.; Lin, Y.C.; Guo, Y.L. Relation between air pollution and allergic rhinitis in Taiwanese schoolchildren. *Respir. Res.* **2006**, *7*, 23. [[CrossRef](#)] [[PubMed](#)]
18. Lee, Y.L.; Shaw, C.K.; Su, H.J.; Lai, J.S.; Ko, Y.C.; Huang, S.L.; Sung, F.C.; Guo, Y.L. Climate, traffic-related air pollutants and allergic rhinitis prevalence in middle-school children in Taiwan. *Eur. Respir. J.* **2003**, *21*, 964–970. [[CrossRef](#)] [[PubMed](#)]
19. Ng, T.P.; Tan, W.C. Epidemiology of allergic rhinitis and its associated risk factors in Singapore. *Int. J. Epidemiol.* **1994**, *23*, 553–558. [[CrossRef](#)] [[PubMed](#)]
20. Zhang, F.; Wang, W.; Lv, J.; Krafft, T.; Xu, J. Time-series studies on air pollution and daily outpatient visits for allergic rhinitis in Beijing, China. *Sci. Total Environ.* **2011**, *409*, 2486–2492. [[CrossRef](#)] [[PubMed](#)]

21. Liu, M.M.; Wang, D.; Zhao, Y.; Liu, Y.Q.; Huang, M.M.; Liu, Y.; Sun, J.; Ren, W.H.; Zhao, Y.D.; He, Q.C.; et al. Effects of outdoor and indoor air pollution on respiratory health of Chinese children from 50 kindergartens. *J. Epidemiol.* **2013**, *23*, 280–287. [[CrossRef](#)] [[PubMed](#)]
22. Lu, C.; Deng, Q.H.; Ou, C.Y.; Liu, W.W.; Sundell, J. Effects of ambient air pollution on allergic rhinitis among preschool children in Changsha, China. *Chin. Sci. Bull.* **2013**, *58*, 4252–4258. [[CrossRef](#)]
23. Peng, R.D.; Dominici, F.; Pastor-Barriuso, R.; Zeger, S.L.; Samet, J.M. Seasonal analyses of air pollution and mortality in 100 U.S. cities. *Am. J. Epidemiol.* **2005**, *161*, 585–594. [[CrossRef](#)] [[PubMed](#)]
24. Ito, K.; Mathes, R.; Ross, Z.; Nádas, A.; Thurston, G.; Matte, T. Fine particulate matter constituents associated with cardiovascular hospitalizations and mortality in New York City. *Environ. Health Perspect.* **2011**, *119*, 467. [[CrossRef](#)] [[PubMed](#)]
25. Zhang, F.; Krafft, T.; Ye, B.; Zhang, J.; Luo, H.; Li, M.; Zhang, Y.; Zhou, L.; Xu, J.; Wang, W. The lag effects and seasonal differences of air pollutants on allergic rhinitis in Beijing. *Sci. Total Environ.* **2013**, *442*, 172–176. [[CrossRef](#)] [[PubMed](#)]
26. Hajat, S.; Haines, A.; Atkinson, R.W.; Bremner, S.A.; Anderson, H.R.; Emberlin, J. Association between air pollution and daily consultations with general practitioners for allergic rhinitis in London, United Kingdom. *Am. J. Epidemiol.* **2001**, *153*, 704–714. [[CrossRef](#)] [[PubMed](#)]
27. Dominici, F.; McDermott, A.; Zeger, S.L.; Samet, J.M. On the use of generalized additive models in time-series studies of air pollution and health. *Am. J. Epidemiol.* **2002**, *156*, 193–203. [[CrossRef](#)] [[PubMed](#)]
28. Wood, S. *Generalized Additive Models: An Introduction with R*; CRC Press: Boca Raton, FL, USA, 2006.
29. Qian, Z.; He, Q.; Lin, H.M.; Kong, L.; Liao, D.; Dan, J.; Bentley, C.M.; Wang, B. Association of daily cause-specific mortality with ambient particle air pollution in Wuhan, China. *Environ. Res.* **2007**, *105*, 380–389. [[CrossRef](#)] [[PubMed](#)]
30. Welty, L.; Zeger, S.L. Are the acute effects of particulate matter on mortality in the National Morbidity, Mortality, and Air Pollution Study the result of inadequate control for weather and season? A sensitivity analysis using flexible distributed lag models. *Am. J. Epidemiol.* **2005**, *162*, 80–88. [[CrossRef](#)] [[PubMed](#)]
31. Gasparrini, A.; Armstrong, B.; Kenward, M.G. Distributed lag non-linear models. *Stat. Med.* **2010**, *29*, 2224–2234. [[CrossRef](#)] [[PubMed](#)]
32. Chan, C.K.; Yao, X. Air pollution in mega cities in China. *Atmos. Environ.* **2008**, *42*, 1–42. [[CrossRef](#)]
33. Zhang, F.; Krafft, T.; Zhang, D.; Xu, J.; Wang, W. The association between daily outpatient visits for allergic rhinitis and pollen levels in Beijing. *Sci. Total Environ.* **2012**, *417*, 39–44. [[CrossRef](#)] [[PubMed](#)]
34. Zhao, J.; Bai, J.; Shen, K.; Xiang, L.; Huang, S.; Chen, A.; Huang, Y.; Wang, J.; Ye, R. Self-reported prevalence of childhood allergic diseases in three cities of China: A multicenter study. *BMC Public Health* **2010**, *10*, 551. [[CrossRef](#)] [[PubMed](#)]
35. Zhang, F.Y.; Xu, J.; Wang, L.; Lu, J.M.; Li, Y.H.; Ni, Y.; Wang, W.Y.; Thomas, K. Air Quality, Patterns and Otolaryngology Health Effects of Air Pollutants in Beijing in 2013. *Aerosol Air Qual. Res.* **2016**, *16*, 1464–1472. [[CrossRef](#)]
36. Ma, Y.; Chen, R.; Pan, G.; Xu, X.; Song, W.; Chen, B.; Kan, H. Fine particulate air pollution and daily mortality in Shenyang, China. *Sci. Total Environ.* **2011**, *409*, 2473–2477. [[CrossRef](#)] [[PubMed](#)]
37. Kan, H.; Chen, B.; Hong, C. Health impact of outdoor air pollution in China: Current knowledge and future research needs. *Environ. Health Perspect.* **2009**, *117*, A187. [[CrossRef](#)] [[PubMed](#)]
38. Chen, W.; Tong, D.; Zhang, S.; Dan, M.; Zhang, X.; Zhao, H. Temporal variability of atmospheric particulate matter and chemical composition during a growing season at an agricultural site in northeastern China. *J. Environ. Sci.* **2015**, *38*, 133–141. [[CrossRef](#)] [[PubMed](#)]
39. Shang, Y.; Sun, Z.; Cao, J.; Wang, X.; Zhong, L.; Bi, X.; Li, H.; Liu, W.; Zhu, T.; Huang, W. Systematic review of Chinese studies of short-term exposure to air pollution and daily mortality. *Environ. Int.* **2013**, *54*, 100–111. [[CrossRef](#)] [[PubMed](#)]
40. Chen, T.M.; Gokhale, J.; Shofer, S.; Kuschner, W.G. Outdoor air pollution: Ozone health effects. *Am. J. Med. Sci.* **2007**, *333*, 244–248. [[CrossRef](#)]
41. De Marco, R.; Poli, A.; Ferrari, M.; Accordini, S.; Giammanco, G.; Bugiani, M.; Villani, S.; Ponzio, M.; Bono, R.; Carrozzi, L.; et al. The impact of climate and traffic-related NO<sub>2</sub> on the prevalence of asthma and allergic rhinitis in Italy. *Clin. Experiment. Allergy* **2002**, *32*, 1405–1412. [[CrossRef](#)]

42. Villeneuve, P.; Doiron, M.S.; Stieb, D.; Dales, R.; Burnett, R.T.; Dugandzic, R. Is outdoor air pollution associated with physician visits for allergic rhinitis among the elderly in Toronto, Canada? *Allergy* **2006**, *61*, 750–758. [[CrossRef](#)] [[PubMed](#)]
43. Pope, C.A.; Burnett, R.T.; Krewski, D.; Jerrett, M.; Shi, Y.; Calle, E.E.; Thun, M.J. Cardiovascular mortality and exposure to airborne fine particulate matter and cigarette smoke shape of the exposure-response relationship. *Circulation* **2009**, *120*, 941–948. [[CrossRef](#)] [[PubMed](#)]
44. Wang, H.; Devalia, L.; Duddle, M.; Hamilton, S.A.; Davies, R.J. Effect of six-hour exposure to nitrogen dioxide on early-phase nasal response to allergen challenge in patients with a history of seasonal allergic rhinitis. *J. Allergy Clin. Immunol.* **1995**, *96*, 669–676. [[CrossRef](#)]
45. Glück, U.; Gebbers, O. Epithelial changes in seasonal allergic rhinitis throughout the year: Evidence of coexistent air pollution and local secretory IgA deficiency? *ORL* **2000**, *62*, 68–75. [[CrossRef](#)] [[PubMed](#)]
46. Peden, D.B.; Setzer, R.R.W.; Devlin, R.B. Ozone exposure has both a priming effect on allergen-induced responses and an intrinsic inflammatory action in the nasal airways of perennially allergic asthmatics. *Am. J. Respir. Crit. Care Med.* **1995**, *151*, 1336–1345. [[CrossRef](#)] [[PubMed](#)]
47. Hwang, B.F.; Jaakkola, J.J. Ozone and other air pollutants and the risk of oral clefts. *Environ. Health Perspect.* **2008**, *116*, 1411. [[CrossRef](#)] [[PubMed](#)]
48. Ichinose, T.; Hiyoshi, K.; Yoshida, S.; Takano, H.; Inoue, K.; Nishikawa, M.; Mori, I.; Kawazato, H.; Yasuda, A.; Shibamoto, T. Asian sand dust aggravates allergic rhinitis in guinea pigs induced by apanese cedar pollen. *Inhal. Toxicol.* **2009**, *21*, 985–993. [[CrossRef](#)] [[PubMed](#)]
49. Fukuoka, A.; Matsushita, K.; Morikawa, T.; Takano, H.; Yoshimoto, T. Diesel exhaust particles exacerbate allergic rhinitis in mice by disrupting the nasal epithelial barrier. *Clin. Exp. Allergy* **2016**, *46*, 142–152. [[CrossRef](#)] [[PubMed](#)]
50. Mösges, R.; Klimek, L. Today's allergic rhinitis patients are different: New factors that may play a role. *Allergy* **2007**, *62*, 969–975. [[CrossRef](#)] [[PubMed](#)]



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).