

Table S1. Indoor air questionnaire (Finnish Institute of Occupational Health® 2006–2008, version 2.0) results from May 2016 and January 2017.

	Reference data	Building Section 1		<i>p</i> -Value	Building Section 2		<i>p</i> -Value
	[1-3]	2016	2017		2016	2017	
Background information							
Number of answers		15	16		16	17	
Response rate (%)	71	79	80		84	85	
Females (%)	21	87	88		94	94	
Daily smokers (%)		13	6		0	0	
Average age (years)		41	42		41	38	
Average employment in this work place (years)	5	5			4	4	
Work environment (%)							
		%	%				
Draught	22	7	44	0.04*	13	47	0.06*
Room temperature too high	17	0	0	-	0	6	1.00*
Varying temperature	16	20	31	0.69*	19	27	0.69*
Room temperature too low	13	27	56	0.10	19	53	0.04
Stuffy air	34	53	38	0.38	80	71	0.69*
Dry air	35	13	44	0.11*	14	41	0.13*
Insufficient ventilation	32	47	31	0.38	75	59	0.33
Smell of mold	9	7	0	0.48*	7	6	1.00*
Unpleasant odour	17	40	19	0.25*	38	24	0.47*
Environmental tobacco smoke	4	0	0	-	13	0	0.23*
Noise	17	47	56	0.59	19	50	0.06
Dim light or reflections	14	7	13	1.00*	6	0	0.49*
Dust or dirt	25	27	25	1.00*	25	35	0.71*
Work regarded as interesting and stimulating (%)							
Often	75	73	88	0.39*	88	82	1.00*
Sometimes	20	27	13		13	18	
Seldom or never	4	0	0		0	0	
Too much work to do (%)							
Often	20	0	13	0.08*	13	18	0.70*
Sometimes	59	40	63		56	65	
Seldom or never	21	60	25		31	18	
Opportunity to influence work conditions (%)							
Often	35	27	25	0.51*	25	24	0.58*
Sometimes	44	60	75		63	47	
Seldom or never	21	13	0		13	29	
Fellow workers help with problems in the work (%)							
Often	72	87	88	1.00*	88	76	0.05*
Sometimes	22	13	13		0	24	
Seldom or never	6	0	0		13	0	
Allergic diseases (%)							
Asthma	8	0	0	-	19	18	1.00*
Hay fever	38	67	56	0.55*	50	41	0.61
Atopic eczema	28	40	19	0.25*	13	12	1.00*
Stress (%)							
Very much	10	7	0	0.45*	13	24	0.74*
Some	28	27	47		50	35	
None/only a little	63	67	53		38	41	
Symptoms (%)							
Fatigue	16	7	19	0.60*	19	29	0.69*
Heavy-headedness	9	20	13	0.65*	6	35	0.09*
Headache	7	13	6	0.60*	19	29	0.69*
Difficulty concentrating	3	0	6	1.00*	6	18	0.60*
Eye irritation	17	27	31	1.00*	20	41	0.27*
Irritated, stuffy, or running nose	20	13	25	0.65*	19	35	0.44*
Hoarse/dry throat	14	13	38	0.22*	31	35	0.81
Cough	5	0	13	0.48*	6	18	0.60*
Cough disturbing sleep	1	0	6	1.00*	0	0	-
Dry or flushed facial skin	11	7	25	0.33*	13	25	0.65*
Hands: dry, itching, red skin	15	0	19	0.23*	7	24	0.34*
Shortness of breath	3	7	0	0.48*	0	6	1.00*
Wheezing	1	7	0	0.48*	0	6	1.00*
Fever or chills	2	0	0	-	7	6	1.00*
Joint pain	3	0	0	-	0	0	-
Muscular pain	4	0	0	-	0	0	-
Other		0	0	-	9	12	0.83*

Statistically significant changes at a 10% confidence interval ($p < 0.1$) are bolded. The p-values marked with * were determined by the Fisher's exact test (SPSS). Reference data are based on analysis of the comprehensive questionnaire data collected by FIOH.